

A Facile Synthesis of α -*N*-Ribosyl-Asparagine and α -*N*-Ribosyl-Glutamine Building Blocks.

Gaetano Speciale, Anna Bernardi, Filippo Nisic*

Universita' degli Studi di Milano, Dipartimento di Chimica, via Golgi 19, 20133 Milano, Italy

Abstract

Adenosine diphosphate ribosylation (ADP-ribosylation) is a widely occurring post-translational modification of proteins at nucleophilic side chain of amino acid residues. Elucidation of ADP-ribosylation events would benefit greatly from the availability of well-defined ADP-ribosylated peptides and analogues thereof. In this paper we present a novel approach to the chemical synthesis of ribosylated amino acid building blocks using traceless Staudinger ligation. We describe an efficient and stereoselective synthesis of α -*N*-ribosyl-asparagine (α -*N*-ribosyl-Asn) and α -*N*-ribosyl-glutamine (α -*N*-ribosyl-Gln) building blocks starting from 5-*tert*-butyldiphenylsilyl- β -D-ribofuranosyl azide. The *N*-glycosyl aminoacids are produced in good yields as pure α -anomers, suitably protected for peptide synthesis.


Keywords: ADP-ribosylation; glycoconjugates; ribofuranosyl aminoacids; Staudinger ligation; stereoselective synthesis.

Introduction

Glycosylated peptides have a fundamental role in biological systems: in fact, more than half of all proteins carry carbohydrate moieties, generating different glycoforms whose exact composition often controls protein function and distribution in biological systems [1]. The details of glycan regulation of protein activity and stability are still under intense scrutiny and the synthesis of well-defined glycopeptides is therefore an important target [2], which still presents many challenges to organic chemistry. One of the processes that is currently being elucidated is adenosine diphosphate ribosylation (ADP-ribosylation), a wide-occurring post-translational modification effected by enzymes that transfer ADP-ribose from NAD⁺ to Asn, Glu, Asp, Arg or Cys residues of proteins, so altering their function [3]. An important contribution for the clarification of the role of ADP-ribosylation events came in 2010, when Filippov and coworkers [4] reported the synthesis of ribosylated oligopeptides using α -*N*-ribosyl-asparagine (α -*N*-ribosyl-Asn) and α -*N*-ribosyl-glutamine (α -*N*-ribosyl-Gln) building blocks. Despite the relevance of this work, the approach

* Corresponding author. Tel.: +39-02-503-14092; e-mail: filippo.nisic@unimi.it


suffers from a poorly selective synthesis of the required ribosylated building blocks **4 α** and **5 α** (Scheme 1), which were prepared by PtO₂ reduction of azide **3**, followed by EDC-mediated coupling of the resulting mixture of epimeric amines with Z-Asp-OBn or Z-Glu-OBn, respectively. The *N*-ribosyl-aminoacids were obtained as a 3:1 α : β anomeric mixtures, from which the desired α -anomers were to be chromatographically isolated. Acetyl transfer to the anomeric nitrogen from the 2-*O*-acetate also occurs during the reaction, further reducing the process yields, particularly for the asparagine derivative **4 α** .



Scheme 1. Filippov's synthesis of ribosylated amino acids (from *ref 4*).

We have recently developed a protocol for the stereoselective synthesis of either α - or β -*N*-glycofuranosyl amides using the Staudinger traceless ligation [5]. In this reaction, a starting anomeric glycosyl azide is partially reduced by an appropriate phosphine and then intramolecularly acylated in order to afford the amide (Scheme 2) [6]. We have shown that for furanosyl azides the outcome of the process is controlled by the configuration and the protection state of the contiguous hydroxyl group, so that 1,2-*cis* amides (α -, in the *ribo* series) are obtained when this group is unprotected and 1,2-*trans* amides (β -, in the *ribo* series) when it is acetylated. For unprotected furanoses, the anomeric configuration of the product amides appears to be dictated by O-P coordination, as supported by ³¹P-NMR studies [5b].

As an application of this methodology, we here report the facile stereoselective synthesis of **4 α** and **5 α** that could be obtained in good yields and with full stereoselectivity with a single synthetic operation starting from β -5-*O*-*tert*-butyldiphenylsilyl-ribosylazide **2**.



Scheme 2. Traceless Staudinger ligation of furanosyl azides with functionalized phosphines (from *ref 5a*).

Experimental Section

Solvents were dried by standard procedures: dichloromethane and *N,N*-diisopropylethylamine were dried over calcium hydride; *N,N*-dimethylacetamide (DMA), 1,3-dimethyltetrahydro-2(1*H*)pyrimidinone (DMPU), chloroform and pyridine were dried over activated molecular sieves. Reactions requiring anhydrous conditions were performed under nitrogen. ^1H , ^{13}C and ^{31}P -NMR spectra were recorded at 400 MHz on a Bruker AVANCE-400 instrument. Chemical shifts (δ) for ^1H and ^{13}C spectra are expressed in ppm relative to internal Me_4Si as standard. Chemical shifts (δ) for ^{31}P are expressed in ppm relative to internal H_3PO_4 as standard. Signals were abbreviated as s, singlet; bs, broad singlet; d, doublet; t, triplet; q, quartet; m, multiplet. Mass spectra were obtained with a Bruker ion-trap Esquire 3000 apparatus (ESI ionization) or FT-ICR Mass Spectrometer APEX II & Xmass software (Bruker Daltonics) - 4.7 Magnet and Autospec Fission Spectrometer (FAB ionization). Thin layer chromatography (TLC) was carried out with pre-coated Merck F₂₅₄ silica gel plates. Flash chromatography (FC) was carried out with Macherey-Nagel silica gel 60 (230-400 mesh).

General procedure for the synthesis of 2-(diphenylphosphanyl)-4-fluorophenyl esters **6**

A solution of the *o*-diphenylphosphinophenol **7** (1 equiv.), the commercially available *N*-carbobenzyloxy-L-asparagine acid 1-benzyl ester (*Z*-Asp-OBn, 1.2 equiv.) and *N,N*-dimethylaminopyridine (0.1 equiv.) in dry CH_2Cl_2 (0.1 M) were added, at room temperature and under nitrogen, to a suspension of *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDC, 1.4 equiv.) and dry *N,N*-diisopropylethylamine (1.4 equiv.) in dry CH_2Cl_2 . The mixture was

stirred at RT for 2 h, monitoring by TLC (60:40 hexane/AcOEt). The reaction mixture was diluted with CH₂Cl₂ and washed with 5 % aqueous HCl and water: the organic layer was dried over Na₂SO₄ and concentrated. The crude product obtained was purified by flash chromatography, as indicated in each case.

1-Benzyl 5-[2-(Diphenylphosphanyl)-4-fluorophenyl] N-(Benzyloxy-carbonyl)-L-aspartate (6a)

The crude product was purified by flash chromatography (hexane/AcOEt 60:40). yield = 85 %.

¹H-NMR (400 MHz, CDCl₃, 25°C): δ = 7.37-7.23 (m, 20H, Ph), 7.02-6.84 (m, 2H, H-2, H-3), 6.49 (m, 1H, H-1), 5.69 (d, *J*_{NH-CH} = 8.6 Hz, 1H, NH), 5.17 (s, 2H, CH₂O), 5.13 (s, 2H, CH₂O), 4.62 (m, *J*_{CH-CH₂} = 4.4 Hz, 1H, CH), 3.00 (dt, *J* = 4.9 Hz, *J* = 17.4 Hz, 1H, H_a, CH₂), 2.67 (dd, *J* = 4.3 Hz, *J* = 17.4 Hz, 1H, H_b, CH₂). ¹³C-NMR (100 MHz, CDCl₃, 25°C): δ = 170.4, 169.2, 161.9, 159.4 (CO_(Cbz)), 134.6, 134.0, 132.0, 131.7, 129.0, 128.9, 129.0, 128.7, 128.6, 128.4, 128.3, 128.2 (C_{Ar}), 124.0, 116.9, 116.6, 67.8 (CH₂O), 67.3 (CH₂O), 50.4 (CH), 36.5 (CH₂). ³¹P-NMR (161 MHz, CDCl₃; 25°C): δ = - 14.3 ppm. FAB-MS: *m/z* 636 (M+1).

1-Benzyl 5-[2-(Diphenylphosphanyl)-4-fluorophenyl] N-(Benzyloxycarbonyl)-L-glutamate (6b)

The crude product was purified by flash chromatography (hexane/AcOEt 70:30). yield = 85 %.

¹H-NMR (400 MHz, CDCl₃, 25°C): δ = 7.37-7.24 (m, 20 H, Ph), 7.05 (m, 1H, H-2), 6.99 (m, 1H, H-3), 6.45 (m, 1H, H-1), 5.37 (d, *J*_{NH-CH} = 8.1 Hz, 1H, NH), 5.16 (s, 2H, CH₂-O), 5.07 (s, 2H, CH₂-O), 4.38 (m, 1H, CH), 2.26 (m, 2H, CH₂-CO), 2.05 (m, 1H, H_a, CH₂-CH), 1.80 (m, 1H, H_b, CH₂-CH). ¹³C-NMR (100 MHz, CDCl₃, 25°C): δ = 170.7, 161.7, 159.3, 156.1 (CO_(Cbz)), 134.4, 134.3, 134.2, 134.0, 129.1, 129.0, 128.8, 128.5, 128.4 (C_{Ar}), 124.2, 124.1, 120.0, 119.8, 116.8, 116.5, 67.6 (CH₂-O), 67.4 (CH₂-O), 53.5 (CH), 30.1 (CH₂-CO), 27.4 (CH₂). ³¹P-NMR (161 MHz, CDCl₃, 25°C): δ = - 14.31 ppm. FAB-MS: *m/z* 650 (M+1).

General procedure for the Staudinger Traceless Ligation of ribofuranosyl azides.

Phosphine **6a** or **6b** (2 equiv.) was added, at room temperature, to a 0.1 M solution of ribofuranosyl azide **2** (1 equiv.) in 98:2 *N,N*-dimethylacetamide and DMPU. The solution was stirred for 20 h at 70 °C, then 10% water was added and the mixture was stirred for an additional 2 h at the same temperature. The solvent was evaporated under reduced pressure, and the residue was purified by flash chromatography as indicated in each case.

General procedure for the acetylation of the ribofuranosyl amides.

Ac₂O (6 eq), pyridine (6 equiv.) and a catalytic amount of *N,N*-dimethylaminopyridine were added, at room temperature, to a solution of substrate (1 equiv.) in dry CH₂Cl₂ (0.1 M). The solution was stirred for 24 h and then was concentrated in vacuo. The residue was dissolved in AcOEt and washed with aqueous 5 % HCl, aqueous 5 % NaHCO₃ and water. The organic layer was dried over Na₂SO₄ and concentrated. The crude was then purified by flash chromatography (hexane:AcOEt, 80:20→50:50).

Purification afforded products with spectroscopic data coherent with the fully characterized compounds published by Filippov and coworkers ⁴.

***N*^δ-(5-*tert*-Butyldiphenylsilyl- α -D-ribofuranosyl)-*N* ^{α} -benzyloxycarbonyl asparagine benzyl ester (12 α)**

The crude product was purified by flash chromatography (CHCl₃/MeOH 95:5). yield = 60 %.

¹H-NMR (400 MHz, CD₃OD, 25°C): δ = 7.71-7.64, 7.42-7.24 (3 x m, 20H, Arom.), 5.73 (d, $J_{1,2}$ = 4.8 Hz, 1H, H-1), 5.15, 5.07 (2 x s, 2H, CH₂Bn, CH₂Cbz), 4.65 (m, 1H, CH α -Asn), 4.25 (t, J = 5 Hz, 1H, H-3), 4.15 (t, J = 4.8 Hz, 1H, H-2), 3.97 (m, 1H, H-4), 3.72 (dd, $J_{4,5}$ = 3.2 Hz, $J_{5,5'}$ = 11.3 Hz, 2H, H-5 H-5'), 2.84 (m, 2H, CH₂ β -Asn), 1.04 (s, 9H, t-Bu). ¹³C-NMR (100 MHz, CD₃OD, 25°C): 172.9, 172.4 (CO α -Asn, CO γ -Asn), 158.6 (CO Cbz), 138.2 (Cq Arom.), 137.3 (Cq Arom.), 136.9, 136.8 (Arom.), 134.6, 134.4 (Cq Arom.), 131.1, 131.0, 129.7, 129.6, 129.5, 129.5, 129.3, 129.3, 129.0, 129.0 – 128.8, 128.4, 128.1 (Arom.), 84.7 (C-4), 81.9 (C-1), 72.8 (C-3), 72.1 (C-2), 68.4, 67.9 (CH₂ Bn, CH₂ Cbz), 65.3 (C-5), 52.4 (CH, α -Asn), 38.9 (CH₂, β -Asn), 27.4 (CH₃, t-Bu), 20.2 (Cq, t-Bu). FT-ICR (ESI) calcd. for C₄₀H₄₆N₂O₉Si [M+Na]⁺ 749.29726; found 749.29721.

***N*^δ-(5-*tert*-Butyldiphenylsilyl- α -D-ribofuranosyl)-*N* ^{α} -benzyloxycarbonyl glutamine benzyl ester (13 α)**

The crude product was purified by flash chromatography (CHCl₃/MeOH 95:5). yield = 76 %.

¹H-NMR (400 MHz, CD₃OD, 25°C): δ = 7.71-7.64, 7.43-7.23 (3 x m, 20H, Arom.), 5.72 (d, $J_{1,2}$ = 4.6 Hz, 1H, H-1), 5.15, 5.06 (2 x s, 2H, CH₂Bn, CH₂Cbz), 4.25 (m, 1H, CH α -Gln), 4.23 (t, J = 4.9 Hz, 1H, H-3), 4.15 (t, J = 4.8 Hz, 1H, H-2), 3.95 (m, 1H, H-4), 3.70 (dd, $J_{4,5}$ = 3.2 Hz, $J_{5,5'}$ = 10.2 Hz, 2H, H-5 H-5'), 2.35 (m, 2H, CH₂ γ -Asn), 2.13 (m, 1H, CH₂ β -Asn), 1.95 (m, 1H, CH₂ β -Asn), 1.02 (s, 9H, t-Bu). ¹³C-NMR (100 MHz, CD₃OD, 25°C): 136.7, 136.6 (Arom.), 134.6, 134.4 (Cq Arom.), 130.8, 130.8, 129.7, 129.5, 129.4, 129.1, 128.9, 128.8, 128.7 (Arom.), 84.5 (C-4), 81.6 (C-1), 72.7 (C-3), 71.9 (C-2), 67.9, 67.6 (CH₂ Bn, CH₂ Cbz), 65.0 (C-5), 55.1 (CH α -Gln), 33.3 (CH₂

β -Gln), 28.0 (CH2 γ -Gln), 27.3 (t-Bu), 22.9 (Cq t-Bu). FT-ICR (ESI) calcd. for $C_{41}H_{48}N_2O_9Si$ $[M+Na]^+$ 763.31291; found 763.31298.


Results and Discussion

Previous research in our laboratories had shown that transfer of aminoacid chains in Staudinger ligations of glycosyl azides was particularly effective using fluorinated phosphines **6** [7]. Thus we envisaged that phosphines **6a** and **6b** (Scheme 3), functionalized with Z-Asp-OBn and Z-Glu-OBn acyl chains, respectively, would be the reagent of choice to prepare the target α -N-ribosyl-Asn and α -N-ribosyl-Gln building blocks. Phosphines **6a** and **6b** were synthesized in excellent yields by EDC-mediated acylation of 2-diphenylphosphanyl-5-fluoro-phenol **7** [8] and purified by filtration on short pads of silica.



Scheme 3. Synthesis of functionalized phosphines **6a** and **6b**.


An early trial of ligation was conducted using the unprotected β -D-ribofuranosyl azide **1** [9] with phosphines **6a** and **6b** for 20 h at 70 °C in a 98:2 DMA:DMPU solution, followed by water quenching (Scheme 4). In both ligation reactions 45 % of a 3:2 mixture of α -ribofuranosyl amide (**8 α** or **9 α**) and β -ribopyranosyl amide (**10** or **11**) was obtained. The isomer distribution was determined by $^1\text{H-MNR}$ analysis of the crude reaction mixtures: the α -ribofuranosyl amide (**8 α** or **9 α**) was characterized by the presence of the H1 proton signal at 5.65 ppm and by the high chemical shift of C4 (84.3 ppm), which is diagnostic for the furanose form. The structures of **10** and **11** were assigned on the basis of the vicinal coupling constant of the anomeric protons ($J_{1,2} = 8$ Hz, H1 = 5.12 ppm) and of the low chemical shift value of C4 (68.8 ppm), that confirms the pyranose structure. The β -glycopyranosyl isomers **10** and **11** obtained in these reactions must derive from ring-expansion occurring after a ring-opening process.



a) 70°C, 20h, DMA/DMPU 98:2; b) H₂O, 2 h, 70°C

Scheme 4. Ligation of **1** with phosphines **6a** and **6b**.

To avoid ring expansion, the 5-hydroxy group of **1** was protected as a *tert*-butyldiphenylsilyl ether and the 5-*tert*-butyldiphenylsilyl- β -D-ribofuranosyl azide **2** was synthesized in good yields as described by Filippov [4]. Ligation of **2** with **6a** or **6b** was carried out as above (20 h at 70 °C in a 98:2 DMA:DMPU) (Scheme 5). In both cases, 400 MHz ¹H-NMR analysis of the crude showed only one signal in the furanose anomeric region, implying that no pyranose was formed, as expected, and that the α/β ratio of the resulting ribofuranosyl amides **12 α** and **13 α** was $\geq 99:1$. The two compounds were isolated by flash chromatography, which afforded the desired building blocks in 60% and 76% yields, respectively, as single α isomers. After acetylation (Ac₂O, cat. DMAP, CH₂Cl₂, quant.), **4 α** and **5 α** were obtained. Their spectroscopic data were fully consistent with the already published characterization [4].



a) 70°C, 20h, DMA/DMPU 98:2; b) H₂O, 2 h, 70°C; c) Ac₂O, DMAP cat., CH₂Cl₂.

Scheme 5. Synthesis of ribofuranosyl amides **4 α** and **5 α** .

Conclusion

In conclusion, we have described an improved synthesis of the building blocks **4** α and **5** α : application of the Staudinger Traceless Ligation protocol allowed to obtain the ribosylated amino acids from **2** in good yields and with excellent selectivity for the α anomer. This represents a major advantage over pre-existing synthesis of these glycosyl aminoacids [4, 10] and should stimulate further research in the field of ADP-ribosylation.

Acknowledgements. This work was supported by the University of Milan (PUR) and Ministero dell'Universita' e della Ricerca (PRIN). Exact mass were obtained from CIGA.

References

1. a) Varki, A. *Glycobiology* **1993**, *3*, 97-130. b) Dwek, R. A. *Chem. Rev.* **1996**, *96*, 683-720. c) Reuter, G.; Gabius, H. *J. Cell. Mol. Life Sci.* **1999**, *55*, 368-422. d) Gamblin, D. P.; Scanlan, E. M.; David, B. G. *Chem. Rev.* **2009**, *109*, 131-163.
2. a) Marcaurelle, L. A.; Bertozzi, C. R. *Glycobiology* **2002**, *12* (6), 69R-77R. b) Warren, J. D.; Miller, J. S., Keding, S. J.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2004**, *126*, 6576.
3. Corda, D.; Di Girolamo, M. *EMBO J.* **2003**, *22*, 1953-1958.
4. van der Heden van Noort, G. J.; van der Horst, M. G.; Overkleeft, H. S.; van der Marel, G. A.; Filippov, D. V. *J. Am. Chem. Soc.* **2010**, *132*, 5236-5240.
5. a) Nisic, F.; Bernardi, A. *Carbohydr. Res.* **2011**, *346*, 465-471; b) Nisic, F.; Speciale, G.; Bernardi, A. *Chem. Eur. J.* **2012**, *18*, 6895-6906.
6. Bianchi, A.; Bernardi, A. *J. Org. Chem.* **2006**, *71*, 4565-4577.
7. Nisic, F.; Andreini, M.; Bernardi, A. *Eur. J. Org. Chem.* **2009**, 5744-5751.
8. Heinicke, J.; Köhler, M.; Peulecke, N.; He, M.; Kindermann, M. K.; Keim, W.; Fink, G. *Chem. Eur. J.* **2003**, *9*, 6093-6107.
9. Stimac, A.; Kobe, J. *Carbohydr. Res.* **1992**, *232*, 359-365.
10. Bonache, M. A.; Nuti, F.; Le Chevalier Isaad A.; Real-Fernandez F.; Chelli M.; Rovero, P.; Papini A. M.; *Tetrahedron Lett.* **2009**, *50*, 4151-4153.