

# Synthesis of new molecules based on 1-pyrindane ring as a new strategy in the neurodegenerative diseases treatment

Cidália Pereira, Fabio Rizzo-Aguiar, José Enrique Rodriguez-Borges, Xerardo Garcia-Mera

## Common synthetic pathway for functionalization

#### First 1-pyrindane derivatives achieved





# Synthesis of new molecules based on 1-pyrindane ring as a new strategy in the neurodegenerative diseases treatment

Cidália Pereira, Fabio Rizzo-Aguiar, José Enrique Rodriguez-Borges, Xerardo Garcia-Mera

## Most recent approaches for bis-functionalization

$$R_{1}$$
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{2}$ 
 $R_{4}$ 
 $R_{1}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{1}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{1}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{4}$ 
 $R_{5}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{6}$ 
 $R_{1}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{6}$ 
 $R_{1}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{7}$ 
 $R_{1}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{7}$ 
 $R_{1}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{7}$ 
 $R_{1}$ 
 $R_{2}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{5}$ 
 $R_{7}$ 
 $R_{1}$ 
 $R_{2}$ 
 $R_{3}$ 
 $R_{4}$ 
 $R_{5}$ 
 $R_{5$ 

