

Bioavailability of Rapeseed Oil Fortified with Ethyl Sinapate [†]

Dobrochna Rabiej-Kozioł ^{*}, Marek P. Krzemiński, Iwona Wiśniewska and Aleksandra Szydłowska-Czeraniak

Faculty of Chemistry, Nicolaus Copernicus University in Toruń, Gagarina 7, 87-100 Toruń, Poland; mkrzem@umk.pl (M.P.K.); 291003@stud.umk.pl (I.W.); olasz@umk.pl (A.S.C.)

^{*} Correspondence: d.rabiej@umk.pl[†] Presented at the 25th International Electronic Conference on Synthetic Organic Chemistry, 15–30 November 2021; Available online: <https://ecsoc-25.sciforum.net/>.

Abstract: In recent years, phenolic acid esters have been proposed as valuable additives in the food and cosmetic industries. Therefore, ethyl sinapate (ESA) was synthesized by an enzymatic approach. Hence, the bioaccessibility of antioxidant compounds after in vitro digestion process of rapeseed oil fortified with ESA was evaluated. Antioxidant activity (AA) of oil samples before and after digestion was analyzed by three spectrophotometric methods. The addition of synthesized ESA in the concentration of 0.02 and 0.5% caused an increase in AA of rapeseed oils from two to ten times. Therefore, the obtained phenolipid can be used as an efficient antioxidant in the oil industry.

Keywords: bioavailability; rapeseed oil; ethyl sinapate; antioxidant activity

1. Introduction

Rapeseed oil is the most popular vegetable oil in Poland. It is a valuable source of bioactive compounds such as polyphenols, phospholipids, and sterols. Additionally, it contains a high amount of essential fatty acids. Unfortunately, oils containing a high amount of unsaturated acids are easily oxidized. Therefore, it is desirable to protect the oil from the oxidation process [1]. However, recent studies have demonstrated the toxic effect of commercially available, artificial phenolic antioxidants, e.g., butylated hydroxyanisole (BHA) and butylated hydroxytoluene (BHT), and especially their metabolites [3]. This fact prompts the search for new compounds dedicated to the fat industry. Unfortunately, naturally phenolic antioxidants have hydrophilic properties. Hence, phenolic acids are esterified to obtain their amphiphilic derivatives. The research focused on phenolic acids because they are widespread in the plant world and have many health beneficial properties such as antioxidant, chelating, free radical scavenging, antiallergic, anti-inflammatory, antimicrobial, antiviral, anticarcinogenic [2,4–5]. Lyophilization of phenolic acids was carried out in chemical, enzymatic and chemo-enzymatic ways [2]. However, enzymatic synthesis offers milder reaction conditions. Additionally, it is known that phenolic acid alkyl esters possess antioxidant properties and enhanced AA of food products [2,4–5].

To the best of our knowledge, there is no data on the influence of ESA on AA of the enriched refined rapeseed oil after in vitro gastrointestinal digestion process. In general, antioxidants should be bioavailable and bioaccessible to be biologically active. At the same time, their bioavailability depends on the release of molecules from the food matrix during the digestive process [6,7].

Given the above, this study aimed at the enzymatic synthesis of ESA. Moreover, in order to increase the nutritional value of rapeseed oil and to valorize ESA, in the present study, the digestive stability of phenolic compounds from rapeseed oils enriched with ESA at two concentration levels was investigated and changes in their AA using the standardized static in vitro digestion model was monitored.

Citation: Rabiej-Kozioł, D.; Krzemiński, M.P.; Wiśniewska, I.; Szydłowska-Czeraniak, A. Bioavailability of Rapeseed Oil Fortified with Ethyl Sinapate. *Chem. Proc.* **2021**, *3*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Julio A. Seijas

Published: 15 November 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

2. Materials and Methods

2.1. Reagents

All reagents, reactants, and solvents were purchased from Merck (Warszawa, Poland). Silica gel (pore size 60Å, 230–400 mesh, Kieselgel, Macherey-Nagel, Dueren, Germany) was purchased from Alchem, (Toruń, Poland). Biocatalyst used in this work, Novozym 435 (*Candida Antarctica*) lipase B immobilized on a macroporous acrylic resin, were a generous gift from Novozymes (Madrid, Spain).

2.2. Samples

The refined rapeseed oil in the original packaging (polyethylene terephthalate (PET) bottle) was kindly provided by a local vegetable oil factory.

2.3. NMR Analysis

The structure of the ESA was confirmed by nuclear magnetic resonance (NMR) spectroscopy. ¹H NMR spectrum was recorded at Bruker Avance III 400 MHz spectrometer (Bruker Corporation, Karlsruhe, Germany) at 298 ± 1K. The sample was dissolved in CDCl₃ containing TMS as an internal standard. Chemical shifts were recorded in δ values in parts per million (ppm), and coupling constants (J) were reported in hertz (Hz).

2.4. Enzymatic Synthesis

Lipase-catalyzed esterification of sinapic acid with ethanol was conducted in a 20 ml vial equipped with a tight plastic cap. Sinapic acid (5 mM) with 10 mL of ethanol (dried over 3Å molecular sieves) was mixed. After dissolving the sinapic acid, the reaction was started by the addition of 60 mg/mL of the biocatalyst. Molecular sieves (3Å, 40 mg/mL) were also used to remove water formed as a by-product. The reaction was carried out in an orbital shaker at 120 rpm at 65 °C for five days (Incu-Shaker™ Mini Shaking Incubator-Benchmark Scientific; NY, USA). The biocatalyst and molecular sieves were filtered and washed on the Buchner funnel. The ethanol was evaporated under vacuum using a rotary evaporator (Laborota 4003, Heidolph Instruments, Schwabach, Germany). ESA was purified on silica gel using dichloromethane:ethyl acetate (90:10) as an eluent. The synthesized ESA [ethyl (E)-3-(4-hydroxy-3,5-dimethoxyphenyl)-propenoate] had the following characteristics: light yellow solid, ¹H NMR (CDCl₃) δ: 0.90 (t, J = 7.0 Hz, 3H), 3.95 (s, 6H), 4.19 (t, J = 6.8 Hz, 2H), 6.31 (d, J = 15.9 Hz, 1H), 6.82 (s, 2H), 7.69 (d, J = 15.9 Hz, 1H).

2.5. Addition of the ESA to Refined Rapeseed Oil

The refined rapeseed oil was fortified with the ESA at two different concentrations: 0.02% and 0.5%. The ESA was weighed, respectively, 0.01 and 0.25 g, added to 50.0 g of oils and transferred into Erlenmeyer flasks. Samples were placed in an ultrasonic cleaner bath (Sono Swiss, SW 6H, Labo Plus, Warszawa, Poland) with ultrasound input power of 180 kW for 5 min to completely dissolve of ESA.

2.6. In Vitro Digestion

The in vitro digestion process was carried out according to the procedure described by Seiquer et al. [6]. Before the digestion process, 1 g of oil sample and 9 mL of distilled water were placed in an ultrasonic cleaner bath (Sono Swiss, SW 6H, Labo Plus, Warszawa, Poland) with ultrasound input power of 180 kW for 5 min. The first step was a digestive process in the stomach (gastric digestion). An initial pH was adjusted to 2 with 1M HCl, then pepsin/HCl solution was added to each oil sample and put on an orbital shaker (Incu-Shaker™ Mini Shaking Incubator-Benchmark Scientific; NY, USA) at 37 °C at 110 rpm for 2 h. After this time, the pH was adjusted to 6 by the dropwise addition of 1 M NaHCO₃. Then 2.5 mL of pancreatic enzyme solution was added to simulate the next step, duodenal digestion. The pH was adjusted to 7.5 with 1 M NaHCO₃. The solutions

were again incubated for 2 h at 37 °C at rpm 110. After the digestion process, the enzymes were inactivated by heating for 4 min at 100 °C. The solutions were frozen for 15 min and then centrifuged also for 15 min to separate fractions. For each test, the top oil layer formed was weighed and processed for methanol extracts.

2.7. Antioxidant Activity

2.7.1. Sample Preparation

The methanolic extracts of the studied oils were prepared according to the procedure described in our previous work [8]. Briefly, 2.00 g of oil was weighed into test tubes and extracted with 5 mL of methanol for 30 min using an orbital shaker (SHKA25081 CE, Labo Plus, Warszawa, Poland). Then, extracts were frozen to separate oil from methanol (−20 °C, 30 min) and transferred quantitatively into glass bottles. Each oil sample was extracted in triplicate. Extracts were stored in a refrigerator to carried out AA analyzes.

2.7.2. Analytical Methods

The AA of refined rapeseed oil without and with ESA, before and after each step of the digestion process was determined by three spectrophotometric methods: 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS), 2,2-diphenyl-1-picrylhydrazyl (DPPH), and Folin–Ciocalteu (FC) described in our previous work [8]. The UV-Vis spectra of the obtained solutions were recorded using a Hitachi U-2900 spectrophotometer (Tokyo, Japan) in a 1 cm quartz cell.

The obtained results were expressed in μmol of Trolox equivalents per 100 g of sample.

2.8. Statistical Analysis

The AA of oil samples was determined five times within one day by the modified ABTS, DPPH, and FC methods. The obtained results were presented as mean (c) \pm standard deviation (SD).

3. Results

The enzymatic synthesis was carried out in milder conditions without organic solvents, while the yield of this reaction was 21%.

Bioaccessibility is the amount of a food constituent present in the gut due to its release from the solid food matrix, which affects the organism through the intestinal barrier [6,7]. Therefore, the AA of oil samples was determined before and after each simulated digestion step, and the data are presented in Table 1. The AA of non-supplemented oil was similar to the results reported in our previous work [9]. As seen, the antioxidant potential of rapeseed oils analyzed by three methods differed, probably due to different mechanisms of the used analytical methods. The refined rapeseed oil enriched with ESA had higher ABTS (1939.12–4036.24 $\mu\text{mol TE}/100\text{ g}$), DPPH (580.74–5183.95 $\mu\text{mol TE}/100\text{ g}$), and FC (292.28–581.98 $\mu\text{mol TE}/100\text{ g}$) results than oils without the new antioxidant (ABTS = 1277.62 $\mu\text{mol TE}/100\text{ g}$, DPPH = 520.32 $\mu\text{mol TE}/100\text{ g}$, and FC = 120.01 $\mu\text{mol TE}/100\text{ g}$). Furthermore, an increase in the radical scavenging activity determined by ABTS (2.08-fold) and DPPH (8.9-fold) assays and total phenolics analyzed by FC method (1.99-fold) for rapeseed oils spiked with increasing concentrations of ESA (0.02 and 0.5%) was found. Differences in antioxidant properties could be observed during the digestion process. However, after the two-stage digestion process, no significant changes in AA were noted for non-supplemented oil (Table 1). On the contrary, Seiquer et al. observed a 3-20-fold increase of AA measured by ABTS, DPPH, FRAP, and FC methods for extra virgin argan oil. It can be explained that, compounds in argan oil could be transformed into different forms with higher antioxidants potential during digestion [6].

Table 1. Antioxidant activity before and after two-step model in vitro digestion of refined rapeseed oil fortified with ESA.

Step of Digestion	Sample	ABTS ± SD	DPPH ± SD [μmol TE/100g]	FC ± SD
before digestion	RO	1277.62 ± 189.47	520.32 ± 5.69	120.01 ± 15.53
	RO + 0.02% ESA	1939.12 ± 22.98	580.74 ± 40.86	292.28 ± 6.99
	RO + 0.5% ESA	4036.24 ± 688.55	5183.95 ± 493.69	581.98 ± 15.71
after 1 step	RO	1631.97 ± 82.83	616.44 ± 32.69	231.31 ± 12.44
	RO + 0.02% ESA	1148.05 ± 80.40	533.83 ± 39.35	261.41 ± 19.00
	RO + 0.5% ESA	10,507.47 ± 626.17	3570.44 ± 333.11	422.40 ± 11.97
after 2 step	RO	761.37 ± 60.04	356.70 ± 44.66	61.46 ± 8.82
	RO + 0.02% ESA	1703.82 ± 20.42	447.67 ± 45.72	504.66 ± 21.23
	RO + 0.5% ESA	11,597.04 ± 646.61	4177.28 ± 374.25	486.23 ± 27.08
after 1 and 2 steps	RO	1489.95 ± 106.01	574.55 ± 42.96	127.29 ± 8.75
	RO + 0.02% ESA	1518.43 ± 77.76	525.71 ± 16.94	352.13 ± 15.03
	RO + 0.5% ESA	10,485.70 ± 1000.22	4684.67 ± 253.84	768.01 ± 7.47

The first step of digestion caused a 1.28–1.93-fold increase in AA of enriched rapeseed oils. In contrast, the second step 0.51–0.68-fold decreased the AA of those oils compared to non-supplemented oil before digestion. The fortification of rapeseed oil with 0.02% ESA resulted in the AA decrease in each digestion step, except the total phenolic contents determined after the second step by the FC method. Moreover, rapeseed oil fortified with 0.5% ESA had lower amounts of total phenolics and other compounds capable to scavenge DPPH radicals. Surprisingly, ABTS radical scavenging activity measured both hydrophilic and hydrophobic antioxidants, demonstrated an increase of 160% in gastric digestion and 187% in duodenal digestion of oil with 0.5% ESA. On the other hand, oil enriched with 0.5% of ESA characterized significantly higher anti-oxidant properties than non-supplemented oil. The AA values determined by ABTS, DPPH, and FC methods were about 7.04, 8.15, and 6.03-fold higher, respectively, than the antioxidant potential of the refined rapeseed oil without the synthetic antioxidant after the digestion process. For comparison, a significant decrease in ABTS values of extra virgin olive oil after in vitro gastric and small intestinal digestion was reported by other authors [10].

4. Conclusions

The ESA was successfully synthesized in an enzymatic way without organic solvents. The in vitro two-step digestion process slightly affects the antioxidant potential of the oil sample without ESA. Antioxidant properties of fortified oil strongly depend on the concentration of the added phenolic acid ester. The fortification of rapeseed oil with 0.02% ESA did not improve their radical scavenging activity measured by ABTS and DPPH methods, while total phenolic content measured by the FC method increased by about two times.

Nevertheless, an increase of ESA up to 0.5% significantly increased AA of the refined rapeseed oil. Additionally, the presence of ESA in a higher amount in oil probably caused the release of antioxidant compounds that reacted with the ABTS radicals and consequently improved antioxidant properties compared to oil samples before the digestion process. Therefore, ESA could be used as an efficient additive to enhance the antioxidant potential of vegetable oils.

Author Contributions: Conceptualization A.S.-C. and D.R.-K.; data curation, D.R.-K. and M.P.K.; formal analysis, D.R.-K.; investigation, I.W., D.R.-K.; methodology, D.R.-K., A.S.-C.; M.P.K.; supervision, A.S.-C. and M.P.K.; visualization, D.R.-K., M.P.K. and A.S.-C.; writing—original draft, D.R.-K.; writing—review and editing, A.S.-C. and M.P.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Polish National Science Center for the financial support: grant no. 2018/29/N/ NZ9/02748.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Acknowledgments: Rabiej-Kozioł D. is grateful for financial support from the Polish National Science Center for the financial support: grant no. 2018/29/N/ NZ9/02748. Authors wish to thank Novozymes Spain Ltd for presenting Novozyme 435.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Wroniak M.; Rękas A. Nutritional value of cold-pressed rapeseed oil during long term storage as influenced by the type of packaging material, exposure to light & oxygen and storage temperature. *J. Food Sci. Technol.* **2016**, *53*, 1338–1347. doi:10.1007/s13197-015-2082-y.
2. Szydłowska-Czerniak A.; Rabiej D.; Krzemiński M. Synthesis of novel octyl sinapate to enhance antioxidant capacity of rapeseed–linseed oil mixture. *J. Sci. Food Agric.* **2018**, *98*, 1625–1631. <https://doi.org/10.1002/jsfa.8637>
3. Li X.Q.; J, C.; Sun Y.Y.; Yang, M.L. Chu, X.G. Analysis of synthetic antioxidants and preservatives in edible vegetable oil by HPLC/TOF-MS. *Food Chem.* **2009**, *113*, 692–700. <https://doi.org/10.1016/j.foodchem.2008.07.072>
4. Menezes J.C.J.M.D.S.; Kamat S.P.; Cavaleiro J.A.S.; Gaspar A.; Garrido J.; Borges F. Synthesis and antioxidant activity of long chain alkyl hydroxycinnamates. *Eur. J. Med. Chem.* **2011**, *46*, 773–777. doi:10.1016/j.ejmech.2010.12.016.
5. Garrido J.; Gaspar A., Garrido, E.M., Miri R.; Tavakkoli M.; Pourali S.; Saso L.; Borges F.; Firuzi O. Alkyl esters of hydroxycinnamic acids with improved antioxidant activity and lipophilicity protect PC12 cells against oxidativ estress. *Biochimie* **2012**, *94*, 961–967. doi:10.1016/j.biochi.2011.12.015.
6. Seiquer I.; Rueda A.; Olalla M.; Cabrera-Vique C. Assessing the bioavailability of polyphenols and antioxidant properties of extra virgin argan oil by simulated digestion and Caco-2 cell assays. Comparative study with extra virgin olive oil. *Food Chem.* **2015**, *88*, 496–503. doi:10.1016/j.foodchem.2015.05.006.
7. Rubió L.; Macià A.; Castell-Auví A.; Pinent M.; Blay T.; Ardévol A.; Romero M.-P.; Motilva M.-J. Effect of the co-occurring olive oil and thyme extracts on the phenolic bioaccessibility and bioavailability assessed by in vitro digestion and cell models. *Food Chem* **2014**, *149*, 277–284. doi:10.1016/j.foodchem.2013.10.075.
8. Szydłowska-Czerniak A.; Łaszewska A. Effect of refining process on antioxidant capacity, total phenolics and prooxidants contents in rapeseed oils. *LWT–Food Sci Technol* **2015**, *64*, 853–859. doi:10.1016/j.lwt.2015.06.069
9. Rabiej D.; Szydłowska-Czerniak A. Antioxidant activity of rapeseed oil with octyl sinapate, an amphiphilic antioxidant, stored in various types of packages under various conditions. *J. Food Nutr. Res.* **2020**, *59*, 16–22.
10. Dinnella C.; Minichino P.; D’andrea A.M.; Monteleone E. Bioaccessibility and antioxidant activity stability of phenolic compounds from extra-virgin olive oils during in vitro digestion. *J. Agric. Food Chem.* **2007**, *55*, 8423–8429. <https://doi.org/10.1021/jf072244+>.