

# Filling biodiversity knowledge gaps: Sponges (Porifera: Demospongiae) recorded off San Jorge Gulf (Argentina), SW Atlantic Ocean<sup>†</sup>

Laura Schejter<sup>1,2,\*</sup> and Diego A. Giberto<sup>1,2,\*</sup>

<sup>1</sup> Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP)

<sup>2</sup> Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)

\* Correspondence: schejter@inidep.edu.ar (LS); diegogiberto@inidep.edu.ar (DAG)

<sup>†</sup> Presented the 2nd International Electronic Conference on Diversity - Animals, Plants and Microbes, online, 15–31 March 2022.

**Abstract:** The invertebrate by-catch of the Argentinean squid *Illex argentinus* collected by bottom trawls off San Jorge Gulf during an expedition performed in December 2020 was employed to characterize the bottom assemblages of the area, with special emphasis on sponges. According to the results, sponges were conspicuous and abundant components in these benthic communities, representing up to 40% of the total invertebrate bycatch (average: ~130 kg.nm<sup>2</sup> per site). *Tedania (T.) mucosa* was the dominant species. Other recorded species were *T. (T.) charcoti*, *T. (T.) murcochi*, *Isodictya verrucosa*, *Iophon proximum*, *Clathria (C.) microxa*, *C. (C.) discreta*, *Haliclona bilamellata*, *Siphonochalina fortis* and *Myxilla (M.) mollis*. The results of this study contribute to fill in the gaps on the distribution records of the sponges in the continental shelf of Argentina in a previously poorly-studied area.

**Keywords:** sponge richness; benthic communities; bycatch

## 1. Introduction

In Argentina, knowledge on the biodiversity of marine Porifera is closely related to the concentration of the sampling effort. Coastal areas such as Buenos Aires and Tierra del Fuego have several records, while the majority of the continental shelf have few or no records at all [1]. Pioneering studies, such as those developed after the “Challenger” and “Discovery” expeditions, attempt to determine and identify the sponge species inhabiting the Argentinian waters [2–10]. Later, several authors contributed to the study of sponges in specific areas [11–20]. The revision made by López Gappa and Landoni [1] provided the first checklist of marine sponges in Argentina and highlighted noticeable differences in records and sponge richness among different areas. In the past 15 years, there were new contributions to the study of sponges and also new species were described. The shelf break sponge-fauna was studied by Schejter et al. [21, 22] and Bertolino et al. [24], while interesting contributions were performed for Burdwood bank region by Schejter et al. [25, 26, 27] and for Bahía San Antonio by Gastaldi et al. [28]. However, according to our knowledge, there are no specific records of sponge species in the area located off San Jorge Gulf, Patagonia, only sparse mentions as a general group in technical guides or reports (i.e. [29]).

Considering the former background, the aim of the present contribution was to provide information on sponge richness and abundance from this scarcely studied area (off San Jorge Gulf, Patagonia) taking advantage of the bottom by-catch collected during a stock assessment expedition of the Argentinean squid *Illex argentinus*. Complementary,

Academic Editor: Matthieu Chauvat  
Published: 14 March 2022

**Publisher’s Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

we also provide some general information on the benthic communities of the area and other conspicuous by-catch invertebrate species.

## 2. Materials and Methods

In December 2020, the RV “Victor Angelescu” performed a regular stock assessment expedition in order to evaluate the population of the Argentinean squid *Illex argentinus* off San Jorge Gulf, between 44-46°S and isobaths 80m-120m depth. Sampling was developed using bottom trawls. At 23 random sites (Table 1), the invertebrate by-catch was preserved frozen and transported to the Benthos Laboratory of the INIDEP. Sorting of the species was performed at the laboratory and sponge morphospecies were separated and weighted (wet weight) using an Ohaus balance (precision 0.002kg). By-catch species were identified using local guides [29, 30], while sponge species were identified using the literature mentioned in the Introduction section. Biomass values (as densities) were calculated for each taxa, using the swept area for each sampling site, and it was expressed in kilograms per square nautical miles ( $\text{kg.nm}^{-2}$ ).

**Table 1.** Location of the sampling stations at the study area. Benthic by-catch and Porifera biomass values are also provided. .

Sampling Station	Latitude (S)	Longitude (W)	Depth (m)	Total benthic by-catch ( $\text{kg.nm}^{-2}$ )	Porifera ( $\text{kg.nm}^{-2}$ )
1	46.084	62.041	107	6.54	4.04
2	45.307	61.542	103	110.83	19.02
3	45.462	61.143	110	18.76	17.8
4	45.287	61.068	114	297.5	14.2
5	45.086	61.075	110	563.3	266.01
6	45.046	61.342	106	24.94	13.98
7	45.186	61.501	104	1687	390.65
8	45.237	63.466	100	43.59	36.46
9	45.472	64.040	98	40.57	38.72
10	45.426	62.386	97	114.1	109.26
11	45.189	62.386	106	669.1	656.46
12	45.054	62.309	105	75.59	72.62
13	44.563	62.096	104	568.3	458.32
14	44.496	61.145	108	667.2	36.55
15	44.390	61.388	106	93.47	82.03
16	44.450	62.540	100	403.3	137.27
17	44.206	63.410	89	505.1	138.39
18	44.229	63.015	96	89.95	74.70
19	44.269	62.288	100	98.06	57.28
20	44.161	61.047	104	650.9	39.68
21	44.086	62.186	97	271.2	130.71
22	44.163	61.403	105	127.5	63.69
23	44.019	61.182	103	36.49	8.26

## 3. Results

Our results showed that the benthic assemblage was composed by 42 taxa. Sponges were conspicuous and abundant components in the benthic communities off San Jorge Gulf, Argentina. They represented 40% of the total invertebrate by-catch in the area (average of 124.61 kg.nm<sup>-2</sup> per station, maximum value of 656.5 kg.nm<sup>-2</sup> at station 11, minimum value of 4.04 kg.nm<sup>-2</sup> at station 1) (Table 1). However, tunicates were the most important group in terms biomass (45%, average of 140.3 kg.nm<sup>-2</sup>), and the solitary ascidian *Paramolgula* sp. accounted for an average of 128.4 kg.nm<sup>-2</sup> per station. An unidentified colonial tunicate accounted for an average of 11.36 kg.nm<sup>-2</sup>. Other conspicuous, but less important groups in terms of biomass in this benthic community were crustaceans (6%), being the spider crab *Libidoclaea granaria* the most important one in terms of biomass, reaching in average 11.67 kg.nm<sup>-2</sup>. Polychaetes represented 5% of the benthic assemblage and the most conspicuous taxa was *Chaetopterus variopedatus* (average 15.61 kg.nm<sup>-2</sup>). Echinoderms only contributed in 2,3% to the benthic assemblage. It is worth to mention that egg cases of the catshark *Schroederichthys biviuis* were recorded at stations 13, 18 and 21, always attached to the sponge *Tedania (Tedaniopsis) mucosa* (Figure 1A).

In particular, 12 sponge taxa were recorded: *Tedania (T.) mucosa*, *T. (T.) charcoti*, *T. (Tedania) murdochi* Topsent, 1913, *Isodictya verrucosa*, *Iophon proximum*, *Clathria (Clathria) microxa*, *C. (C.) discreta*, *Siphonochalina fortis*, *Myxilla (M.) mollis*, *Haliclona bilamellata*, and also 2 unidentified species belonging to Suberitidae and Haplosclerida (Figure 1). *Tedania (T.) mucosa* Thiele, 1905, which was the most common and abundant sponge species recorded in the area, was registered in 22 sites (95% of the sites), reaching up to 98 % of the total wet biomass at station 11 (652.7 kg.nm<sup>-2</sup>), with an average of 100.8 kg.nm<sup>-2</sup>. The main characteristics of the sponge species recorded are listed below:

- *Tedania (T.) mucosa* specimens (Figure 1A) were massive, beige to red-brown in color, with a smooth surface and an evident ectosomal skeleton made of tornota arranged in palisade. This species was recorded in all the sampled sites, except at station 4. At stations 13, 18 and 21, egg cases of the catshark *Schroederichthys biviuis* were found strongly attached to the sponges, as described in [31]. Spicules are smooth styles of 230-255 by 10 µm; mucronate tornota of 190-220 by 5-7.5 µm and two categories of onichaete of 140-200 µm and 70-85 µm.


- *Tedania (Tedaniopsis) charcoti* specimens (Figure 1K) were massive, but fragile and friable in the dried state. The surface is uneven and looks porous, with no conspicuous ectosome. This species was recorded at stations 4, 5, 6, 11, 15, 17, 18 and 20. Spicules are styles of 245-380 by 5-10 µm; tornota of 180-250 by 5-7.5 µm and two categories of onichaete of 230-270 µm and 65-105 µm. *Tedania (T.) charcoti* accounted for 14.72 kg.nm<sup>-2</sup> in average in the studied area. This is a common species in Argentina [21], and our specimens fit very well with the previous records, although spicule dimensions in the holotype and Antarctic specimens are usually bigger.

- *Tedania (Tedania) murdochi* specimens (Figure 1H) were massive and rigid, beige in color, with some evident oscula. It was recorded at stations 10 and 17. Spicule dimensions are smaller than the other two recorded species: styles are 235-270 by 10-15 µm; tornota, 180-210 by 5-7.5 µm and onichaete are 155-205 µm.

- *Isodictya verrucosa* specimens (Figure 1B) have the typical morphology of the genus, yellow-beige in color, with a plumoreticulate skeleton. This species was recorded at stations 12 and 13. Spicules are oxeads of 310-400 by 15 µm and small isochelae of 25-30 µm. As noted by [21], megascleres are shorter than in the holotype and Antarctic specimens, but the morphology of the spicules very much agree with the description of the species. This is the second mention of the species for the SW Atlantic Ocean, after Schejter et al. [21].

- *Iophon proximum* specimens (Figure 1D) are massive and variable in shape, porous and became dark brown in color after preservation. It was found also associated (attached) with other sponge species. It was recorded at stations 2, 5, 7, 10, 14, 20 and 21. This is a common and highly variable species and an extended description and revision could be found at Desqueyroux-Faundez and van Soest [32].

**Figure 1.** Sponge species recorded in the present study. A. *Tedania (T.) mucosa* with an egg case of the catshark *Schroederichthys bivivus* attached; B. *Isodictya verrucosa*; C. *Clathria (C.) microxa* attached to a dead *Flabellum* skeleton; D. *Iophon proximum*; E. *Clathria (C.) discreta*; F. Haplosclerida; G. *Myxilla (M.) mollis*; H. *Tedania (T.) murdochi*; I. *Siphonochalina fortis*; J. *Haliclona bilamellata*; K. *Tedania (T.) charcoti*; L. Suberitidae.



- *Clathria (Clathria) microxa* specimen (Figure 1C) is bushy and was found growing attached to a dead *Flabellum* coral. It was recorded at station 5. Spicules are two types of styles, 370-440 by 15-20  $\mu\text{m}$  and 320-445 by 5-10  $\mu\text{m}$ , acanthostyles of 100-115 by 5  $\mu\text{m}$ , toxas 230-460  $\mu\text{m}$  and microxeas of 45  $\mu\text{m}$ . This is the third mention of the species for the SW Atlantic Ocean, after Schejter et al. [21, 22].

- *Clathria (C.) discreta* specimen (Figure 1E) was found at station 14. This is a common, well-registered species in Argentina and detailed information could be found at Burton [8, 10] and Schejter et al. [21].

- *Siphonochalina fortis* specimen (Figure 1I) was found at station 17, and it is characterized by their ramose morphology, highly compressible, with a skeleton conformed by quadrangular meshes of fibres with embedded oxeas of 55-60 by 3.5-5  $\mu\text{m}$ . It is a common species in Argentina.

- *Myxilla (Myxilla) mollis* specimen (Figure 1G) was recorded at station 7 and it is a very common species in Argentina. A detailed description could be found at Bertolino et al. [24].

- *Haliclona bilamellata* specimen (Figure 1J) has a massive to tubular morphology, soft and sticky, and was recorded at station 17. It has oxeas of 160-185 by 10  $\mu\text{m}$ , and also some smaller and thinner of 140 by 5  $\mu\text{m}$ . Our specimen agrees with the descriptions provided in Burton [8] and Sará [12].

- Several unidentified specimens belonging to the Family Suberitidae (Figure 1L) were found at stations 17, 18, 21 and 22, and two fragments belonging to the order Haplosclerida (Figure 1F) were found at station 11. These specimens will require more studies to get a proper identification at a species level.

#### 4. Discussion

The results of this study contribute to better explore the benthic assemblage off San Jorge Gulf, Patagonia and fill in the gaps on the records and distribution of the sponges in this particular unknown region. Among the 42 registered taxa, 12 sponge species were recorded in this area. Although all these species were previously recorded in other sectors of the Argentinian waters, this new evidence contribute to build the distribution pattern of sponge species in Argentina. Additionally, it is worth to mention that the studied area presented an unusually high abundance of sponges and tunicates in the benthic communities that may reach up to 40% and 45% in wet weight, respectively, of the total by-catch. This is probably an indication that this area was relatively preserved from intense bottom fishing (i.e.[33]) and may constitute a potential refuge for juveniles of species of commercial and non-commercial interest, as suggested for some regions of the San Jorge Gulf [34]. It must be noted that Argentinean hake (*Merluccius hubbsi*) fishing (developed using bottom trawlers) is banned at the studied area, but the Argentine red shrimp (*Pleoticus muelleri*) fishing (developed also by a trawler fleet) is allowed. Finally, and considering the finding of egg cases of the catshark *S. bivi*, this area may also have interesting characteristics for oviparous chondrictian species and should be better explored in order to understand if it should be managed in order to preserve target species from anthropogenic impacts.

**Funding:** The expedition was financed by INIDEP. The study was partially financed by PICT 2019-4233.

**Acknowledgments:** Authors thanked the INIDEP colleagues that collect the invertebrate by-catch samples during the stock assessment of the Argentinean short-finned squid.

**Conflicts of Interest:** "The authors declare no conflict of interest."

#### References

- López Gappa J.; Landoni N. A., Biodiversity of Porifera in the Southwest Atlantic between 35 S and 56 S. *Revista del Museo Argentino de Ciencias Naturales* **2005**, *7*, 191–219.
- Ridley S.O.; Dendy A. Preliminary report on the Monaxonida collected by H.M.S. Challenger. Part I. *Annals and Magazine of Natural History* **1886**, *18*, 325–351, 470–493.
- Ridley S.; Dendy A. Report on the Monoaxonida collected by H.M.S. Challenger during the year 1873-1876. In *Report on the Scientific Results of the voyage of H.M.S. Challenger, 1873-1876, Zoology*, 1887. Vol. 20, 1–275.
- Sollas W.J. Preliminary account of the Tetractinellid sponges dredged by H.M.S. Challenger 1872-76. Part I. The Choristida. *Scientific Proceedings of the Royal Dublin Society*, 1886. Vol. 5, 177–199.
- Sollas W.J. Report on the Tetractinellida collected by H.M.S. Challenger during the years 1873-1876. In *Report on the scientific results of the voyage of H.M.S. Challenger, 1873-1876. Zoology*, 1888. Vol. 25, 1-458.
- Schulze F.E. Report on the Hexactinellida collected by H.M.S. Challenger during the years 1873-76. In *Report on the Scientific Results of the voyage of H.M.S. Challenger, 1873-1876. Zoology*, 1887. Vol. 21, 1-513.
- Topsent E. Spongiaires de l'expédition antarctique nationale ecossaise. *Transactions of the Royal Society of Edinburgh* **1913**, *49*, 579-643.
- Burton M. Sponges. *Discovery Report* **1932**, *6*, 237–392.
- Burton M. Sponges. *Further Zoological Results Swedish Antarctic Expedition 1901-1903 under the direction of Dr. Otto Nordenskjöld* **1934**, *3*, 1–58.
- Burton M. Las esponjas marinas del Museo Argentino de Ciencias Naturales. *Anales del Museo Argentino de Ciencias Naturales Bernardino Rivadavia* **1940**, *40*, 95–121.
- Boury-Esnault N. Spongiaires. Campagne de la "Calypso" au large des côtes atlantiques de l'Amérique du Sud (1961-1962). *Annales de l'Institut océanographique* **1973**, *49*, 263–294.
- Sará M. Demospongie di acque superficiali della Terra del Fuoco (Spedizione A.M.F. Mares-G.R.S.T.S. e S.A.I.). *Bolletino dei Musei e degli Istituti Biologici dell'Università di Genova* **1978**, *46*, 7–117.

13. Mothes de Moraes B., Marques Pauls S. Algunas esponjas monoaxonidas (Porifera: Demospongiae) do litoral sul do Brasil, Uruguay e Argentina. *Iheringia Série Zoologia* **1979**, *54*, 57–66. 192  
193
14. Cuartas E.I. Poríferos de la Provincia Biogeográfica Argentina. *Physis A* **1986**, *44*, 37–41. 194
15. Cuartas E.I. Demospongiae (Porifera) de Mar del Plata (Argentina) con la descripción de *Cliona lisa* sp.n. y *Plicatellopsis reptans* sp.n. *Neritica* **1991**, *6*, 43–63. 195  
196
16. Cuartas E.I. Poríferos intermareales de San Antonio Oeste, provincia de Río Negro, Argentina (Porifera: Demospongiae). *Neotrópica* 1992a. *38*, 111–118. 197  
198
17. Cuartas E.I. Poríferos de la Provincia Biogeográfica Argentina. III. Poecilosclerida (Demospongiae) del litoral marplatense. *Physis A* 1992 b.47: 73–88. 199  
200
18. Cuartas E.I. Algunas Demospongiae (Porifera) de Mar del Plata, Argentina, con descripción de *Axociella marplatensis* sp.n. *Iheringia Serie Zoología* 1992 c. *73*, 3-12. 201  
202
19. Cuartas E.I. Redescrición de *Clathria burtoni* “nomen novum” de *C. prolifera* Burton 1940. (Porifera: Demospongiae). *Annali del Museo Civico di Storia Naturale “G. Doria”* **1995**, *90*, 571–576. 203  
204
20. Genzano G.; Cuartas E.I.; Excoffon A. Porifera y cnidaria de la campaña “Oca Balda” 05/88. *Talassas* **1991**, *9*, 63–78. 205
21. Schejter L.; Calcinai B.; Cerrano C.; Bertolino M.; Pansini M.; Giberto D.; Bremec C. Porifera from the Argentine Sea: Diversity in Patagonian scallop beds. *Italian Journal of Zoology* **2006**, *73*, 373–385. 206  
207
22. Schejter L.; Bertolino M.; Calcinai B.; Cerrano C.; Bremec C., Epibiotic sponges on the hairy triton *Fusitriton magellanicus* in the SW Atlantic Ocean, with the description of *Myxilla (Styloptilon) canepai* sp. nov. *Aquatic Biology* **2011**, *14*, 9–20. 208  
209
23. Schejter L.; Chiesa I. L.; Doti B. L.; Bremec C. *Mycale (Aegogropila) magellanica* (Porifera: Demospongiae) in the southwestern Atlantic Ocean: endobiotic fauna and new distributional information. *Scientia Marina* **2012**, *76*, 753–761. 210  
211
24. Bertolino M.; Schejter L.; Calcinai B.; Cerrano C.; Bremec C. Sponges from a submarine canyon of the Argentine Sea. In *Porifera Research: Biodiversity, Innovation, Sustainability*, Custódio, M. R.; Hajdu, E.; Lôbo-Hajdu, G.; Muricy, G., Eds. Museu Nacional: Rio de Janeiro, 2007. 189–201. 212  
214
25. Schejter L.; Rimondino C.; Chiesa I.; Díaz de Astarloa J. M.; Doti B. L.; Elías R.; Escolar M.; Genzano G.; López Gappa J.; Tatián M.; Zelaya D.G.; Cristobo J.; Perez C.D.; Cordeiro R.T.; Bremec C.S. Namuncurá Marine Protected Area: an oceanic hot spot of benthic biodiversity at Burdwood Bank, Argentina. *Polar Biol.* **2016**, *39*, 2373–2386. 215  
216  
217
26. Schejter L.; Bertolino M.; Calcinai B. Description of *Antho (Plocamia) bremecae* sp. nov and checklist of Microcionidae (Demospongiae: Poecilosclerida) from Burdwood Bank and neighboring areas, SW Atlantic Ocean. *Zootaxa* **2017**, *4312*, 580–594. 218  
219
27. Schejter L.; Albano M. Benthic communities at the marine protected area Namuncurá/ Burdwood bank, SW Atlantic Ocean: detection of vulnerable marine ecosystems and contributions to the assessment of the rezoning process. *Polar Biol.* **2021**, *44*, 2023–2037. 220  
221  
222
28. Gastaldi M.; De Paula T. S.; Narvarte M.; Lobo-Hajdú G.; Hajdu E. Marine sponges (Porifera) from the Bahía San Antonio (North Patagonian Gulfs, Argentina), with additions to the phylogeography of the widely distributed *Cliona* aff. *celata* and *Hymeniacion perlevis*, and the description of two new species. *Marine Biol. Res.* **2018**, *14*, 682–716. 223  
224  
225
29. Roux A.; Piñero R.; de la Garza J. *Guía para la identificación de la fauna asociada a la pesquería de langostino patagónico*. Instituto Nacional de Investigación y Desarrollo Pesquero, Mar del Plata, 2007. 32pp. 226  
227
30. Bremec C; Marecos A.; Schejter L.; Lasta M. Guía técnica para la identificación de invertebrados epibentónicos asociados a los bancos de vieira patagónica (*Zygochlamys patagonica*) en el Mar Argentino. *Publicaciones Especiales INIDEP*, Mar del Plata, 2003. 28pp. 228  
229  
230
31. Vazquez D. M.; Belleggia M.; Schejter L.; Mabragna E. Avoiding being dragged away: Finding egg cases of *Schroederichthys bivius* (Chondrichthyes: Scyliorhinidae) associated with benthic invertebrates. *J. Fish Biol.* **2018**, *92*, 248–253. 231  
232
32. Desqueyroux-Faundez R.; van Soest R. A review of Iophonidae, Myxillidae and Tedaniidae occurring in the South East Pacific (Porifera: Poecilosclerida). *Revue Suisse de Zoologie* **1996**, *103*, 3–79. 233  
234
33. Wassenberg T.J.; Dews G.; Cook S.D. The impact of fish trawls on megabenthos (sponges) on the north-west shelf of Australia. *Fisheries Research* **2002**, *58*, 141–151 235  
236
34. Giberto D. A.; Roux A.; Bremec C. S. Capítulo 4. Ecosistemas norpatagónicos: Golfo San Jorge y Península Valdés. In *Comunidades bentónicas en regiones de interés pesquero de la Argentina*, Bremec C. S.; Giberto D. A. (Eds). Publicación Especial INIDEP. Mar del Plata, Argentina, 2017; pp. 47–56. 237  
238  
239