

Plant-Based Nanoantibiotics: An Effective Strategy to Overcoming on Antibiotic Resistance [†]

Sepidar Seyyedi Mansour ¹, Maria Carpena ¹, Javier Echave ¹, Franklin Chamorro ¹, Antia G. Pereira ^{1,2}, Marta Barral-Martinez ¹, Anton Soria-Lopez ¹, Lucia Cassani ^{1,4}, Aurora Silva ^{1,3}, Jianbo Xiao ^{1,5}, Jesus Simal-Gandara ^{1,*} and Miguel A. Prieto ^{1,2,*}

- ¹ Nutrition and Bromatology Group, Department of Analytical and Food Chemistry, Faculty of Food Science and Technology, University of Vigo, Ourense Campus, E32004 Ourense, Spain; sepidar.seyyedi@uvigo.es (S.S.M.); mcarpena@uvigo.es (M.C.); javier.echave@uvigo.es (J.E.); franklin.noel.chamorro@uvigo.es (F.C.); antia.gonzalez.pereira@uvigo.es (A.G.P.); marta.barral@uvigo.es (M.B.-M.); anton.soria@uvigo.es (A.S.-L.); lucia.victoria.cassani@uvigo.es (L.C.); mass@isep.ipp.pt (A.S.); jianbo.xiao@uvigo.es (J.X.)
 - ² Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolonia, 5300-253 Bragança, Portugal
 - ³ REQUIMTE/LAQV, Instituto Superior de Engenharia do Porto, Instituto Politécnico do Porto, Rua Dr António Bernardino de Almeida 431, 4200-072 Porto, Portugal
 - ⁴ Instituto de Investigaciones en Ciencia y Tecnología de Materiales (INTEMA, CONICET), Mar del Plata (7600), Colón 10850, Argentina
 - ⁵ International Joint Research Laboratory of Intelligent Agriculture and Agri-products Processing, Jiangsu University, Zhenjiang 212013, China
- * Correspondence: jsimal@uvigo.es (J.S.-G.); mprieto@uvigo.es (M.A.P.)
[†] Presented at the The 2nd International Electronic Conference on Antibiotics—Drugs for Superbugs: Antibiotic Discovery, Modes of Action And Mechanisms of Resistance, 15–30 June 2022; Available online: <https://eca2022.sciforum.net/>.

Citation: Seyyedi Mansour, S.; Carpena, M.; Echave, J.; Chamorro, F.; Pereira, A.G.; Barral-Martinez, M.; Soria-Lopez, A.; Cassani, L.; Silva, A.; Xiao, J.; et al. Plant-based nanoantibiotics: An effective strategy to overcoming on antibiotic resistance. *Med. Sci. Forum* **2022**, *2*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor:

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Commonly used antibiotics use multiple administrations for providing a continuous bactericidal effect but can increase systemic toxicity, as well as lead to bacterial drug resistance [1,2]. The use of plant-based antibacterial compounds in combination with conventional antibiotics to treat drug-resistant infections could be an alternative to overcoming the problem of bacterial resistance [3]. The combination of antibiotics with plant-based antibacterial compounds has been shown synergistic advantages due to the inhibition of drug efflux and the presence of alternative mechanisms of action [4]. Moreover, the nanoencapsulation of antibiotics is another effective technique to overcome antibiotic resistance [1]. Nano-encapsulated antimicrobials have a better performance in comparison with traditional antibiotics due to their small size which leads to better interaction with bacterial cells [5]. The different nanocarriers are effective in efficiently administering antibiotics by improving pharmacokinetics and accumulation while reducing the adverse effects [6]. Additionally, the surface engineering of nanocarriers provides benefits such as targeting and modulating various resistance mechanisms [7]. Furthermore, most nanocarriers are suitable platforms for co-loading of plant-based antibacterial compounds and traditional antibiotics to provide synergistic effects [8]. This study outlines recent attempts to combat infectious diseases, with a focus on the use of plant-based nanoantibiotics as novel tools to address today's issues in infectious disease treatment.

Keywords: nanocarriers; antibacterial compounds; plants extracts; synergistic activity

1. Introduction

Infectious diseases were the greatest cause of death worldwide at the turn of the 20th century. The introduction of antimicrobial drugs was largely responsible for the decline in morbidity and mortality from these diseases throughout the last century. Antibiotics

are frequently used to prevent and cure bacterial infections in medical care. On the other hand, free antibiotics have a low in vivo ingestion rate and are unable to effectively accumulate at the infection site [8]. As a result, a sizeable antibiotic dosage is necessary to obtain an effective bactericidal concentration and resulting in systemic drug distribution. Antibiotic misuse and overuse, together with the fast mutation and evolution of bacteria, have resulted in a growing number of side effects that influence human health and result in the rise of drug-resistant microorganisms capable of surviving traditional antibiotic therapies. Long-term or high-dose antibiotic therapy may produce toxicity in human tissues and organs to variable degrees [9]. Widespread antibiotic use contributes to the expansion of drug-resistant bacteria, which pose a severe threat to human health. Thus, the long-term treatment of bacterial illnesses requires the discovery of novel and effective antibacterial medicines [10]. To avoid these undesirable effects associated with antibiotics, current research is focusing on the use of plant-derived extracts as antibiotic substitutes [11]. In plant-based edible materials, a large range of natural compounds with antimicrobial activity has been found, although the efficacy of natural antimicrobials can be influenced by a variety of factors [12]. The bioavailability of these plant-based compounds is limited by their hydrophobicity, volatility, and instability, which dramatically reduces their biological and antibacterial action. A variety of techniques for encapsulating these plant-derived antibacterial substances have been reported to meet these issues. By altering antibiotic delivery systems on the nanoscale, nanotechnology is increasingly being used to build effective antimicrobial delivery systems [13]. In the context of research on this novel technique, this study proposes using nanotechnology as a new paradigm in controlling infectious diseases, particularly in overcoming antimicrobial medication resistance, with an emphasis on plant-based nanoantibiotics.

2. Plant-Based Antimicrobials

Numerous plants produce secondary metabolites and other compounds to ward off pathogens such as microorganisms (bacteria, fungi, and viruses), insects, and herbivores. These plant-derived chemicals are classified according to their botanical source, chemical structure, and mode of action [14]. Recently, there has been renewed interest in isolating natural antimicrobials from waste materials recovered from the food and agricultural industries, including such waste products of fruits and vegetables. Among them, some of the most significant plant-based antimicrobial compounds are highlighted below.

2.1. Phenols and Polyphenols

Phenols and polyphenols are secondary metabolites produced by plants to protect them from bacteria, insects, and herbivores. Simple phenols, phenolic acids, flavonoids, quinines, tannins, and coumarins are some of the main antimicrobial classes of polyphenols [15]. The toxicity of simple phenols and phenolic acids to microorganisms is dependent on the hydroxylation site(s) and degree of hydroxylation, with a more oxidized structure exhibiting more inhibitory activity [16]. Other studies believe that phenolic acids' antibacterial effect is due to membrane disruption, which causes leaking of critical intracellular components, direct changes in microbial metabolism, and inactivation of essential substrates for microbial development. Figure 1 schematically depicts phenolics' capacity to inhibit bacteria by modifying their surface characteristics [17].




Figure 1. The schematic of the used liposomes and polymer-based NPs for encapsulation of natural antimicrobials.

2.2. Essential Oils

Essential oils (EOs) are complex extracts from aromatic plants that contain a variety of aldehydes, terpenes, and phenols. They are widely known for their biological and antibacterial activities. EOs and their components have demonstrated remarkable antimicrobial activity against multidrug-resistant (MDR) pathogens such as methicillin-resistant *Staphylococcus aureus* [18]. Because of its impact on one or more cell components, EOs can affect bacterial cell integrity, resulting in growth inhibition or, at high enough concentrations, cell death. Thus, using EOs to fight or inhibit MDR bacteria development shows promise [19].

3. Nanoencapsulation of Plant-Based Antimicrobials

An efficient antimicrobial agent should interact with the desired target while interacting with nonspecific targets as little as possible to improve potency and avoid adverse effects. Plant-derived antimicrobials like EOs have some limitations in this aspect since they are highly influenced by environmental conditions. In particular, poor water solubility decreases bioavailability and significantly reduces antibacterial activity, whereas volatility makes prolonged release and regulated delivery difficult [3]. To address these problems, a range of techniques for encapsulating plant-derived antibacterial compounds as colloidal systems have been reported, including nanoemulsions, lipid-based nanocarriers (e.g., liposomes, and solid lipid nanocapsules), and polymer-based carriers [20]. Moreover, the majority of nanocarriers are suitable platforms for the co-loading of plant-based antibacterial agents and conventional antibiotics to generate synergistic effects [8].

3.1. Antimicrobial Nanoemulsions

Oil-in-water nanoemulsions, including lipid nanoparticles (NPs) disseminated in water, are the most often employed nano-enabled delivery method for natural based antimicrobials. Antimicrobial nanoemulsions can be classified according to the characteristics and location of the antimicrobials they carry. Nanoemulsions are colloidal dispersions composed of 2 incompatible fluids, one distributed in the other as small particles ($d < 200$ nm). Although these systems are thermodynamically unstable, they can be made to be metastable for long enough durations to be useful in a variety of commercial applications [13,21].

3.2. Liposome-Based Nano-Antibiotics

Liposomes are nano- to micro- sized vesicles made up of a bilayer of phospholipids and an aqueous core. Liposomes have been extensively studied as potentially useful carriers of enzymes, proteins, and drugs for the treatment of a variety of diseases [22]. Liposomes are also the most extensively utilized antimicrobial drug delivery vehicle due to their bilayer lipid structure, which mimics the cell membrane and readily fuses with infectious bacteria. Additionally, both hydrophilic and hydrophobic antimicrobial agents can be encapsulated and maintained in the aqueous core and the phospholipids bilayer, respectively, without requiring chemical modification [6].

3.3. Polymer-Based Nano-Antibiotics

Antibiotics encapsulated in polymeric NPs have various advantages: (1) structural stability in biological fluids and under harsh and varied situations for preparing (e.g., spray drying and ultrafine milling) and storage, (2) precisely tunable characteristics (e.g., size, zeta-potentials, and drug release profiles) by attempting to manipulate polymer lengths, surfactants, and organic solvents used throughout NPs preparation, and (3) simple and versatile surface functionalization for conjugating drugs and targeting ligands. Polymeric NPs have been investigated for their ability to administer a variety of antimicrobial agents, with significantly increased therapeutic efficacy in the treatment of a variety of infectious diseases [6,23,24]. Figure 1 illustrates liposomes and polymer-based NPs used to encapsulate natural antimicrobials.

4. Concluding Remarks

Antibiotics have saved a huge number of lives from a variety of infectious diseases for more than half a century. However, the emergence of antibiotic resistance gained by microbial variations poses a significant challenge to the fight against infectious diseases. The use of antimicrobial compounds derived from plants instead of conventional antibiotics has become a significant topic of research. Numerous of these chemicals are incompatible with utilization in their pure form due to solubility, stability, and activity concerns. As a result, nanoemulsion-based delivery systems are being developed to address these constraints. Nanomaterials are prospective antibacterial agents of a new class due to their high surface area to volume ratio and unique physicochemical characteristics.

Taken together, the utilization of plant-derived NPs has demonstrated benefits and applications in medicine and the pharmaceutical sector. More precisely, investigations demonstrate that NPs possess antibacterial characteristics on their own or in conjunction with antibiotics, alleviating the current problem of acquired resistance caused by antibiotic overuse or misuse. With this application in mind, future research should focus on two areas: determining the safety of plant-based NPs (toxicity to humans and the environment) and mitigating the environmental impact of their synthesis. Not only is it necessary to refine the synthesis procedures, but also to investigate the mechanisms by which NPs exert their antibacterial effect, to transform plant-based NPs into a viable strategy capable of meeting society's demand for an effective solution to antibiotic resistance.

Author Contributions: Conceptualization, S.S.M., M.C., J.E., F.C., A.G.-P., M.B.-M., A.S.-L., L.C., A.S., J.X., J.S.-G. and M.A.P.; methodology S.S.M., J.E., F.C., A.G.-P., M.B.-M., A.S.L., and A.S.; validation, L.C., A.S., J.X., J.S.-G. and M.A.P.; investigation, S.S.M., M.C., L.C.; writing—original draft preparation, S.S.M. and M.C.; writing—review and editing, S.S.M. and M.C.; visualization, S.S.M., A.S., L.C. and M.C.; supervision, J.X., J.S.-G. and M.A.P. All authors have read and agreed to the published version of the manuscript.

Funding: The JU receives support from the European Union's Horizon 2020 research and innovation program and the Bio Based Industries Consortium. The project SYSTEMIC Knowledge hub on Nutrition and Food Security, has received funding from national research funding parties in Belgium (FWO), France (INRA), Germany (BLE), Italy (MIPAAF), Latvia (IZM), Norway (RCN), Portugal (FCT), and Spain (AEI) in a joint action of JPI HDHL, JPI-OCEANS and FACCE-JPI launched

in 2019 under the ERA-NET ERA-HDHL (n° 696295). The authors would like to thank the EU and FCT for funding through the programs UIDB/50006/2020; UIDP/50006/2020 and project PTDC/OCE-ETA/30240/2017- SilverBrain-From sea to brain: Green neuroprotective extracts for nanoencapsulation and functional food production (POCI-01-0145-FEDER-030240).

Institutional Review Board Statement:

Informed Consent Statement:

Data Availability Statement:

Acknowledgments: The research leading to these results was supported by MICINN supporting the Ramón y Cajal grant for M.A. Prieto (RYC-2017-22891) and Jianbo Xiao (RYC-2020-030365-I) and the FPU grant for A. Soria-Lopez (FPU2020/06140); by Xunta de Galicia for supporting the program EXCELENCIA-ED431F 2020/12, the post-doctoral grant of L. Cassani (ED481B-2021/152), and the pre-doctoral grants of A.G. Pereira (ED481A-2019/0228) and M. Carpena (ED481A 2021/313). The authors thank the program BENEFICIOS DO CONSUMO DAS ESPECIES TINTORERA-(CO-0019-2021) that supports the work of F. Chamorro. The research leading to these results was supported by the EcoChestnut Project (Erasmus+ KA202) that supports the work of J. Echave. Authors are grateful to Ibero-American Program on Science and Technology (CYTED—AQUA-CIBUS, P317RT0003), to the Bio Based Industries Joint Undertaking (JU) under grant agreement No 888003 UP4HEALTH Project (H2020-BBI-JTI-2019) that supports the work of M. Barral-Martínez.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Nagarajan, K.; Perumal, S.K.; Marimuthu, S.K.; Palanisamy, S.; Subbiah, L. Addressing Antimicrobial Resistance Through Nanoantibiotics: Challenges and Novel Strategies. In *Handbook of Research on Nano-Strategies for Combatting Antimicrobial Resistance and Cancer*; IGI Global: Pennsylvania, PA, USA, 2021; pp. 56–86. <https://doi.org/10.4018/978-1-7998-5049-6.ch003>.
- Pancu, D.F.; Scurtu, A.; Macasoi, I.G.; Marti, D.; Mioc, M.; Soica, C.; Coricovac, D.; Horhat, D.; Poenaru, M.; Dehelean, C. Antibiotics: Conventional therapy and natural compounds with antibacterial activity—a pharmaco-toxicological screening. *Antibiotics* **2021**, *10*, 401. <https://doi.org/10.3390/antibiotics10040401>.
- Abd El-Hack, M.E.; El-Saadony, M.T.; Saad, A.M.; Salem, H.M.; Ashry, N.M.; Abo Ghanima, M.M.; Shukry, M.; Swelum, A.A.; Taha, A.E.; El-Tahan, A.M.; et al. Essential oils and their nanoemulsions as green alternatives to antibiotics in poultry nutrition: A comprehensive review. *Poult. Sci.* **2022**, *101*, 101584. <https://doi.org/10.1016/j.psj.2021.101584>.
- Ali, K.; Dwivedi, S.; Azam, A.; Saquib, Q.; Al-Said, M.S.; Alkhedhairi, A.A.; Musarrat, J. Aloe vera extract functionalized zinc oxide nanoparticles as nanoantibiotics against multi-drug resistant clinical bacterial isolates. *J. Colloid Interface Sci.* **2016**, *472*, 145–156. <https://doi.org/10.1016/j.jcis.2016.03.021>.
- Maruthupandy, M.; Rajivgandhi, G.; Muneeswaran, T.; Song, J.M.; Manoharan, N. Biologically synthesized zinc oxide nanoparticles as nanoantibiotics against ESBLs producing gram negative bacteria. *Microb. Pathog.* **2018**, *121*, 224–231. <https://doi.org/10.1016/j.micpath.2018.05.041>.
- Hadinoto, K.; Cheow, W.S. Nano-antibiotics in chronic lung infection therapy against *Pseudomonas aeruginosa*. *Colloids Surfaces B Biointerfaces* **2014**, *116*, 772–785. <https://doi.org/10.1016/j.colsurfb.2014.02.032>.
- Yang, X.; Ye, W.; Qi, Y.; Ying, Y.; Xia, Z. Overcoming Multidrug Resistance in Bacteria Through Antibiotics Delivery in Surface-Engineered Nano-Cargos: Recent Developments for Future Nano-Antibiotics. *Front. Bioeng. Biotechnol.* **2021**, *9*, 696514. <https://doi.org/10.3389/fbioe.2021.696514>.
- Huang, W.; Zhang, Q.; Li, W.; Yuan, M.; Zhou, J.; Hua, L.; Chen, Y.; Ye, C.; Ma, Y. Development of novel nanoantibiotics using an outer membrane vesicle-based drug efflux mechanism. *J. Control. Release* **2020**, *317*, 1–22. <https://doi.org/10.1016/j.jconrel.2019.11.017>.
- Tamma, P.D.; Avdic, E.; Li, D.X.; Dzintars, K.; Cosgrove, S.E. Association of adverse events with antibiotic use in hospitalized patients. *JAMA Intern. Med.* **2017**, *177*, 1308–1315. <https://doi.org/10.1001/jamainternmed.2017.1938>.
- Omolo, C.A.; Kalhapure, R.S.; Agrawal, N.; Jadhav, M.; Rambharose, S.; Mocktar, C.; Govender, T. A hybrid of mPEG-b-PCL and G1-PEA dendrimer for enhancing delivery of antibiotics. *J. Control. Release* **2018**, *290*, 112–128. <https://doi.org/10.1016/j.jconrel.2018.10.005>.
- Amato, D.N.; Amato, D.V.; Mavrodi, O.V.; Braasch, D.A.; Walley, S.E.; Douglas, J.R.; Mavrodi, D.V.; Patton, D.L. Destruction of Opportunistic Pathogens via Polymer Nanoparticle-Mediated Release of Plant-Based Antimicrobial Payloads. *Adv. Healthc. Mater.* **2016**, *5*, 1094–1103. <https://doi.org/10.1002/adhm.201500974>.
- Das, S.; Singh, V.K.; Dwivedy, A.K.; Chaudhari, A.K.; Deepika; Dubey, N.K. Eugenol loaded chitosan nanoemulsion for food protection and inhibition of Aflatoxin B1 synthesizing genes based on molecular docking. *Carbohydr. Polym.* **2021**, *255*, 117339. <https://doi.org/10.1016/j.carbpol.2020.117339>.
- McClements, D.J.; Das, A.K.; Dhar, P.; Nanda, P.K.; Chatterjee, N. Nanoemulsion-Based Technologies for Delivering Natural Plant-Based Antimicrobials in Foods. *Front. Sustain. Food Syst.* **2021**, *5*, 643208. <https://doi.org/10.3389/fsufs.2021.643208>.

14. Savoia, D. Plant-derived antimicrobial compounds: Alternatives to antibiotics. *Future Microbiol.* **2012**, *7*, 979–990.
15. Das, A.K.; Nanda, P.K.; Chowdhury, N.R.; Dandapat, P.; Gagooua, M.; Chauhan, P.; Pateiro, M.; Lorenzo, J.M. Application of pomegranate by-products in muscle foods: Oxidative indices, colour stability, shelf life and health benefits. *Molecules* **2021**, *26*, 467. <https://doi.org/10.3390/MOLECULES26020467>.
16. Coppo, E.; Marchese, A. Antibacterial Activity of Polyphenols. *Curr. Pharm. Biotechnol.* **2014**, *15*, 380–390. <https://doi.org/10.2174/138920101504140825121142>.
17. Dietrich, H.; Pour Nikfardjam, M.S. Influence of phenolic compounds and tannins on wine-related microorganisms. In *Biology of Microorganisms on Grapes, in Must and in Wine*; Springer, Cham, Switzerland, **2017**, 421–454. https://doi.org/10.1007/978-3-319-60021-5_18.
18. Fadli, M.; Chevalier, J.; Saad, A.; Mezrioui, N.E.; Hassani, L.; Pages, J.M. Essential oils from Moroccan plants as potential chemosensitisers restoring antibiotic activity in resistant Gram-negative bacteria. *Int. J. Antimicrob. Agents* **2011**, *38*, 325–330. <https://doi.org/10.1016/j.ijantimicag.2011.05.005>.
19. Rai, M.; Kon, K. *Fighting Multidrug Resistance With Herbal Extracts, Essential Oils And Their Components*; Academic Press: Cambridge, MA, USA, 2013; ISBN 9780123985392.
20. Anand, U.; Carpena, M.; Kowalska-Góralaska, M.; Garcia-Perez, P.; Sunita, K.; Bontempi, E.; Dey, A.; Prieto, M.A.; Proćków, J.; Simal-Gandara, J. Safer plant-based nanoparticles for combating antibiotic resistance in bacteria: A comprehensive review on its potential applications, recent advances, and future perspective. *Sci. Total Environ.* **2022**, *821*, 153472. <https://doi.org/10.1016/j.scitotenv.2022.153472>.
21. Salvia-Trujillo, L.; Martin-Belloso, O.; Julian McClements, D. Excipient nanoemulsions for improving oral bioavailability of bioactives. *Nanomaterials* **2016**, *6*, 17. <https://doi.org/10.3390/nano6010017>.
22. Huh, A.J.; Kwon, Y.J. “Nanoantibiotics”: A new paradigm for treating infectious diseases using nanomaterials in the antibiotics resistant era. *J. Control. Release* **2011**, *156*, 128–145. <https://doi.org/10.1016/j.jconrel.2011.07.002>.
23. Ungaro, F.; D’Angelo, I.; Coletta, C.; D’Emmanuele Di Villa Bianca, R.; Sorrentino, R.; Perfetto, B.; Tufano, M.A.; Miro, A.; La Rotonda, M.I.; Quaglia, F. Dry powders based on PLGA nanoparticles for pulmonary delivery of antibiotics: Modulation of encapsulation efficiency, release rate and lung deposition pattern by hydrophilic polymers. *J. Control. Release* **2012**, *157*, 149–159. <https://doi.org/10.1016/j.jconrel.2011.08.010>.
24. Zhang, L.; Pornpattananangkul, D.; Hu, C.-M.; Huang, C.-M. Development of Nanoparticles for Antimicrobial Drug Delivery. *Curr. Med. Chem.* **2010**, *17*, 585–594. <https://doi.org/10.2174/092986710790416290>.