

Household Indoor Concentration Levels of Nitrogen Dioxide (NO₂) and Ozone (O₃) in Eskişehir, Türkiye [†]

Seda Naz SARICA ^{1,*}, Özlem ÖZDEN ÜZMEZ ² and Semra MALKOÇ ³

¹ Affiliation 1; sedanazarica@gmail.com

² Affiliation 2; oozden@eskisehir.edu.tr

³ Affiliation 3; satik@eskisehir.edu.tr

* sedanazarica@gmail.com; Tel.: +90 537-669-89-13

[†] Presented at 5th International Electronic Conference on Atmospheric Sciences, 16–31 July 2022; Available online: <https://ecas2022.sciforum.net/>.

Abstract: The concentrations of nitrogen dioxide (NO₂) and ozone (O₃) were determined at five indoor environments (corridor, living area, bedroom, kitchen, bathroom) of two homes located in different regions of Eskişehir, Türkiye. Home 1 is located in the city center and urban residential area while Home 2 is located in a suburban area. In order to determine the indoor and outdoor concentration ratios of the pollutants (I/O), outdoor sampling was also carried out simultaneously with indoor sampling. Sampling studies were performed in one-day periods in four seasons by using passive sampling method. The indoor NO₂ concentrations varied between 8.80–124.18 µg/m³ while O₃ concentrations varied between 4.15–22.10 µg/m³. The highest NO₂ concentrations were determined in the kitchens both in two homes. This can be due to the intensive cooking activities carried out in the kitchens. The variation of O₃ concentrations in the measured indoor environments varied in homes. When the outdoor concentrations were examined, it was seen that NO₂ concentrations were higher in Home 1 and O₃ concentrations were higher in Home 2 in all seasons. This result is related to the location of the homes. The I/O ratios for NO₂ were generally > 1 for the kitchens. Also, all I/O ratios for NO₂ in Home 2 were found > 1 in autumn season. The I/O ratios for O₃ were found to be < 1 in both homes in all seasons. Seasonal variations of the pollutant concentration levels were also observed for indoor environments. Indoor NO₂ concentrations, especially in Home 1, and O₃ concentrations, especially in Home 2, were higher in spring and summer compared to other seasons. The reason for this is thought to be more active natural ventilation due to the warming of the weather in these seasons.

Academic Editor(s): Anthony Lupo

Published: 31 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Keywords: indoor air quality; home environment; NO₂; O₃



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Indoor air quality (IAQ) is a significant concern due to the amount of time spent indoors. Some studies show that people living in urban environments spend an average of 90% of their time indoors [1–3]. Indoor air can be more significantly polluted than outdoor air, even in the biggest and most industrialized cities. For many individuals, the health effects from exposure to indoor air pollution are higher than outside. IAQ is important because it has a significant impact on human health. Also, IAQ is little dependent on socioeconomic standing or education level but is highly impacted by variables such as personal products, furniture, and cleaning materials used [4,5]. These conditions may lead to the presence of biological, chemical and physical pollutants. Ventilation and other techniques are employed to enhance IAQ and minimize pollution; however, ventilation itself may also be considered as a source of contamination and exposure [6].

According to their physical states, pollutants exist in three basic states and have different qualities and impacts. These; solid, liquid and gaseous phases. Solid and liquid pollutants are regarded as particle matter. Major gaseous air pollutants are reactive, participate in photochemical processes or may react directly with material, plant or biological tissue. They occur as a combination in the atmosphere. Nitrogen dioxide (NO_2) and ozone (O_3) are among important air contaminants. These harmful gases are released into the atmosphere in large quantities from various sources all around the world.

Indoor NO_2 concentrations are directly impacted by high outside concentrations (for example, in cities with increased traffic density) [7]. The proximity of buildings to roads or the proximity of areas used for parking of motor vehicles to homes are among the major sources impacting indoor concentrations [8–10]. The most important indoor NO_2 sources are combustion processes (heaters, fireplaces and stoves). Accordingly, indoor NO_2 concentrations often exceed outside concentrations [11].

While O_3 is a layer in the upper atmosphere (stratosphere) that protects humans from ultraviolet (UV) radiation, it is a secondary air pollutant formed through a complex series of photochemical reactions that require reactive hydrocarbons, sunlight and nitrogen dioxide (NO_2) in the troposphere. Ozone concentrations are more likely to be lower in areas with motor vehicle exhaust and traffic, whereas NO_2 concentrations are higher in these areas. Indoor O_3 levels are usually lower when compared to outside levels. Common indoor ozone generators are laser printers, copiers and electrostatic air cleaners [12,13].

In this study, two homes having different outdoor locations and indoor characteristics were selected in Eskişehir, Turkey. Indoor NO_2 and O_3 sampling studies were performed in five various microenvironments such as corridor, living room, bedroom, kitchen and bathroom in each home. In addition to indoor sampling studies, outdoor samplings were also carried out simultaneously to evaluate the effect of outdoor air on indoor air and to determine indoor/outdoor (I/O) ratios for NO_2 and O_3 . Sampling studies were repeated in four seasons as winter, spring, summer and autumn.

2. Materials and Method

2.1. Characteristics of the Selected Homes

In this study, indoor and outdoor NO_2 and O_3 concentrations were determined in two selected homes in Eskişehir, Turkey. Some parameters such as the number of people living and working at homes, the size and locations of the homes, the number of the rooms were considered while choosing homes. 24-h sampling was carried out in the living room, bedroom, kitchen, bathroom and corridor at each home. The Google Earth view of the homes included in the study is shown in Figure 1.




Figure 1. The homes included in the study: (a) The location of the first home (Home 1) in Bahçelievler neighborhood. (b) The location of the second home (Home 2) in Aşağı Söğütönü neighborhood.

Specific characteristics of each home are given in Table 1. Also, the floor plans and sampling points of the selected homes are shown in Figure 2.

Table 1. Specific characteristics of the selected homes.

Home	The Size and Floor	Ventilation Type	Sampling Points	Location	Number of People Living in the Home
Home 1	85 m ² . On the 4th floor of a 4-storey building	Aspirator and natural ventilation	Corridor, living room, bedroom, kitchen, bathroom and outdoor	In an urban residential area. 50 m away from Eskişehir (E90-D200) belt highway. High population and traffic density	3 people live in this home. Two of them are actively going to work
Home 2	90 m ² . On the 1 st floor of a 4-storey building	Aspirator and natural ventilation	Corridor, living room, bedroom, kitchen, bathroom and outdoor	In a suburban area. 10 km from the city center. Low population and traffic density	4 people live in this home. One of them is actively working and one of them is a student




Figure 2. Floor plan and indoor–outdoor sampling points in (a) Home 1, (b) Home 2. 1: Corridor, 2: Living Room, 3: Bedroom, 4: Kitchen, 5: Bathroom, 6: Outdoor

2.2. Sampling Program and Method

In this study, the concentrations of inorganic gas pollutants were determined by passive sampling technique in different microenvironments (corridor, living room, bedroom, kitchen and bathroom) of 2 different homes. 24-h passive sampling studies were carried out in four seasons such as winter, spring, summer and autumn. The most suitable sampling point was determined for each indoor environment to best represent the air of the environment to be sampled, and passive samplers were put at the same points at each sampling period. Indoor and outdoor samplings were performed simultaneously.

The passive samplers used in the study were developed and validated by Eskişehir Technical University Environmental Engineering Department Air Quality Research Group [14–18]. The main parts of the passive sampler are: (1) sampler body, (2) stainless steel mesh barrier, (3) closed cap, (4) filter paper impregnated with specific solution and (5) fixer ring. The passive sampler body was manufactured from polytetrafluoroethylene (PTFE) for NO₂ and delrin for O₃. As collecting medium, Whatman GF/A glass fiber filter papers impregnated with 20% TEA (Triethanolamine) aqueous solution for NO₂ and aqueous solution containing 1% NaNO₂, 2% Na₂CO₃, and 2% glycerol for O₃ were used. Figure 3 shows an example of the NO₂ passive sampler and its parts [19].




Figure 3. Parts of the tailor-made passive sampler [19].

Extractions were performed before the analyses of the samples. Extractions of NO₂ filter papers were carried out in 10 mL ultrapure water (Milli Q) + 20 µL 35% H₂O₂ (Merck) solution, and O₃ filter papers were extracted in 10 mL ultrapure water (Milli Q) for 15 min. All samples were analyzed using Dionex ICS-1100 ion chromatography.

3. Results and Discussion

3.1. Indoor-Outdoor NO₂ and O₃ Concentrations

Table 2 shows determined indoor and outdoor NO₂ concentrations. For Home 1, during winter, spring, and autumn, the highest concentrations were measured in the kitchen (26.73 µg/m³, 51.37 µg/m³, and 54.00 µg/m³). The kitchen’s higher exposure to heavy traffic than other rooms, as well as ventilation provided by opening windows and also cooking and using a stove for breakfast and dinner is considered to be the cause of high concentrations measured in the kitchen at Home 1. For the summer season, the highest concentration was found in the living room (52.31 µg/m³). The reason for this high concentration can be that the living room is located closer to the street and is under the influence of traffic due to natural ventilation. For Home 2, the highest NO₂ concentrations were observed in the kitchen in all seasons. Seasonal variations in the indoor concentrations were observed for both homes. When the outdoor concentrations were examined, the highest concentration was measured for Home 1 in the winter season and for Home 2 in the autumn season. Since it is under the influence of more traffic due to its location, the outdoor concentrations measured for Home 1 were found to be higher in all seasons.

Table 2. Indoor-outdoor NO₂ concentrations (µg/m³) for selected homes for each season.

	Winter	Spring	Summer	Autumn
Home 1				
Corridor	14.27	31.98	39.58	38.27
Living Room	14.65	26.91	52.31	34.78
Bedroom	9.95	25.34	38.91	33.24
Kitchen	26.73	51.37	40.93	54.00
Bathroom	8.80	24.04	38.79	25.77
Outdoor	88.74	39.07	40.70	54.46
Home 2				
Corridor	16.88	22.68	12.19	51.91
Living Room	12.07	15.76	15.23	75.03
Bedroom	14.46	18.82	12.81	63.41
Kitchen	31.51	41.73	22.60	124.18
Bathroom	11.01	12.67	11.67	77.98
Outdoor	34.40	24.62	29.89	42.54

Table 3 shows O₃ concentrations measured indoors. For Home 1, during winter, spring and summer the highest concentrations were measured in the kitchen (6.35 µg/m³, 9.51 µg/m³, 11.34 µg/m³, respectively). The highest concentration was found in the bedroom in autumn (9.08 µg/m³). For Home 2, during winter, spring and summer the highest concentrations were measured in the kitchen (6.84 µg/m³, 11.01 µg/m³, 22.10 µg/m³, respectively). The highest concentration was found in the corridor in autumn (7.36 µg/m³). The most important source of indoor ozone pollution is the outdoor air. Also, the absence of major indoor ozone sources such as printers, photocopiers, and many other devices and appliances designed for indoor use (e.g., air cleaners) are the reasons for lower indoor than outdoor ozone concentrations. In addition, the age of a building and various housing aspects (carpeting, air conditioning, window fans, and window openings) have been significantly associated with indoor ozone levels. When the outdoor concentrations were examined, it was seen that the concentrations measured for Home 2, which is located far away from the city center, were higher than that of Home 1. Due to significantly elevated outdoor ozone concentrations during summer, summer indoor concentrations are typically elevated. The results of this study also support this situation.

Table 3. Indoor-outdoor O₃ concentrations (µg/m³) for selected homes for each season.

	Winter	Spring	Summer	Autumn
Home 1				
Corridor	5.81	5.90	7.12	6.88
Living Room	5.89	7.72	8.35	8.43
Bedroom	4.61	6.86	8.73	9.08
Kitchen	6.35	9.51	11.34	6.27
Bathroom	4.91	6.41	6.91	5.48
Outdoor	62.98	73.81	61.93	55.91
Home 2				
Corridor	4.63	9.74	11.29	7.36
Living Room	4.15	8.77	13.46	5.40
Bedroom	5.57	9.36	12.98	5.53
Kitchen	6.84	11.01	22.10	6.18
Bathroom	5.14	9.09	10.22	6.65
Outdoor	95.99	169.36	157.86	133.60

3.2. NO₂ and O₃ Indoor/Outdoor Concentration Ratios (I/O)

Table 4 shows indoor/outdoor NO₂ concentration ratios (I/O). All ratios were < 1 except for the ratio found for the kitchen in spring and summer and also the living room in the summer at Home 1. The reason for this is that Home 1 is in a high-traffic location, and the outside NO₂ concentration is higher in almost every season than in the indoor environment. Also, all I/O ratios for NO₂ in Home 2 were found > 1 in autumn season.

Table 4. Indoor/Outdoor ratios of NO₂ concentrations.

	Winter	Spring	Summer	Autumn
Home 1				
Corridor	0.16	0.82	0.97	0.70
Living Room	0.17	0.69	1.29	0.64
Bedroom	0.11	0.65	0.96	0.61
Kitchen	0.30	1.31	1.01	0.99
Bathroom	0.1	0.61	0.95	0.47

	Home 2			
Corridor	0.49	0.92	0.41	1.22
Living Room	0.35	0.64	0.51	1.76
Bedroom	0.42	0.76	0.43	1.49
Kitchen	0.92	1.70	0.76	2.92
Bathroom	0.32	0.51	0.39	1.83

Table 5 shows indoor/outdoor O₃ concentration ratios. For both homes, the I/O ratios for all seasons were < 1. This result indicates that there is no significant indoor source of O₃ in either home. The ratios found for Home 2 were lower than the ratios found for Home 1. The most important reason for this is the higher outdoor concentrations in Home 2.

Table 5. Indoor/Outdoor ratios of O₃ concentrations.

	Winter	Spring	Summer	Autumn
Home 1				
Corridor	0.09	0.08	0.11	0.12
Living Room	0.09	0.10	0.13	0.15
Bedroom	0.07	0.09	0.14	0.16
Kitchen	0.10	0.13	0.18	0.11
Bathroom	0.08	0.09	0.11	0.10
Home 2				
Corridor	0.05	0.06	0.07	0.06
Living Room	0.04	0.05	0.09	0.04
Bedroom	0.06	0.06	0.08	0.04
Kitchen	0.07	0.07	0.14	0.05
Bathroom	0.05	0.05	0.06	0.05

4. Conclusions

As a result of the pandemic, people have spent most of their time at homes. For this reason, many people have been exposed to air pollution more indoors than outdoors and have been affected more accordingly. This study was carried out during the COVID-19 pandemic period. In the study, simultaneous indoor-outdoor NO₂ and O₃ concentrations and also indoor/outdoor ratios (I/O) were determined for different home micro-environments in Eskişehir, Turkey. The highest NO₂ concentrations were found in the kitchens of both homes. This could be due to the intensive cooking activities that occur in the kitchens. O₃ concentrations measured in indoor environments varied depending on the season and also the environment. The reason for this can be the locations of the selected homes and the temperature changes in the seasons. In general, seasonal variations in the pollutant concentrations were observed in indoor environments. In spring and summer, indoor NO₂ concentrations, especially in Home 1, and O₃ concentrations, particularly in Home 2, were higher than in other seasons. It is thought that the reason for this is natural ventilation was more active during these seasons due to the warming of the weather. When the outdoor concentrations were evaluated, it was found that NO₂ concentrations were higher in Home 1 and O₃ concentrations were higher in Home 2 in all seasons. The reason for this is the locations of the homes and, accordingly, the level of being affected by the traffic density.

Author Contributions: The data presented in this study includes a part of the master thesis carried out by Seda Naz Sarıca in Eskişehir Technical University. Other authors also serve as thesis advisors (Özlem Özden Üzmez is 1st advisor, Semra Malkoç is 2nd advisor)

Funding: This research is funded by Eskişehir Technical University Research Fund (22ADP080).

Data Availability Statement: The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Conflicts of Interest: The authors declare that there is no conflict of interest.

References

1. Diffey, B.L. An overview analysis of the time people spend outdoors. *Br. J. Dermatol.* **2011**, *164*, 848–854.
2. Klepeis, N.E.; Nelson, W.C.; Ott, W.R.; Robinson, J.P.; Tsang, A.M.; Switzer, P.; Behar, J.V.; Hern, S.C.; Engelmann, W.H. The National Human Activity Pattern Survey (NHAPS): A resource for assessing exposure to environmental pollutants. *J. Expo. Sci. Environ. Epidemiol.* **2001**, *11*, 231–252.
3. World Health Organization. *Children's Health and the Environment*; WHO Training Package for the Health Sector; World Health Organization: Geneva, Switzerland, 2009. Available online: <http://www.who.int/ceh> (accessed on).
4. Fadeyi, M.O.; Weschler, C.J.; Tham, K.W.; Wu, W.Y.; Sultan, Z.M. Impact of human presence on secondary organic aerosols derived from ozone-initiated chemistry in a simulated office environment. *Environ. Sci. Technol.* **2013**, *47*, 3933–3941.
5. Weschler, C.J. Roles of the human occupant in indoor chemistry. *Indoor Air* **2016**, *26*, 6–24.
6. Seguel, J.M.; Merrill, R.; Seguel, D.; Campagna, A.C. Indoor air quality. *Am. J. Lifestyle Med.* **2017**, *11*, 284–295.
7. WHO. *WHO Guidelines for Indoor Air Quality: Selected Pollutants*; World Health Organization, Regional Office for Europe: Copenhagen, Denmark, 2010.
8. Janssen, N.A.H.; van Vliet, P.H.N.; Aarts, F.; Harssema, H.; Brunekreef, B. Assessment of exposure to traffic related air pollution of children attending schools near motorways. *Atmos. Environ.* **2001**, *35*, 3875–3884.
9. Kodama, Y.; Arashidani, K.; Tokui, N.; Kawamoto, T.; Matsuno, K.; Kunugita, N.; Minakawa, N. Environmental NO₂ concentration and exposure in daily life along main roads in Tokyo. *Environ. Res.* **2002**, *89*, 236–244.
10. Nakai, S.; Nitta, H.; Maeda, K. Respiratory health associated with exposure to automobile exhaust. II. Personal NO₂ exposure levels according to distance from the roadside. *J. Expo. Anal. Environ. Epidemiol.* **1995**, *5*, 125–136.
11. Kattan, M.; Gergen, P.J.; Eggleston, P.; Visness, C.M.; Mitchell, H.E. Health effects of indoor nitrogen dioxide and passive smoking on urban asthmatic children. *J. Allergy Clin. Immunol.* **2007**, *120*, 618–624.
12. Aviado, D.M.; Salem, H. Acute effects of air pollutants on the lungs. *Arch. Environ. Heal. Int. J.* **1968**, *16*, 903–907.
13. Lippmann, M. Health effects of tropospheric ozone. *Environ. Sci. Technol.* **1991**, *25*, 1954–1962.
14. Bozkurt, Z.; Üzmez, Ö.Ö.; Döğeroğlu, T.; Artun, G.; Gaga, E.O. Atmospheric concentrations of SO₂, NO₂, ozone and VOCs in Düzce, Turkey using passive air samplers: Sources, spatial and seasonal variations and health risk estimation. *Atmos. Pollut. Res.* **2018**, *9*, 1146–1156.
15. Özden, Ö.; Döğeroğlu, T. Performance evaluation of a tailor-made passive sampler for monitoring of tropospheric ozone. *Environ. Sci. Pollut. Res.* **2012**, *19*, 3200–3209.
16. Özden, Ö.; Dogeroglu, T. A field evaluation of a passive sampler for the simultaneous determination of NO₂ and SO₂ in anurban and rural area. *WSEAS Trans. Environ. Dev.* **2006**, *2*, 1008–1015.
17. Üzmez, Ö.Ö. *Atmosferik uçucu organik bileşiklerin ölçümü için pasif örnekleyici geliştirilmesi ve kullanımı*; Anadolu University: Eskişehir, Turkey, 2015.
18. Özden, Ö. *Hava kalitesinin monitorlanmasında pasif örnekleyicilerin kullanılması*; Anadolu University: Eskişehir, Turkey, 2015.
19. Üzmez, Ö.Ö. Atmospheric Concentrations of Inorganic Pollutants (NO₂, SO₂ and ozone) in Eskişehir: Spatial and Vertical Variations, Weekday-Weekend Differences. *Anadolu Univ. J. Sci. Technol. A-Appl. Sci. Eng.* **2018**, *19*, 523–535.