

Who, Where, When, and How Damages Forest – Challenges for Prediction and Control †

Valentyna Meshkova ^{1,2}

¹ Ukrainian Research Institute of Forestry & Forest Melioration, Pushkinska str. 86, 61024, Kharkiv, Ukraine; Valentynameshkova@gmail.com

² Czech University of Life Sciences Prague, Kamýčká 1283, 16500 Praha-Suchbát, Czech Republic; meshkova@fd.czu.cz

† Presented at the 3rd International Electronic Conference on *Forests* – Exploring New Discoveries and New Directions in Forests, 15 to 31 October 2022. Available online: <https://iecf2022.sciforum.net>

Abstract: The purpose of the research was an integration of published and own data obtained in the forests of Ukraine on the approaches to improving the prediction and control of forest insect pests. Climate change and anthropogenic activity bring changes in forest structure, its vulnerability to pests, pest species composition, and harm. The most dangerous alien species are *Leptoglossus occidentalis*, *Corythucha arcuata*, and *Agrilus planipennis*. Forecasts should consider contemporary pest composition, seasonal development, harm, the area and structure of vulnerable forests, and foliage mass depending on the natural zone, tree health condition, and additional damaging factors.

Keywords: climate change; foliage-browsing insects; alien pests; outbreaks; harmfulness; pest infestation prediction

1. Introduction

Terrestrial forest biodiversity is dominated by plants and the animals that eat them, the majority of which are insects. However, the certain traits of the pests might lead to a severe weakening and/or death of trees, a decrease in seeds harvesting, and reforestation (physiological harm), as well as deterioration in the timber quality (technical harm) or decrease in ornamental value (aesthetic harm) [1, 2]. Many species are pests not to the forest ecosystem, but to the forestry sector, which cannot obtain the expected forest products or seeds for growing planting stock and establishing plantations [3]. Harmfulness are highly variable among pest species the majority of which are insects [1]. Timely adequate protection of various forestry objects, such as fruits and seeds, planting stock, forests, and forest products, have to be based on the forecast data, in particular which species, where and when are able to cause environmentally and economically significant damage to forests and forest management [4]. Methods to surveying, and forecasting the spread of forest pests as well as assessment the injuriousness were developed in Ukraine for pests of nurseries and plantations, foliage-browsing and xylophagous insects [5–7]. The approaches cover the lists of harmful species, scenarios of the most common pests' development, the data on the frequency or severity of outbreaks in different regions, and the characteristics of forest stands that can be susceptible for insect pests' outbreaks. This information is used for seasonal (operational), annual (tactical), and long-term (strategic) predictions [4].

However, in recent decades, climate change resulted in an increase in air temperature, a decrease in precipitation, groundwater level and hydrothermal indices [8]. Such changes lead to shifts in natural zones, worsening conditions for the growth of the main forest-forming species and increasing their susceptibility to insect damage [4]. At the same time, the lengthening of the growing season is favorable for phytophagous insects [9, 10]. Because of the earlier beginning of insect development after hibernation, it is necessary to adjust the deadlines for the removal of harvested wood from the forest, the timing of the use of insecticides, and the pest population assessment.

Citation: Meshkova, V. Who, where, when, and how damages forest – challenges for prediction and control. *Environ. Sci. Proc.* **2022**, *4*, x. <https://doi.org/10.3390/xxxxx>

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Under climate change and the increasing anthropogenic impact, the tree species composition, age structure, and other forest characteristics have also changed [11, 12], which affected forest plots' preferences for pests and outbreaks' duration [13, 14]. Therefore, it is necessary to make adjustments to forecasts of different lead times.

The purpose of the research was to integrate published and own data obtained in the forests of Ukraine on the approaches to improving the outbreak prediction and control of forest insect pests.

2. Materials and Methods

The data sources were datasets from own research Ukrainian forests and the reference materials regarding the population dynamics of forest insect pests, their impact on forest health and outbreak prediction presented in the references.

3. Results and Discussion

The list of potentially dangerous pest species was unchanged for a long time, but the significance of some of them changed [15]. For example, in the late 1940's - early 1950's on the territory of Ukraine, the outbreaks of *Malacosoma neustria* (Linnaeus, 1758), *Euproctis chryorrhoea* (Linnaeus, 1758), *Phalera bucephala* (Linnaeus, 1758) were recorded, the distribution of which significantly decreased already in the 1970s [16].

Among phyllophagous insects in deciduous trees in Kharkiv city, the number of large-size species had significantly decreased for 50 years (1960's – 2010's), while the number of middle-size and small-size species had increased. Indifferent species prevailed at the beginning of this period and prodromal ones did at the end of it. The list of eruptive insects was supplemented by the open-living alien polyphagous species *Hyphantria cunea* (Drury, 1773) and several hidden monophagous species of Gracillariidae: native *Acrocercops brongniardella* (Fabricius, 1798) and *Phyllonorycter populifoliella* (Treitschke, 1833), and alien *Cameraria ohridella* Deschka & Dimic, 1986 and *Phyllonorycter issikii* (Kumata, 1963) [2]. Changes in the pest composition are largely associated with an increase in technogenic pollution, when smaller species, species with a hidden lifestyle and sucking mouthparts, and multivoltine species survive [17].

The extension of the growing season and warmer winters are favorable also for alien species that have spread beyond their natural ranges with planting stock, forest products, various cargoes, etc. [18] Some alien species are extinct in new regions, while others remain in low abundance, and we do not know about them. Some species did not affect the forest health in the area of origin but are pests in new places [19].

Among over 20 alien insects in the forests of Ukraine, the most dangerous are *Leptoglossus occidentalis* Heidemann, 1910, *Corythucha arcuata* (Say, 1832), and *Agrilus planipennis* Fairmaire, 1888 [20]. The feeding of *L. occidentalis* decreases the germination of pine seeds [20]. *C. arcuata* is found only in two regions but is moving northward and may pose a threat to oak stands [21]. *Agrilus planipennis* has spread over several years from the eastern border with Russia to the entire Luhansk and partly Kharkiv region damaging *Fraxnus excelsior* and *F. pennsylvanica*, and moving toward west [22]. Many species of insects that have recently been identified in the territory of Ukraine have previously penetrated into neighboring countries. Data obtained in other areas on distribution patterns, the list of host tree species, the timing of seasonal development, the relationship to temperature and humidity during feeding and wintering, and the presence of natural enemies of alien species can be used to predict the expected impact on the forest.

Incidence, duration, and severity of foliage browsing insect outbreaks in Ukraine were evaluated for 1947–1977 and 1978–2008 according to long-term statistical data [15]. The analysis made it possible the prediction of the regions with the highest threat as well as the years of the next outbreak beginning and collapse for certain pests and regions. However, these parameters can also vary with climate change, anthropogenic pressure, and changes in the stand structure.

The zoning of the territory according to the frequency or intensity of outbreaks of certain pest species is based on the long-term dynamics of the foci area in different regions and forestry enterprises [4]. It was shown that outbreak duration is connected with the pattern of seasonal development for different foliage-browsing insects. The longest is an outbreak of *T. viridana* (5 years on average) with the shortest period of vulnerability to regulating factors (larvae feeding period is about 30 days). The duration of the outbreak of *Diprion pini* with two generations per year is 4 years. Most outbreaks are longer in the East part of Ukraine [15]. Data on mean intervals between outbreaks of foliage-browsing insects in different regions of Ukraine and the durations of these outbreaks can be used for the estimation of years of the next outbreak beginning and collapsing.

Outbreaks of foliage-browsing insects often develop in the same stands at certain time intervals [4, 23, 24]. At the same time, the severity and duration of outbreaks depend on the stand structure, and the consequences of foliage damage depend, to a large extent, on the initial forest health. The forest subcompartments with the highest probability of foliage-browsing insects' foci may be recognized using scoring the characteristics of stands, which are presented in the forest inventory database. For each subcompartment, the type of forest site conditions, age, the relative density of stocking, and the proportion of host plants in the tree species composition are evaluated by points [4]. Additional information about the adjacent clear-cuts, burned areas and other types of plots increasing the risk of pest outbreaks is obtained using GIS technologies. According to the sum of the area of subcompartments with a high threat of certain pest outbreaks, the expected foci area in the forest enterprise is calculated and mapped [25].

Such a list of subcompartments is not permanent, because all mentioned forest parameters may change as a result of natural or anthropogenic factors. Thus, with the age of pine stands, the threat of outbreaks of *Neodiprion sertifer* (Geoffroy, 1785) decreases and the threat of *Dendrolimus pini* (Linnaeus, 1758) foci increases [4]. A decrease in the relative density of stocking is favorable for phytophages of various ecological groups, particularly foliage-browsing ones [11], and xylophages [26]. The change in the microclimate of certain forest subcompartments affects the seasonal development, hibernation, and survival of insects [4].

After a recent long-term outbreak of bark beetles in the pine forests of Polissya (Ukrainian forest zone), the area of stands with a high threat of foliage-browsing insect outbreaks has decreased [23]. It happened because bark beetles colonised primarily pure pine stands with a low relative density of stocking [26, 27, 28]. As the trees died in these stands, selective and then clear sanitary cuttings were carried out, which reduced the area of stands attractive to foliage-browsing insects. At the same time, the fragmentation of forests increased, which contributed to a greater risk of open forest edges being colonised by insects [14]. It means that the potential foci area of foliage-browsing insects should be regularly specified, at least after the next forest inventory.

The population density of foliage-browsing insects, at which it is advisable to use insecticides, depends on a foliage mass, which is directly related to tree health. Using the data on foliage phytomass of one tree of a certain height and diameter, depending on the natural zone [29] and feeding norms of the most common pests, the standards for calculating the maximum population density of these species have been developed considering the average class of tree health in the inspected stand [7]. Therefore, if the phytomass of a healthy tree is taken as a 1, then the phytomass of a weakened tree (2nd class of health condition) is equal to 0.8 of the phytomass of a healthy tree of this species, for the severely weakened tree (3rd class) it is 0.4, and for the drying tree (4th class) it is 0.16. Consequently, the population density of larvae to consume 100 % of foliage of the healthy tree is 1.25, 2.5, and 6.25 higher than for the trees of the 2nd, 3rd, and 4th classes of health condition [3].

Xylophages colonized the trees that are weakened by fire, windstorm, or anthropogenic impact (recreation, industrial pollution, forestry activities). Such foci collapse when the available substrate decreases [14]. The harmfulness of xylophagous species was evaluated using a scoring system that considered the effect of these insects on the health of

living trees and on timber quality [30]. However, some parameters for the evaluation of the harmfulness of the same insect species depend on the host tree, region, and additional factors that weaken trees [6, 31].

Optimal timing for pest detection, assessment, and control depends on insect seasonal development in a given region and year [4]. The patterns of seasonal development for foliage-browsing insects and possible changes with temperature increase were analysed considering hibernating stage and the presence of summer diapause [13]. Reliable relationships between the dates of stable transition of air temperature through 5, 10 and 15°C in spring and autumn, as well as the dependence of the seasonal development of insects with different patterns of seasonal development on the temperature, were described [4]. This made it possible to draw up a scheme and methodology for determining the feeding periods of such insects for planning their assessment and control. Such an approach will be actual in any climate change. For example, the treatment of forest against larvae of *Tortrix viridana*, *Operophtera brumata* (Linnaeus, 1758), *Lymantria dispar* (Linnaeus, 1758), *Euproctis chryorrhoea*, *Neodiprion sertifer*, *Panolis flammea* (Denis & Schiff., 1775), and *Dendrolimus pini* after hibernation should be carried out after a stable transition of average daily air temperature over 10°C.

3. Conclusions

Existing approaches to predicting the spread and development of forest pests remain relevant in the face of climate change and an increase in anthropogenic pressure.

However, our study demonstrates that the evaluation of insects' significance in forest weakening should take into account changes in the composition of local and alien species, the starting and ending dates of a growing season, the area of vulnerable forests, their tree composition, structure, fragmentation, and foliage mass.

Among alien species the biggest threat to forests we can expect from *Leptoglossus occidentalis*, *Corythucha arcuata*, and *Agrilus planipennis*.

Moreover, the criteria for insect harmfulness have to be revised depending on the natural zone, the mass of foliage in the tree, its health condition, and additional damaging factors which might be challenging for further research.

Funding: This research received no external funding

Conflicts of Interest: The author declares no conflict of interest.

References

1. Kautz, M., Meddens, A. J., Hall, R. J., & Arneith, A. Biotic disturbances in Northern Hemisphere forests – a synthesis of recent data, uncertainties and implications for forest monitoring and modeling. *Glob. Ecol. and Biogeogr.* 2017, 26(5), 533-552; <https://doi.org/10.1111/geb.12558>
2. Kardash, Ye. S. Changes in the complex of phyllophagous insects in deciduous trees of Kharkiv city for 50 years. *Baltic coastal zone* 2021, 24, 27–39. ISSN 2083-5485
3. Meshkova, V. Assessment and prediction of biotic risks in the forests of Ukraine. *Bucovina Forestieră* 2021, 21(1), 83-92, 2021. DOI: 10.4316/bf.2021.007
4. Meshkova, V. Seasonal development of the foliage browsing insects. *Novoe slovo*, 2009. Kharkov. ISBN:978-966-204-669-4.
5. Björkman, C., Bylund, H., Nilsson, U., Nordlander, G., Schroeder, M. Effects of new forest management on insect damage risk in a changing climate. In *Climate change and insect pests*. CABI, UK, 2015, pp. 248–266. ISBN 978-1-78064-378-6
6. Meshkova, V. Evaluation of injuriousness of stem insects in pine forest. *Sci. Bul. UNFU* 2017, 27(8). 101–104. DOI: <https://doi.org/10.15421/40270816>
7. *Methodical guidelines for a survey, assessment and forecasting of the spread of forest pests and diseases for the lowland part of Ukraine* / V.L. Meshkova ed. Kh., Planeta-print, 2020. 90 p. (in Ukrainian) ISBN 978-617-7897-00-1
8. Shvidenko, A., Buksha, I., Krakovska, S., Lakyda, P. Vulnerability of Ukrainian forests to climate change. *Sustainability* 2017, 9 (7), 1152. DOI: <https://doi.org/10.3390/su9071152>
9. Siitonen, J. *Ips acuminatus* kills pines in southern Finland. *Silva Fennica* 2014, 48 (4), article id 1145. DOI: <http://dx.doi.org/10.14214/sf.1145>
10. Andreieva, O. Y., Goychuk, A. F. Forest site conditions and the threat for insect outbreaks in the Scots pine stands of Polissya. *Folia Forestalia Polonica, Series A – Forestry* 2020, 62(4), 270–278. DOI: 10.2478/ffp-2020-0026

11. Andreieva, O. Yu. Change of attractiveness of stands of the state enterprise "Korostenske Forest & Hunting Enterprise" for foliage browsing insects. *For & For Melior* 2021, 138, 97–103. <https://doi.org/10.33220/1026-3365.138.2021.97> (in Ukrainian).
12. Kosunen, M., Kantola, T., Starr, M., Blomqvist, M., Talvitie, M., Lyytikäinen-Saarenmaa, P. Influence of soil and topography on defoliation intensity during an extended outbreak of the common pine sawfly (*Diprion pini* L.). *IForest* 2017, 10, 164–171. DOI: 10.3832/ifor2069-009.
13. Meshkova, V. Proc and cons of Climate Change for Forest Phytophagous insects. *Proc. of the 1st Intern Electr Conf on Entomol*, 1–15 July 2021, MDPI: Basel, Switzerland, doi:10.3390/IECE-10373
14. Meshkova, V. The Lessons of Scots Pine Forest Decline in Ukraine. *Environ. Sci. Proc.* 2021, 3(1), 28. <https://doi.org/10.3390/IECF2020-07990>
15. Meshkova, V., Davydenko, K. Foliage browsing insects outbreaks in Ukraine before and after global warming. *Ber. Freiburger Forst. Forsch.* 2011, 89, 18-25. ISSN 1436–1566. https://www.fva-bw.de/fileadmin/publikationen/fff_bericht/fff_h_89.pdf
16. Meshkova, V. Foliage-browsing Lepidoptera (Insecta) in deciduous forests of Ukraine for the last 70 years. *Proc. FASU* 2021, 22, 173-179. DOI: <https://doi.org/10.15421/412115>
17. Jactel, H., Koricheva, J., & Castagneyrol, B. Responses of forest insect pests to climate change: not so simple. *Current opinion in insect science* 2019, 35, 103-108. <https://doi.org/10.1016/j.cois.2019.07.010>
18. Sandvik, H., Hilmo, O., Henriksen, S., Elven, R., Åsen, P. A., Hegre, H., ... & Westergaard, K. B. Alien species in Norway: Results from quantitative ecological impact assessments. *Ecol. Solut. and Evid.* 2020, 1(1), e12006. DOI: 10.1002/2688-8319.12006
19. Liebhold, A. M., Brockerhoff, E. G., Kalisz, S., Nuñez, M. A., Wardle, D. A., & Wingfield, M. J. Biological invasions in forest ecosystems. *Bio. Invas.* 2017, 19(11), 3437-3458. DOI 10.1007/s10530-017-1458-5
20. Meshkova, V. Alien phytophagous insects in forest and urban stands of Ukraine. *Buc. For.* 2022, 22(1): 29-40 <http://www.bucovina-forestiera.ro/index.php/bf/article/view/300/> DOI: 10.4316/bf.2022.004
21. Meshkova, V., Nazarenko, S., Glod, O. The first data on the study of *Corythucha arcuata* (Say, 1832) (Heteroptera: Tingidae) in Kherson region of Ukraine. *Proc. FASU* 2020, 21, 30–38. <https://doi.org/10.15421/412023>
22. Davydenko, K., Skrylnik, Y., Borysenko, O., Menkis, A., Vysotska, N., Meshkova, V., Olson, A., Elfstrand, M., Vasaitis, R. Invasion of Emerald ash borer *Agrilus planipennis* and ash dieback pathogen *Hymenoscyphus fraxineus* in Ukraine—A concerted action. *Forests* 2022, 13, 789. <https://doi.org/10.3390/f13050789>
23. Andreieva, O.Y., Boliukh, O.G. The outbreaks of Common Pine Sawfly (*Diprion pini* L.) in the forest fund of Zhytomyr region. *Sci. Bull. UNFU* 2019, 29 (7), 84–89. DOI: <https://doi.org/10.15421/40290717> (in Ukrainian)
24. Meshkova, V., Nazarenko, S., Koliienkina, M. *Diprion pini* L. (Hymenoptera, Symphyta, Diprionidae) population dynamics in the Low Dnieper region. *Folia Forestalia Polonica, Series A – Forestry* 2019, 61 (1), 22–29. DOI: <https://doi.org/10.2478/ffp-2019-0002>.
25. Meshkova, V., Borysenko, O. GIS-based prediction of the foliage browsing insects' outbreaks in the pine stands of the SE "Kreminske FHE". *Proc. FASU* 2017, 15, 12–18. DOI: <https://doi.org/10.15421/411714>
26. Meshkova, V. L., Borysenko, O. I., & Pryhornytskyi, V. I. Forest site conditions and other features of Scots pine stands favorable for bark beetles. *Proc. FASU* 2018, 16, 106-114; <https://doi.org/10.15421/411812>
27. Andreieva, O.Yu., Vyshnevskiy, A.V., Boliukh, S.V. Population dynamics of bark beetles in the pine forests of Zhytomyr region. *Sci. Bull. UNFU*, 2019, 29(8), 31–35. DOI: <https://doi.org/10.36930/40290803> (in Ukrainian)
28. Meshkova, V., & Bobrov, I. Parameters of *Pinus sylvestris* health condition and *Ips acuminatus* population in pure and mixed stands of Sumy region. *Proc. FASU* 2020, 20, 131-140. DOI: <https://doi.org/10.15421/412012>
29. Lakyda, P., Vasylyshyn, R., Lashenko, A., & Terentiev, A. *Standards of evaluation of components of aboveground trees biomass of trees of the main forest-forming species of Ukraine*. ECO-inform, Kyiv, 2011, 186 p. (in Ukrainian).
30. Mozolevskaya, E. G. Evaluation of stem pests injuriousness. *Sci. pap. of MFTI* 1974, 65: 124–132 (In Russian).
31. Skrylnik, Yu., Koshelyaeva, Y., Meshkova, V. Harmfulness of xylophagous insects for silver birch (*Betula pendula* Roth.) in the left-bank forest-steppe of Ukraine. *Fol. For. Pol., Ser A – Forestry* 2019, 61 (3), 161–175. DOI: <https://doi.org/10.2478/ffp-2019-0016>.