

Proceeding Paper

Co-Axial Wet-Spun Fibers: An Innovative Strategy for Chronic Wound Healing Applications [†]

Catarina S. Miranda ^{1,*}, A. Francisca G. Silva ², Sílvia M. M. A. Pereira-Lima ², Susana P. G. Costa ², Natália C. Homem ³ and Helena P. Felgueiras ¹

1 Centre for Textile Science and Technology (2C2T), University of Minho, 4710-057 Braga, Portugal; e-mail@e-mail.com

2 Centre of Chemistry, University of Minho, 4710-057 Braga, Portugal; e-mail@e-mail.com (A.F.G.S.); e-mail@e-mail.com (S.M.M.A.P.-L.); e-mail@e-mail.com (S.P.G.C.)

3 Digital Transformation CoLab (DTx), University of Minho, 4710-057 Braga, Portugal; e-mail@e-mail.com

* Correspondence: catarina.miranda@2c2t.uminho.pt

[†] Presented at the 3rd International Electronic Conference on Applied Sciences; Available online: <https://asec2022.sciforum.net/>.

Citation: Miranda, C.S.; Silva, A.F.G.; Pereira-Lima, S.M.M.A.; Costa, S.P.G.; Homem, N.C.; Felgueiras, H.P. Co-Axial Wet-Spun Fibers: An Innovative Strategy for Chronic Wound Healing Applications. *Eng. Proc.* **2022**, *4*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s):

Published: 1 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Chronic wounds (CW) are a worldwide concern, causing serious strives on the health and quality of patients' life. In CW, human neutrophil elastase (HNE) enzyme gets highly expressed during inflammation, reaching abnormally elevated concentrations. Additionally, prevalence of *Staphylococcus aureus*-induced infections remains very high and difficult to treat. Considering these phenomena, a drug delivery system made of co-axial wet-spun fibers, loaded with the tetrapeptide Ala-Ala-Pro-Val (AAPV, a known inhibitor of HNE activity) and N-carboxymethyl chitosan (NCCM, responsive to neutral-basic pH's, characteristic of CW and endowed with antibacterial features), was proposed. AAPV was synthesized by solid-phase peptide synthesis, whereas NCCM was synthesized from low molecular weight chitosan in a chloroacetic acid mixture. HNE inhibition tests were conducted to establish the AAPV IC₅₀ in 1.50 µg/mL and the NCCM minimum bactericidal concentration (MBC) against *Staphylococcus aureus* in 6.40 mg/mL. These determinations were used to establish fiber loading amounts. Core-shell structures were produced with 10% *w/v* polycaprolactone (PCL) at the core and 2% *w/v* sodium alginate (SA) solution at the shell. NCCM was mixed with SA at 2 × MBC so neutral-basic pH-triggered solubility (characteristic of CW) would allow pores to be opened in the outer layer for accessing the core, where AAPV was combined with PCL. Fourier-transform infrared spectroscopy and brightfield microscopy were used to confirm the presence of the four components on the fibers and the co-axial architecture, respectively. Fibers presented maximum elongations of over 100%. Release kinetics studies conducted via UV-visible absorption spectroscopy mapped NCCM liberation overtime but were incapable of detecting AAPV, since polymer degradation masked AAPV absorption peaks. Time-kill kinetics studies against *S. aureus* demonstrated the effectiveness of NCCM in eliminating this bacterium, particularly after 6 h of incubation. On its turn, AAPV guaranteed HNE inhibition. Data demonstrated the potential of SA-NCCM-PCL-AAPV co-axial systems to work as stepwise, pH-triggered delivery platforms.

Keywords: wound healing; drug delivery; therapeutic tetrapeptides