

Structural and Functional Annotation of an Uncharacterized Protein of *Helicobacter pylori*: An In-Silico Approach [†]

Md. Morshed Alam ^{1,2}, Abu Saim Mohammad Saikat ^{2,3,*} and Md. Ekhlas Uddin ^{2,4}

¹ Department of Biochemistry & Molecular Biology, Primeasia University, Dhaka 1213, Bangladesh; morshedariyan35@gmail.com

² Department of Computational Biology and Bioinformatics, Advanced Bioscience Center for Collaborative Research (ABSCCR), Rajshahi 6250, Bangladesh; ekhlasbt03@gmail.com

³ Department of Biochemistry and Molecular Biology, Life Science Faculty, Bangabandhu Sheikh Mujibur Rahman Science and Technology University, Gopalganj 8100, Bangladesh

⁴ Department of Biochemistry and Molecular Biology, Gono Bishwabidyalay, Dhaka 1344, Bangladesh

* Correspondence: asmsaikat.bmb@gmail.com

[†] Presented at the 2nd International Electronic Conference on Processes: Process Engineering—Current State and Future Trends (ECP 2023), 17–31 May 2023; Available online: <https://ecp2023.sciforum.net/>.

Abstract: *Helicobacter pylori* (*H. pylori*) is a gram-negative, slow-growing bacterium, microaerophilic, and a unique gastric pathogen causing chronic inflammation in the gastric mucosa with a possibility of developing gastric cancer has one-third of its proteins still uncharacterized. In this study, computational analysis has been done on the structural, functional, and epitopic characteristics of a hypothetical protein (HP) from *H. pylori* named HPF63-1454. The model prediction and primary, secondary, and three-dimensional structures of the chosen HP were built. The newly created model appeared to have high quality once refinement and structure validation was completed. The anticipated tertiary structure was assessed using the Swiss Model assessment. The best materials are chosen using structural analyses considering data from the Z scores, Swiss Interactive Workplace, and Ramachandran plot statistics. This investigation aimed to determine the importance of the *H. pylori* protein HPF63-1454. Therefore, this research will increase our understanding of pathophysiology and allow us to target the protein complex specifically. *H. pylori* infections are now treated with readily available antibiotics and acid-suppressing medications, but the species soon established itself as a hardy pest. Long-term and taxing antibiotic therapy for *H. pylori* infection is increasingly beginning to fail, necessitating novel, creative therapeutic modalities.

Keywords: *Helicobacter pylori*; uncharacterized protein; Ramachandran plot; pathogenesis; bacteria

Citation: Alam, M.M.; Saikat, A.S.M.; Uddin, M.E. Structural and Functional Annotation of an Uncharacterized Protein of *Helicobacter pylori*: An In-Silico Approach. *Eng. Proc.* **2023**, *5*, x. <https://doi.org/10.3390/xxxxx>
Published: 17 May 2023



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Helicobacter pylori (*H. pylori*) is a microaerophilic, gram-negative, spiral-shaped, slow-growing bacterium. This ubiquitous bacterium infects a substantial portion of the global populace. Australian scientists and doctors Robin Warren and Barry Marshall discovered in 1982 [1,2], an etiological agent for peptic ulcer and gastric inflammatory illnesses. A brand-new genus of bacteria that lives at the mucosa-to-lumen transition in the gut. They have characteristics required for colonization of the juxta-mucosal environment mucus, including microaerophilia, spiral shape, and urease positivity [3,4].

The primary cause of gastritis, gastroduodenal ulcer disease, adenocarcinoma (gastric), and mucosa-associated tissue lymphoma is *H. pylori* infection. *H. pylori* infection has drawn attention as an etiological cofactor in this cancerous illness since it is linked to an elevated risk of gastric adenocarcinoma [4]. It may disrupt a wide range of biological processes inside and outside the stomach, which could impact or determine the development of various disorders external to the stomach [5]. The most significant single risk factor for stomach malignancies is *H. pylori*-induced gastritis, but only a tiny percentage of infected

people acquire a malignancy. Host responses, strain-specific bacterial components, and host-microbe interactions affect cancer risk. In many developing nations, the frequency among middle-aged adults is over 80%, compared to 20 to 50% in industrialized countries. The disease is acquired primarily among early childhood families and is brought on by oral consumption of the bacteria [6–8].

This uncharacterized protein could be helpful as a pharmacological target and marker if annotated. According to the findings of various bioinformatics tools, software, and databases, the selected protein plays a significant part in the pathogenesis of *H. pylori*.

2. Materials and Methods

2.1. Protein Sequence Retrieval

The selected protein's amino acid (aa) sequence was obtained with the accession number BAW58640 (version: BAW58640.1) in the FASTA format from the NCBI [9,10].

2.2. Protein Characterization

The protein characterization was performed using The ExPASy server to measure the aa sequence composition, instability index, aliphatic index, GRAVY, and extinction coefficients. Moreover, we used the elevation of the theoretical isoelectric point (pI) for the HPF63_1454 protein [11,12].

2.3. Functional Annotation Anticipation

We utilized the conserved domain (CD) prediction in the protein of HPF63_1454. We predicted protein motif determination using the CD search functionality on the NCBI platform and the ScanProsite tool of the ExPASy server, respectively [13–15].

2.4. Prediction of Secondary Structure

The prediction of the secondary structure of the selected protein HPF63_1454 was anticipated using SPIRED V.4.0 software (server-based), and the SOPMA framework was used for the element anticipation [16,17].

2.5. Three-Dimensional Structure Prediction and Validation

The tertiary (3D) structure of protein predicted by the SWISS-MODEL and the structural quality assessment studies were performed using the PROCHECK Sever of the (SAVES v.6.0) [18–20].

3. Results and Discussion

3.1. Protein Sequence Retrieval

Protein sequences determine the fundamental components of biological structure and function. The NCBI protein sequence database is a protein sequence database collected from multiple sources, including RefSeq and Gene-Bank. The protein obtained from the NCBI protein database with the accession number BAW58640 (version: BAW41740.1) is present in the locus BAW58640 containing 451 aa.

3.2. Protein Characterization

The physicochemical characteristics of the selected protein were demonstrated using the ProtParam tool. The protein HPF63_1454 contains a 451 aa long sequence with the protein having a molecular weight (MW) of 53125.98 Da. The protein consists of 451 aa, where Leu (56) was the most abundant amino acid, followed by Lys (54), Glu (41), Asp (34), Ile (29), Ser (29), Asn (23), Thr (23), Tyr (21), Ala (20), Phe (18), Val (16), Arg (15), Pro (13), Gln (12), Gly (11), His (11), Met (10), Trp (8), and Cys (7) (Table 1).

Table 1. Amino acid composition of the selected protein.

Amino Acid	Quantity	Percentage
Ala (A)	20	4.4%
Arg (R)	15	3.3%
Asn (N)	23	5.1%
Asp (D)	34	7.5%
Cys (C)	7	1.6%
Gln (Q)	12	2.7%
Glu (E)	41	9.1%
Gly (G)	11	2.4%
His (H)	11	2.4%
Ile (I)	29	6.4%
Leu (L)	56	12.4%
Lys (K)	54	12.0%
Met (M)	10	2.2%
Phe (F)	18	4.0%
Pro (P)	13	2.9%
Ser (S)	29	6.4%
Thr (T)	23	5.1%
Trp (W)	8	1.8%
Tyr (Y)	21	4.7%

3.3. Functional Annotation Prediction

The NCBI CDD tool anticipated two different domains, including DUF1524 and COG1479. The DUF1524 domain is predicted to have a conserved HXXP motif associated with endonuclease activities (Pfam accession: pfam07510; interval: 290–444; E-value: 2.17×10^{-18}). Endonucleases cleave the phosphodiester bond present in nucleotide monomers. When cutting DNA, some do not care about the sequence, while others only do it at particular nucleotide sequences [21,22].

Moreover, The COG1479 domain contains differences between two nucleases (Hnh nuclease and ParB nuclease), which is (Pfam accession: COG1479; interval: 33–188; E-value: 1.35×10^{-6}). Its identical motif is noticed in protein families. Its rules as endonucleases. The actual severing of DNA strands by the nuclease domain can result in single-stranded or double-stranded breaks.

3.4. Secondary Structure Prediction

Secondary structure decisiveness of the chosen protein consensus projection from multiple alignments significantly improves protein secondary structure prediction. The secondary-structural component of the protein (HPF63_1454) was a prediction made by the SOPMA software where the alpha helix (Hh), extended strand (Ee), beta-turn (Tt), random coil (Cc) was 287 (63.64%), 35 (7.76%), 7 (1.55%), and 122 (27.05%), respectively.

3.5. Three-Dimensional Structure Prediction and Validation of the Selected Protein

The three-dimensional architecture of the desired protein was anticipated using the SWISS-Model program [23,24]. Moreover, the PROCHECK program of the SAVES methods was utilized for structural quality assessment of the modeled protein, where the arrangement of the ψ angle and the ϕ angle is shown. The protein's modeled tertiary structure was validated by the residues in the most preferred areas, which were swallowed by 92.0% of the protein (BAW58640.1) (Figure 1).




Figure 1. (a) Three-dimensional structure modeled by the SWISS-Model tool, and (b) The PROCHECK tool validated the Ramachandran plot statistics of the modeled protein structure.

4. Conclusions

The bacterium *Helicobacter pylori* increase the risk of developing gastric cancer. Although it is a highly deadly condition, the development of malignancy allows for a method to detect those at an increased risk daily. However, gastric infection with this bacterium is widespread, and most infected people do not go on to catch the disease. This study used the anticipated ligand-binding active sites in *Helicobacter pylori* to demonstrate the structural and functional annotation of HPF63-1454. They analyze the protein structure that led to the placement of amino acid sequences in the desired location. The physicochemical characteristics and available enrichment estimation are steady in comprehending protein operations. In most arrangements, alpha-helix, random coils, extended strands, and beta twists were found to predominate, as confirmed by the secondary assumption and assessment structures. The findings of this study may contribute to the modulation of new target identification and drug discovery against the protein strain to reduce infections caused by the bacterium.

Author Contributions: Conceptualization, A.S.M.S.; methodology, A.S.M.S; software, M.M.A. and A.S.M.S; validation, A.S.M.S.; formal analysis, A.S.M.S; investigation, A.S.M.S; resources, A.S.M.S; data curation, M.M.A. and A.S.M.S; writing—original draft preparation, M.M.A.; writing—review and editing, A.S.M.S. and M.E.U.; visualization, M.M.A. and A.S.M.S; supervision, A.S.M.S; project administration, A.S.M.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Upon request to corresponding author.

Acknowledgments: The authors want to acknowledge the Department of Computational Biology and Bioinformatics, Advanced Bioscience Center for Collaborative Research (ABSCCR), Rajshahi 6250, Bangladesh, for supporting this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Go, M.F. Review article: Natural history and epidemiology of *Helicobacter pylori* infection. *Aliment. Pharmacol. Ther.* **2002**, *16* (Suppl. S1), 3–15. <https://doi.org/10.1046/j.1365-2036.2002.0160s1003.x>.
2. Zhang, C.; Yamada, N.; Wu, Y.L.; Wen, M.; Matsuhisa, T.; Matsukura, N. *Helicobacter pylori* infection, glandular atrophy and intestinal metaplasia in superficial gastritis, gastric erosion, erosive gastritis, gastric ulcer and early gastric cancer. *World J. Gastroenterol.* **2005**, *11*, 791–796. <https://doi.org/10.3748/wjg.v11.i6.791>.
3. Malfertheiner, P.; Megraud, F.; O'Morain, C.; Bazzoli, F.; El-Omar, E.; Graham, D.; Hunt, R.; Rokkas, T.; Vakil, N.; Kuipers, E.J. Current concepts in the management of *Helicobacter pylori* infection: The Maastricht III Consensus Report. *Gut* **2007**, *56*, 772–781. <https://doi.org/10.1136/gut.2006.101634>.
4. Parsonnet, J.; Friedman, G.D.; Vandersteen, D.P.; Chang, Y.; Vogelstein, J.H.; Orentreich, N.; Sibley, R.K. *Helicobacter pylori* infection and the risk of gastric carcinoma. *N. Engl. J. Med.* **1991**, *325*, 1127–1131. <https://doi.org/10.1056/NEJM199110173251603>.
5. Franceschi, F.; Gasbarrini, A.; Polyzos, S.A.; Kountouras, J. Extragastric Diseases and *Helicobacter pylori*. *Helicobacter* **2015**, *20* (Suppl. S1), 40–46. <https://doi.org/10.1111/hel.12256>.
6. Polk, D.B.; Peek, R.M., Jr. *Helicobacter pylori*: Gastric cancer and beyond. *Nature reviews. Cancer* **2010**, *10*, 403–414. <https://doi.org/10.1038/nrc2857>.
7. Feldman, R.A. Epidemiologic observations and open questions about disease and infection caused by *H. pylori*. In *H. pylori Molecular and Cellular Biology*; Horizon Scientific: Wymondham, UK, 2007; pp. 29–51.
8. Rowland, M.; Kumar, D.; Daly, L.; O'connor, P.; Vaughan, D.; Drumm, B. Low rates of *Helicobacter pylori* reinfection in children. *Gastroenterology* **1999**, *117*, 336–341.
9. NCBI Resource Coordinators Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res.* **2018**, *46*, D8–D13. <https://doi.org/10.1093/nar/gkx1095>.
10. Saikat, A.S.M.; Paul, A.K.; Dey, D.; Das, R.C.; Das, M.C. In-Silico Approaches for Molecular Characterization and Structure-Based Functional Annotation of the Matrix Protein from *Nipah henipavirus*. *Chem. Proc.* **2022**, *12*, 21. <https://doi.org/10.3390/ecsoc-26-13522>.
11. Wilkins, M.R.; Gasteiger, E.; Bairoch, A.; Sanchez, J.C.; Williams, K.L.; Appel, R.D.; Hochstrasser, D.F. Protein identification and analysis tools in the ExPASy server. *Methods Mol. Biol.* **1999**, *112*, 531–552. <https://doi.org/10.1385/1-59259-584-7:531>.
12. Saikat, A.S.M.; Islam, R.; Mahmud, S.; Imran, M.A.S.; Alam, M.S.; Masud, M.H.; Uddin, M.E. Structural and Functional Annotation of Uncharacterized Protein NCGM946K2_146 of *Mycobacterium Tuberculosis*: An In-Silico Approach. *Proceedings* **2020**, *66*, 13. <https://doi.org/10.3390/proceedings2020066013>.
13. Lu, S.; Wang, J.; Chitsaz, F.; Derbyshire, M.K.; Geer, R.C.; Gonzales, N.R.; Gwadz, M.; Hurwitz, D.I.; Marchler, G.H.; Song, J.S.; et al. CDD/SPARCLE: The conserved domain database in 2020. *Nucleic Acids Res.* **2020**, *48*, D265–D268. <https://doi.org/10.1093/nar/gkz991>.
14. de Castro, E.; Sigrist, C.J.; Gattiker, A.; Bulliard, V.; Langendijk-Genevaux, P.S.; Gasteiger, E.; Bairoch, A.; Hulo, N. ScanProsite: Detection of PROSITE signature matches and ProRule-associated functional and structural residues in proteins. *Nucleic Acids Res.* **2006**, *34*, W362–W365. <https://doi.org/10.1093/nar/gkl124>.
15. Saikat, A.S.M.; Uddin, M.E.; Ahmad, T.; Mahmud, S.; Imran, M.A.S.; Ahmed, S.; Alyami, S.A.; Moni, M.A. Structural and Functional Elucidation of IF-3 Protein of *Chloroflexus aurantiacus* Involved in Protein Biosynthesis: An In Silico Approach. *BioMed Res. Int.* **2021**, *2021*, 9050026. <https://doi.org/10.1155/2021/9050026>.
16. Combet, C.; Blanchet, C.; Geourjon, C.; Deléage, G. NPS@: Network protein sequence analysis. *Trends Biochem. Sci.* **2000**, *25*, 147–150. [https://doi.org/10.1016/s0968-0004\(99\)01540-6](https://doi.org/10.1016/s0968-0004(99)01540-6).
17. Saikat, A.S.M.; Islam, R.; Mahmud, S.; Imran, M.A.S.; Alam, M.S.; Masud, M.H.; Uddin, M.E. Structural and Functional Annotation of Uncharacterized Protein NCGM946K2_146 of *Mycobacterium Tuberculosis*: An In-Silico Approach. *Proceedings* **2020**, *66*, 13. <https://doi.org/10.3390/proceedings2020066013>.
18. Waterhouse, A.; Bertoni, M.; Bienert, S.; Studer, G.; Tauriello, G.; Gumienny, R.; Heer, F.T.; de Beer, T.A.P.; Rempfer, C.; Bordoli, L.; et al. SWISS-MODEL: Homology modelling of protein structures and complexes. *Nucleic Acids Res.* **2018**, *46*, W296–W303. <https://doi.org/10.1093/nar/gky427>.
19. Laskowski, R.A.; MacArthur, M.W.; Moss, D.S.; Thornton, J.M. PROCHECK: A program to check the stereochemical quality of protein structures. *J. Appl. Crystallogr.* **1993**, *26*, 283–291.
20. Saikat, A.S.M.; Ripon, A. Structure Prediction, Characterization, and Functional Annotation of Uncharacterized Protein BCRIVMBC126_02492 of *Bacillus cereus*: An In Silico Approach. *Am. J. Pure Appl. Biosci.* **2020**, *2*, 104–111. <https://doi.org/10.34104/ajpab.020.01040111>.
21. Nichols, N.M. Endonucleases. In *Current Protocols in Molecular Biology*; John Wiley & Sons: Hoboken, NJ, USA, 2011; Chapter 3, Unit3.12. <https://doi.org/10.1002/0471142727.mb0312s93>.
22. Saikat, A.S.M.; Das, R.C.; Das, M.C. Computational Approaches for Structure-Based Molecular Characterization and Functional Annotation of the Fusion Protein of *Nipah henipavirus*. *Chem. Proc.* **2022**, *12*, 32. <https://doi.org/10.3390/ecsoc-2613530>.
23. Saikat, A.S.M. An In Silico Approach for Potential Natural Compounds as Inhibitors of Protein CDK1/Cks2. *Chem. Proc.* **2022**, *8*, 5. <https://doi.org/10.3390/ecsoc-25-11721>.
24. Tuncbag, N.; Gursoy, A.; Keskin, O. Prediction of protein-protein interactions: Unifying evolution and structure at protein interfaces. *Phys. Biol.* **2011**, *8*, 035006. <https://doi.org/10.1088/1478-3975/8/3/035006>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.