

# Comparative Evaluation of Antioxidant Activities of Flours from Durum Wheat Varieties <sup>†</sup>

Ioanna S. Kosma <sup>1</sup>, Afroditi Michalaki <sup>2</sup>, Ioannis Geraris Kartelias <sup>2</sup> and Haralabos C. Karantonis <sup>2,\*</sup>

<sup>1</sup> Laboratory of Food Chemistry, Department of Chemistry, University of Ioannina, 45110 Ioannina, Greece; i.kosma@uoi.gr

<sup>2</sup> Laboratory of Food Chemistry and of Technology and Quality of Animal Origin Food, Department of Food Science and Nutrition, School of the Environment, University of the Aegean, 81400 Myrina, Lemnos, Greece; fnsd20011@fns.aegean.gr (A.M.); fnsd21002@fns.aegean.gr (I.G.K.)

\* Correspondence: chkarantonis@aegean.gr; Tel.: +30-225-408-3111

<sup>†</sup> Presented at the 4th International Electronic Conference on Foods, 15–30 October 2023; Available online: <https://foods2023.sciforum.net/>.

**Abstract:** Antioxidants are known to play a crucial role in maintaining cellular health by neutralizing harmful free radicals. Among various dietary sources of antioxidants, wheat-based products, particularly flours, have gained significant attention due to their potential health benefits. Durum wheat, a widely cultivated species, serves as a primary ingredient in numerous food products. However, limited research has been conducted to assess the antioxidant activity of flours obtained from durum wheat varieties. In this study, we aimed to comparatively evaluate the antioxidant and antiplatelet potential of flours from 22 selected durum wheat varieties cultivated in Greece. We focused on three major parameters for antioxidant activity measurement: total phenolic content, 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity, and ferric reducing antioxidant power (FRAP). The total phenolic content was determined using the Folin-Ciocalteu method, while the DPPH and FRAP assays were employed to assess the ability of flours to scavenge free radicals and reduce ferric ions, respectively. Antiplatelet activity was evaluated by the platelet activating factor inhibition assay (PAF) in platelet rich plasma. Analysis of the data revealed notable differences in total phenolics, antioxidant and antiplatelet activities among the tested samples. The total phenolic content ranged from 624.0 ± 3.5 to 950.0 ± 5.3 µg of gallic acid equivalent/g flour with the variety Zeta E having the highest content. Antioxidant activities based on the DPPH and FRAP assays ranged from 0.56 ± 0.02 to 2.26 ± 0.08 and 1.93 ± 0.02 to 3.65 ± 0.03 µmol of trolox equivalent/g flour respectively with the varieties Marco Aurelio and Zeta E exerting the highest antioxidant activities in DPPH and FRAP, respectively. In addition, the IC<sub>50</sub> values for antiplatelet activity ranged from 0.72 ± 0.21 to 3.06 ± 0.17 in mg of flour, with the variety of Zoi exhibiting the highest antiplatelet activity. Overall, this comparative evaluation highlights the differences of antioxidant and antiplatelet activities among flours obtained from 22 different durum wheat varieties cultivated in Greece. The results from this study aid in the selection of wheat varieties with superior antioxidant and antiplatelet capacities for use in bakery food formulation and dietary recommendations.

**Citation:** Kosma, I.S.; Michalaki, A.; Kartelias, I.G.; Karantonis, H.C. Comparative Evaluation of Antioxidant Activities of Flours from Durum Wheat Varieties. *Biol. Life Sci. Forum* **2023**, *26*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Name

Published: date



**Copyright:** © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Keywords:** durum wheat flour; in vitro antioxidant activity; total phenolics; DPPH; FRAP; antiplatelet activity; platelet activating factor

## 1. Introduction

There is a growing focus on the connection between modern diets and lifestyles and various health issues, such as cardiovascular disease and malignancies in the digestive system. In recent years, there has been a notable rise in the consumption of nutritious food due to heightened consumer awareness and health-related considerations.

At the same time, it is projected that the global production of wheat would surpass 770 million tons in the year 2024. Wheat constitutes a significant proportion of Greece's agricultural output. According to a reliable source, it was projected that Greek wheat output will exceed the threshold of one million tons in the year 2024 [1]. One of the primary agricultural crop cultivated in Greece is durum wheat (*Triticum turgidum* L. var. durum). The use of durum wheat flour in the food industry provides both nutritional and economic advantages to Greece.

Bakery products from durum wheat are considered a staple meal due to its significant contribution to energy and nutrition, mostly attributed to its substantial carbohydrate and protein composition. Furthermore, wheat has essential nutrients and phytochemical compounds that are crucial for maintaining optimal health. Several phytochemicals included in wheat grains have been well acknowledged for their substantial biological impacts [2–4]. Wheat has the potential to possess secondary metabolites, such as polyphenols. Chemical compounds such as phenolic acids, coumarins, flavonoids, stilbenes, and lignans are included under the aforementioned group [3,4]. Previous studies have shown evidence of the antioxidant, anti-inflammatory, antimutagenic, and anticarcinogenic characteristics associated with polyphenols [4,5].

It has been shown that oxidative stress, inflammation and thrombosis is implicated in the development of several health conditions, including cancer, atherosclerosis, rheumatoid arthritis, and neurological disorders [6,7]. The capacity of antioxidants to eliminate free radicals and mitigate oxidative harm is well recognized [6]. Therefore, it is recommended to increase the consumption of foodstuffs that are high in antioxidants.

The aim of the study was to comparatively evaluate the antioxidant and antiplatelet potential of flours from 22 selected durum wheat varieties (*Triticum turgidum* L. var. durum) cultivated in Greece so that it is possible to be selected flours from those varieties that are with the greatest possible antioxidant and antiplatelet activities.

## 2. Materials and Methods

### 2.1. Materials

Twenty-two flour samples from different durum wheat (*Triticum turgidum* L. var. durum) varieties were kindly offered by the Institute of Applied Biosciences at the Centre for Research and Technology Hellas (INAB|CERTH).

### 2.2. Extract Preparation

Flour samples were extracted with aquas methanol 80% (*v/v*) according to Armelin et al. [8] with some modifications. Briefly, samples were agitated in an orbital shaker (GFL 3017, GFL; Burgwedel, Germany) at 200 rpm for 2 h at room temperature and then centrifuged at 1200× *g* force for 10 min (OrtoAlresa, Digicen 21R, Spain). The supernatant was stored at −40 °C until further analyzed for total phenolics, antioxidant activities and antiplatelet activity.

### 2.3. Total Phenolics

Total phenolics were measured in triplicate by using the Folin–Ciocalteu's method as previously described [9] using a spectrophotometer Lambda 25 (Perkin Elmer, Norwalk, CT, USA). Results were expressed as equivalent concentrations of gallic acid (mg GAE per g flour).

### 2.4. In Vitro Antioxidant Activities

The antioxidant activities were evaluated by the DPPH and FRAP assays as previously described [9]. Each sample was examined in triplicate. Trolox solutions were prepared in appropriate concentrations for quantitation purposes and the results expressed as Trolox equivalents in μmol per g of flour.

### 2.5. In Vitro Antiplatelet Activity

The in vitro antiplatelet activity was evaluated on an aggregometer (Chrono-log Corporation, 500CA, Havertown, PA, USA) by the ability of samples to inhibit lipid mediator of Platelet Activating Factor (PAF) towards human platelet rich plasma (PRP) as describe previously [10]. The in vitro antiplatelet activity was expressed as  $\text{mg}^{-1}$  required for 50% inhibition of PAF activity towards human PRP.

### 2.6. Statistics


Statistical analysis was performed with the SPSS software (IBM© SPSS® ver 28.0, IBM UK Ltd., Portsmouth, UK). Data were analyzed for normality using the Shapiro-Wilk test. Homogeneity of variance was checked by Levene's test. Data between the flours from different cultivars were compared statistically using one-way analysis of variance with post-hoc Bonferroni test for  $p < 0.05$ .

## 3. Results and Discussion

Data for total phenolics, antioxidant activities and antiplatelet activity were all normally distributed, as assessed by Shapiro-Wilk's test ( $p > 0.05$ ) and they passed the Levene's test for homogeneity of variance ( $p > 0.05$ ) and one way ANOVA with post-hoc Bonferroni test was applied to find any statistical differences in total phenolics, antioxidant activities and antiplatelet activity between the flour samples.

### 3.1. Total Phenolics

Total phenolic content for the flour samples range from as low as  $0.620 \pm 0.003$  mg gallic acid/g (Thraki) to as high as  $0.950 \pm 0.005$  mg gallic acid/g (Zeta E). This indicates significant variation in the phenolic content among the different samples. Svevo, with a total phenolic content of 0.89 mg gallic acid/g, has one of the highest phenolic contents among the samples. Lemnos, Canavaro, and Vendeta all have phenolic content value of 0.73 mg gallic acid/g, showing that these three samples have similar phenolic content. The results are presented in Figure 1. This variation of phenolic compounds among the different sample indicate possible differences in their antioxidant potential.




**Figure 1.** Total phenolics in 22 durum wheat (*Triticum turgidum* L. var. durum) flour samples. Different letters denote statistical difference based on one way ANOVA with with post-hoc Bonferroni test for  $p < 0.05$ .

### 3.2. In Vitro Antioxidant Activities

The DPPH values also vary widely among the samples, with values ranging from  $0.56 \pm 0.02$  to  $2.26 \pm 0.08$ . Higher Trolox equivalent (TE) values ( $\mu\text{mol}$  per g of flour) based


on DPPH assay indicate better scavenging of free radicals. Samples like “Marco Aurelio”, “Simeto”, “Svevo”, “Secolo”, “Dunavis” and “Zeta E” have notably higher values, suggesting stronger antioxidant activity in these samples (Figure 2). Similar to DPPH, TE ( $\mu\text{mol}$  per g of flour) based on FRAP assay vary across the samples, ranging from  $2.12 \pm 0.03$  to  $3.65 \pm 0.02$ . Again, “Marco Aurelio” exhibits the highest TE value, followed by samples like “Mexicali”, “Zoi”, “Pesti”, “Sifnos”, “Secolo”, and “Zeta E” indicating stronger antioxidant potential for those flours (Figure 2).



**Figure 2.** Antioxidant activities in 22 durum wheat (*Triticum turgidum* L. var. durum) flour samples based on DPPH and FRAP assays. The results are expressed as mean  $\pm$  SD of triplicate measurements and were expressed as trolox equivalents (TE) in  $\mu\text{mol}$  trolox per one gram of flour sample. Different letters denote statistical difference based on one way ANOVA with with post-hoc Bonferoni test for  $p < 0.05$ .

### 3.3. In Vitro Antiplatelet Activity

Higher values in ( $\text{mg}^{-1}$ ) suggest stronger antiplatelet activity. The sample with the highest antiplatelet activity is “Zoi” with a value of 1.38, followed closely by “Pigreco” and “Anna” with equal values of 1.31. On the other hand, the sample with the lowest antiplatelet activity is “Vendeta” with a value of 0.32.



**Figure 3.** Antiplatelet activity in 22 durum wheat (*Triticum turgidum* L. var. durum) flour samples based on in vitro inhibition of Platelet Activating Factor (PAF). The results are expressed as mean  $\pm$  SD of triplicate measurements and were expressed as the reverse of the amount of flour ( $\text{mg}^{-1}$ ) required for 50% inhibition of PAF activity ( $1/IA_{50}$ ) in human platelet rich plasma (hPRP). Different

letters denote statistical difference based on one way ANOVA with with post-hoc Bonferroni test for  $p < 0.05$ .

#### 4. Conclusions

The data provide valuable information about the antioxidant and antiplatelet potential of flour samples from different durum wheat (*Triticum turgidum* L. var. *durum*) cultivars. Community can use these data to make informed decisions about which samples may offer greater health benefits in terms of antioxidant and antiplatelet properties that are implicated in the development of various chronic diseases [6,7].

**Author Contributions:** Conceptualization, H.C.K.; methodology, I.S.K. and H.C.K.; software, H.C.K.; investigation, I.S.K., A.M., I.G.K. and H.C.K.; resources, H.C.K.; data curation, A.M. and H.C.K.; writing—original draft preparation, H.C.K.; writing—review and editing, H.C.K.; supervision, H.C.K.; project administration, H.C.K.; funding acquisition, H.C.K. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research has been co-financed by the European Union and Greek national funds through the Operational Program Competitiveness, Entrepreneurship and innovation, under the call RESEACH—CREAT—INNOVATE (project code: T2EDK-01244).

**Institutional Review Board Statement:**

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:**

**Acknowledgments:** This research has been co-financed by the European Union and Greek national funds through the Operational Program Competitiveness, Entrepreneurship and innovation, under the call RESEACH—CREAT—INNOVATE (project code: T2EDK-01244).

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

#### References

1. Food and Agriculture Organisation of the United Nations (FAOSTAT). Available Online: [Http://Www.Fao.Org/Faostat/En/#data](http://www.fao.org/faostat/en/#data) (Accessed on 13 September 2023).
2. Graziano, S.; Marando, S.; Prandi, B.; Boukid, F.; Marmiroli, N.; Francia, E.; Pecchioni, N.; Sforza, S.; Visioli, G.; Gulli, M. Technological Quality and Nutritional Value of Two Durum Wheat Varieties Depend on Both Genetic and Environmental Factors. *J. Agric. Food Chem.* **2019**, *67*, 2384–2395. <https://doi.org/10.1021/acs.jafc.8b06621>.
3. Graziano, S.; Marmiroli, N.; Visioli, G.; Gulli, M. Proteins and Metabolites as Indicators of Flours Quality and Nutritional Properties of Two Durum Wheat Varieties Grown in Different Italian Locations. *Foods* **2020**, *9*, 315. <https://doi.org/10.3390/foods9030315>.
4. Giacosa, A.; Peroni, G.; Rondanelli, M. Phytochemical Components and Human Health Effects of Old versus Modern Italian Wheat Varieties: The Case of Durum Wheat Senatore Cappelli. *Nutrients* **2022**, *14*, 2779. <https://doi.org/10.3390/nu14132779>.
5. Şahin, Y.; Yıldırım, A.; Yücesan, B.; Zencirci, N.; Erbayram, Ş.; Gürel, E. Phytochemical Content and Antioxidant Activity of Einkorn (*Triticum Monococcum* Ssp. *Monococcum*), Bread (*Triticum aestivum* L.), and Durum (*Triticum Durum* Desf.) Wheat. *Prog. Nutr.* **2017**, *19*, 450–459. <https://doi.org/10.23751/pn.v19i4.5847>.
6. Kiran, T.R.; Oflu, O.; Karabulut, A.B. Oxidative Stress and Antioxidants in Health and Disease. *J. Lab. Med.* **2023**, *47*, 1–11. <https://doi.org/10.1515/labmed-2022-0108>.
7. Demopoulos, C.A.; Karantonis, H.C.; Antonopoulou, S. Platelet Activating Factor—A Molecular Link between Atherosclerosis Theories. *Eur. J. Lipid Sci. Technol.* **2003**, *105*, 705–716. <https://doi.org/10.1002/ejlt.200300845>.
8. Armellini, R.; Peinado, I.; Pittia, P.; Scampicchio, M.; Heredia, A.; Andres, A. Effect of Saffron (*Crocus Sativus* L.) Enrichment on Antioxidant and Sensorial Properties of Wheat Flour Pasta. *Food Chem.* **2018**, *254*, 55–63. <https://doi.org/10.1016/j.foodchem.2018.01.174>.
9. Michalaki, A.; Karantonis, H.C.; Kritikou, A.S.; Thomaidis, N.S.; Dasenaki, M.E. Ultrasound-Assisted Extraction of Total Phenolic Compounds and Antioxidant Activity Evaluation from Oregano (*Origanum Vulgare* Ssp. *Hirtum*) Using Response Surface Methodology and Identification of Specific Phenolic Compounds with HPLC-PDA and Q-TOF-MS/MS. *Molecules* **2023**, *28*, 2033. <https://doi.org/10.3390/molecules28052033>.
10. Achilladelis, P.; Petsas, A.S.; Karantonis, H.C. Effect of Fortification of Tahini with Natural Plant Origin Raw Materials on Its Bioactivity. *Appl. Sci.* **2023**, *13*, 9626. <https://doi.org/10.3390/app13179626>.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.