

Enhancing Sorghum Productivity with Methyur, Kamethur, and Ivin Plant Growth Regulators[†]

Tsygankova V.A.^{1*}, Voloshchuk I.M.¹, Pilyo S.G.¹, Klyuchko S.V.¹ and Brovarets V.S.¹

¹Department for Chemistry of Bioactive Nitrogen-Containing Heterocyclic Compounds, V.P. Kukhar Institute of Bioorganic Chemistry and Petrochemistry, National Academy of Sciences of Ukraine, 1, Academician Kukhar str., 02094, Kyiv-94, Ukraine.

* Correspondence: **author:** Tsygankova Victoria Anatolyivna, Dr. Biol. Sci., Senior Staff Scientist, Principal researcher, email: V.Tsygankova@ukr.net, Tel: [+380681224673](tel:+380681224673).

[†]Presented at the 3rd International Electronic Conference on Agronomy French National Institute of Agricultural Research (IECAG 2023), France, 15 - 30 October 2023.

Abstract: Sorghum is one of the most important food, fodder and technical crops grown in the world. Global climate change and environmental pollution with toxic industrial and agricultural waste are the most unfavorable environmental factors affecting the growth and development of sorghum, which leads to a decrease in product quality. The development of new environmentally friendly plant growth regulators to improve growth and increase the productivity of sorghum is an urgent task of modern agriculture. Currently, considerable attention is paid to the development of new environmentally friendly plant growth regulators based on 6-methyl-2-mercapto-4-hydroxypyrimidine sodium and potassium salts (Methyur and Kamethur) and N-oxide-2,6-dimethylpyridine (Ivin). Thanks to the use of plant growth regulators Methyur, Kamethur and Ivin, it is possible to increase the productivity of agricultural crops and their adaptive properties to stress factors of abiotic nature. This work examines the use of plant growth regulators Methyur, Kamethur and Ivin to increase the productivity of sorghum. Field experiments were carried out on grain sorghum (*Sorghum bicolor* L.) cv. Yaron and sweet sorghum (*Sorghum saccharatum* L.) cv. Favorite. Seeds sterilized with 1% KMnO₄ solution were treated either with distilled water (control sample) or with solutions of any plant growth regulators Methyur, Kamethur or Ivin, applied at a concentration of 10⁻⁷ M for 24 hours (experimental sample). Each control and experimental sample contained 50 plant seeds, the experiments were carried out in triplicate. Then the soaked seeds were planted in the soil. Yield indicators such as panicle length (in cm) and fresh weight of grain (in grams), determined in experimental samples of sorghum plants, were calculated as % in relation to similar indicators determined in control samples of sorghum plants. It was shown that the yield indicators of sorghum plants grown for 4 month in the field, treated with Methyur, Kamethur and Ivin at a concentration of 10⁻⁷M exceeded those of control plants. Panicle length (in %) of experimental grain sorghum (*Sorghum bicolor* (L.) Moench) cv. Yaron increased: by 7% - in plants treated with Kamethur, by 20% - in plants treated with Methyur, by 17% - in plants treated with Ivin, compared to control. Panicle length (in %) of experimental sweet sorghum (*Sorghum saccharatum* (L.) Moench) cv. Favorite increased: by 36% - in plants treated with Kamethur, by 37% - in plants treated with Methyur, by 25% - in plants treated with Ivin, compared to control. Grain fresh weight (in %) of experimental grain sorghum (*Sorghum bicolor* (L.) Moench) cv. Yaron increased: by 22% - in plants treated with Kamethur, by 26% - in plants treated with Methyur, by 13% - in plants treated with Ivin, compared to control. Grain fresh weight (in %) of experimental sweet sorghum (*Sorghum saccharatum* (L.) Moench) cv. Favorite increased: by 24% - in plants treated with Kamethur, by 38% - in plants treated with Methyur, by 35% - in plants treated with Ivin, compared to control. Based on the results obtained, a conclusion was made about the high growth-stimulating effect of plant growth regulators, similar to the phytohormones auxins and cytokinin, and the dependence of their effect on their composition. It is proposed to use new en-

Citation: To be added by editorial staff during production.

Academic Editor: Firstname Lastname

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

environmentally friendly plant growth regulators Methyur, Kamethur and Ivin to improve growth and increase the productivity of sorghum, while reducing the use of environmentally toxic agrochemicals for plant protection and improving the environmental condition of the entire agricultural system.

Keywords: productivity of sorghum; plant growth regulators; Methyur; Kamethur; Ivin

1. Introduction

Sorghum (*Sorghum bicolor* (L.) Moench) is one of the important cereal food, fodder and technical crops grown in many countries around the world [1, 2]. Sorghum ranks fifth in the world after wheat, rice, corn and barley. Currently, sorghum is grown on almost all continents, over the past 50 years, the sorghum sown areas in the world amount to almost 44 million hectares [2].

The main advantage of sorghum is its high drought tolerance and unpretentiousness to soils, which makes this crop especially important in the context of global climate change [3]. However, there are problems with growing sorghum under adverse environmental conditions; for this purpose, plant growth regulators are used to improve growth and increase productivity of sorghum [4, 5].

This article describes the results of our previous work [6], in which the effect of new plant growth regulators based on pyrimidine and pyridine derivatives, such as Methyur (sodium salt of 6-methyl-2-mercapto-4-hydroxypyrimidine), Kamethur (potassium salt of 6-methyl-2-mercapto-4-hydroxypyrimidine) and Ivin (N-oxide-2,6-dimethylpyridine), synthesized in the Department for Chemistry of Bioactive Nitrogen-Containing Heterocyclic Compounds, V.P. Kukhar Institute of Bioorganic Chemistry and Petrochemistry of the National Academy of Sciences of Ukraine, on growth and productivity of grain sorghum (*Sorghum bicolor* L.) cv. Yarona, and sweet sorghum (*Sorghum saccharatum* L.) cv. Favorite was studied in the field.

2. Materials and methods

The chemical structure and relative molecular weight of new plant growth regulators Methyur and Kamethur (derivatives of sodium and potassium salts of 6-methyl-2-mercapto-4-hydroxypyrimidine) and Ivin (N-oxide-2,6-dimethylpyridine) are shown in Figure 1.




Figure 1. Chemical structure and relative molecular weight of plant growth regulators Methyur, Kamethur and Ivin.

Field experiments were carried out on grain sorghum (*Sorghum bicolor* L.) cv. Yarona, and sweet sorghum (*Sorghum saccharatum* L.) cv. Favorite. Seeds sterilized with 1% KMnO_4 solution were treated either with distilled water (control sample) or with solutions of any plant growth regulators Methyur, Kamethur or Ivin, applied at a concentration of 10^{-7} M for 24 hours (experimental sample). Each control and experimental sample contained 50 plant seeds. Then the soaked seeds were planted in the soil. The analysis of growth parameters: the average length of root (in mm) and the average fresh weight (in

gram) of sorghum grown for 2 month in the field, and productivity parameters: the average panicle length (in cm) and the average fresh weight of grain (in gram) of sorghum grown for 4 month in the field was carried out according to the guidelines [7]. Statistical processing of the data of the experiments performed in three replications was carried out according to the Student's-t variance test with a significance level of $P \leq 0.05$; the values are average \pm SD [8].

The growth parameters (the average length of root (in mm) and the average fresh weight (in gram)) of sorghum grown for 2 month in the field and productivity parameters (the average panicle length (in cm) and the average fresh weight of grain (in gram)) of sorghum grown for 4 month in the field, determined in experimental samples of sorghum plants, were calculated as % in relation to similar indicators determined in control samples of sorghum plants

3. Results

3.1. Study of the effect of plant growth regulators on sorghum growth

Field studies have shown that the growth indicators of experimental grain sorghum (*Sorghum bicolor* (L.) Moench) cv. Yarona exceeded that of control plants (Figure 2, A, B and C). Root length (in mm) increased: by 15% - in plants treated with Kamethur, by 14% - in plants treated with Methyur, by 29% - in plants treated with Ivin, compared to the control. Plant fresh weight (in gram) increased: by 67% - in plants treated with Kamethur, by 53% - in plants treated with Methyur, by 21% - in plants treated with Ivin, compared to control.




Figure 2. Growth indicators of 2-month-old grain sorghum (*Sorghum bicolor* (L.) Moench) cv. Yarona grown in the field: A - root length (in mm), B - plant fresh weight of (in gram), C - sorghum roots.

Growth indicators of experimental sweet sorghum (*Sorghum saccharatum* (L.) Moench) cv. Favorite exceeded that of control plants (Figure 3, A, B and C). Root length (in mm) increased: by 20% - in plants treated with Kamethur, by 40% - in plants treated

with Methyur, by 25% - in plants treated with Ivin, compared to the control. Plant fresh weight (in gram) increased: by 7% - in plants treated with Kamethur, by 57% - in plants treated with Methyur, by 30% - in plants treated with Ivin, compared to control.





Figure 3. Growth indicators of 2-month-old sweet sorghum (*Sorghum saccharatum* (L.) Moench) cv. Favorite grown in the field: A - root length (in mm), B - plant fresh weight of (in gram), C – sorghum roots.

3.2. Study of the effect of plant growth regulators on sorghum yield

Field studies have shown that the yield indicators of experimental grain sorghum (*Sorghum bicolor* (L.) Moench) cv. Yarona exceeded that of control plants (Figure 4, A, B and C). Panicle length (in cm) increased: by 7% - in plants treated with Kamethur, by 20% - in plants treated with Methyur, by 17% - in plants treated with Ivin, compared to control. Grain fresh weight (in gram) increased: by 22% - in plants treated with Kamethur, by 26% - in plants treated with Methyur, by 13% - in plants treated with Ivin, compared to control.



1
2
3
4

5

6

7
8
9

10

11
12
13
14
15
16
17
18

19




Figure 4. Yield indicators of 4-month-old grain sorghum (*Sorghum bicolor* (L.) Moench) cv. Yarona grown in the field: A – panicle length (in cm), B - grain fresh weight (in gram), C – panicles with sorghum grains.

Yield indicators of experimental sweet sorghum (*Sorghum saccharatum* (L.) Moench) cv. Favorite exceeded that of control plants (Figure 5, A, B and C). Panicle length (in cm) increased: by 36% - in plants treated with Kamethur, by 37% - in plants treated with Methyr, by 25% - in plants treated with Ivin, compared to control. Grain fresh weight (in gram) increased: by 24% - in plants treated with Kamethur, by 38% - in plants treated with Methyr, by 35% - in plants treated with Ivin, compared to control.




Figure 5. Yield indicators of 4-month-old sweet sorghum (*Sorghum saccharatum* (L.) Moench) cv. Favorite grown in the field: A – panicle length (in cm), B - grain fresh weight (in gram), C – panicles with sorghum grains.

Summarizing the data obtained, it should be concluded that synthetic plant growth regulators Methyur, Kamethur and Ivin have a high stimulating effect on the growth and development of sorghum shoots and roots in the vegetative phase, as well as on the growth and development of sorghum panicles and grain formation in the reproductive phase. Obviously, their high growth-stimulating effect may be explained by their auxin-like and cytokinin-like effects on the processes of proliferation, elongation and differentiation of plant cells [9 - 11]. In addition, the composition of synthetic plant growth regulators has a positive effect on plant growth and development. A plant growth regulator Ivin contains the macronutrient nitrogen; Kamethur contains the macronutrients nitrogen, potassium and sulfur, which are necessary for plant growth and metabolism, and plant adaptation to biotic and abiotic stress factors [12, 13]. Plant growth regulator Methyur, containing the macronutrients nitrogen, sulfur and the chemical element sodium, promotes plant growth and productivity, as well as plant adaptation to salt and osmotic stress [14 - 16].

4. Conclusions

The results of field studies confirmed the possibility of practical application of the new environmentally friendly plant growth regulators based on 6-methyl-2-mercapto-4-hydroxypyrimidine sodium and potassium salts (Methyur and Kamethur) and N-oxide-2,6-dimethylpyridine (Ivin) to improve the growth and increase the productivity of sorghum. Thanks to the use of these environmentally friendly plant growth regulators in a nanomolar, environmentally non-toxic concentration of 10^{-7} M to treat seeds before planting in the soil, it will be possible to improve the growth and increase the productivity of sorghum and its adaptation to stress factors, while reducing the use of environmentally toxic agrochemicals for plant protection and improving the environmental condition of the entire agricultural system.

Author Contributions: Conceptualization, T.V.A. and B.V.S.; methodology, T.V.A.; validation, T.V.A., V.I.M., P.S.G., K.S.V., and B.V.S.; investigation, T.V.A., V.I.M., P.S.G., K.S.V., and B.V.S.; writing—original draft preparation, T.V.A. and V.I.M.; writing—review and editing, T.V.A.; supervision, T.V.A. and B.V.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was carried out with the support of the project “Elaboration of new plant growth regulators on the base of azaheterocycles derivatives” (Resolution of the Bureau of the Academy of Sciences of the National Academy of Sciences of Ukraine from June 8, 2021 No. 4. State registration No 0122U000442, code: 2.1.10.12-22. Decision of the Expert Council on Evaluation of Topics of Fundamental Research Works at the National Academy of Sciences of Ukraine from 07.07.2021 No. 2).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Aruna C., Visarada K.B.R.S., Bhat B.V., Tonapi V.A. (Eds.) Breeding Sorghum for Diverse End Uses. Woodhead Publishing Series in Food Science, Technology and Nutrition. 1st ed., Elsevier Ltd. UK, 2018; 450 p.
2. Mundia C.W., Secchi S., Akamani K. and Wang G. A. Regional Comparison of Factors Affecting Global Sorghum Production: The Case of North America, Asia and Africa’s Sahel. Sustainability, 2019; 11: 2135. doi:10.3390/su11072135.
3. Kidanemariam W. Review on Mechanisms of Drought Tolerance in Sorghum (*Sorghum bicolor* (L.) Moench) Basis and Breeding Methods. Acad. Res. J. Agri. Sci. Res. 2019; 7(2): 87-99. DOI: 10.14662/ARJASR2019.007.
4. Macedo W.R., Araujo D.K., Santos V.M., Castro P.R.C., Fernandes G.M. Plant growth regulators on sweet sorghum: Physiological and nutritional value analysis. Comunicata Scientiae, 2017; 8(1): 170-175. DOI: 10.14295/CS.v8i1.1315.

5. Li H., Wang X.L., Guo X.Q., Rao M., Steinberger Y., Cheng X., Xie G.H. Effects of plant growth regulators on growth, yield and lodging of sweet sorghum. *Res. on Crops*, 2011: 12(2): 372-382. 1
6. Tsygankova V.A., Voloshchuk I.V., Klyuchko S.V., Pilyo S.G., Brovarets V.S., Kovalenko O.A. The effect of pyrimidine and pyridine derivatives on the growth and productivity of sorghum. *International Journal of Botany Studies*, 2022: 7(5): 19-31. 2
7. Voytsehovska, O.V., Kapustyan, A.V., Kosik, O.I., Musienko, M.M., Olkhovich, O.P., Panyuta, O.O., Parshikova, T.V., Glorious, P.S. 2010. *Plant physiology: praktykum*. Parshikova T.V. (Ed). Lutsk: Teren, 420 p. 3
8. Bang H., Zhou X.K., van Epps H.L., Mazumdar M. *Statistical Methods in Molecular Biology*. Series: Methods in molecular biology, New York: Humana press, 2010, 13(620), 636 p. URL: <https://doi.org/10.1007/978-1-60761-580-4> 4
9. Lavy M., Estelle M. Mechanisms of auxin signaling. *Development*. 2016. 143(18): 3226 - 3229. DOI: 10.1242/dev.131870 5
10. Kieber J.J., Schaller G.E. Cytokinin signaling in plant development. *Development*. 2018. 145(4): dev149344: 1 – 7. doi: 10.1242/dev.149344. 6
11. Tsygankova V. A., Andrushevich Ya.V., Shtompel O. I., Solomyanny R. M., Hurenko A. O., Frasinuk, Mrug G.P., Shablykin O.V., Pilyo S. G., Kornienko A. M. & Brovarets V. S. New Auxin and Cytokinin Related Compounds Based on Synthetic Low Molecular Weight Heterocycles. Chapter 16. P. 353-377. In: Aftab T. (Ed.) *Auxins, Cytokinins and Gibberellins Signaling in Plants*. Signaling and Communication in Plants. Springer Nature Switzerland AG. 2022. 377 p. URL: https://doi.org/10.1007/978-3-031-05427-3_16 7
12. Tsygankova, V.A., Blyuss, K.B., Shysha, E.N., Biliavska, L.A., Iutynska, G.A., Andrushevich, Ya.V., Ponomarenko, S.P., Yemets, A.I., Blume, Ya.B. Using microbial biostimulants to deliver RNA interference in plants as an effective tool for biocontrol of pathogenic fungi, parasitic nematodes and insects. Chapter 6. In: *Research advances in plant biotechnology*. Series: Plant Science Research and Practices. Blume Ya.B. (Ed). USA: Nova Science Publishers, Inc., 2020. pp. 205 - 319. 8
13. Pidlisnyuk V., Mamirova A., Newton R.A., Stefanovska T., Zhukov O., Tsygankova V., and Shapoval P. The role of plant growth regulators in *Miscanthus × giganteus* utilisation on soils contaminated with trace elements. *Agronomy*. 2022. 12(12): 2999. DOI: <https://doi.org/10.3390/agronomy12122999>. 9
14. Bilyavska N.O., Voloshina N.Yu., Topchii N.M., Konturska O.O., Palladina T.O. Effects of salt and osmotic stresses and Methyure on foliar photosynthetic apparatus in maize, *Bulletin of Kharkiv National Agrarian University, Series: Biology*. 2009. 3: 35 - 42. URL: <https://repo.btu.kharkov.ua/handle/123456789/8030> 10
15. Palladina T.O., Ribchenko Zh.I., Konturska O.O. Dependence of preparation Metiure adaptogenic effect on plants under salt stress conditions from its molecular structure. *Biotechnologia Acta*. 2012. 5(1): 115 – 120. URL: <https://cyberleninka.ru/article/n/dependence-of-preparation-metiure-adaptogenic-effect-on-plants-under-salt-stress-conditions-from-its-molecular-structure> 11
16. Rudnytska M.V., Palladina T.A. Effect of preparations Methyur and Ivine on Ca²⁺-ATPases activity in plasma and vacuolar membrane of corn seedling roots under salt stress conditions, *Ukr. Biochem. J.* 2017. 89(1): 76 – 81. URL: <https://doi.org/10.15407/ubj89.01.076> 12