

Proceeding Paper

# Thiohydrazides in the Synthesis of Functionalized Extranuclear Heterosteroids †

Yulia Volkova <sup>1,\*</sup>, Alexander Scherbakov <sup>2,3</sup> and Igor Zavarzin <sup>1</sup>

<sup>1</sup> N.D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 47 Leninsky prosp., Moscow 119991, Russia; email1@email.com

<sup>2</sup> N.N. Blokhin National Medical Research Center of Oncology, Kashirskoye shosse 24, Moscow 115522, Russia; email2@email.com

<sup>3</sup> National Research Lobachevsky State University of Nizhny Novgorod, Gagarina 23, Nizhny Novgorod 603950, Russia

\* Correspondence: yavolkova@gmail.com

† Presented at the 27th International Electronic Conference on Synthetic Organic Chemistry (ECSOC-27), 15–30 November 2023; Available online: <https://ecsoc-27.sciforum.net/>.

**Abstract:** Heterocyclic derivatives of hormones have attracted great interest as a privileged scaffold for drug discovery due to their outstanding biological activity. A number of them are potent anti-cancer agents, which are used in the chemotherapy of breast and prostate cancers. Here, the data obtained by the authors in the field of studying functionalized thiohydrazides as simple “versatile agents” for the installation of heterocyclic moiety to the steroid core are summarized. Namely, a flexible synthetic approach to unknown pyrazolines, 1,3,4-thiadiazole, thiadiazine, and pyridazine derivatives of steroids with selective control of heterocyclization patterns are discussed. Steroidal 1,3,4-thiadiazoles were obtained by oxidative heterocyclization of oxamic acid thiohydrazides with 16-hydroxymethylidene- $\Delta^{1,3,5(10)}$ -estratrieno-17-one. An extension of this reaction to steroidal  $\alpha,\beta$ -unsaturated ketones resulted in androst-5-ene-[17,16d]-pyrazolines. Spiro-androstene-17,6'[1',3',4']thiadiazines were exclusively synthesized employing 16 $\beta$ ,17 $\beta$ -epoxyprogrenolone. Using 21-bromopregna-5,16-dien-20-one as a substrate, 17-[1',3',4']thiadiazine-substituted androstenes were prepared. 18-Nor-5 $\alpha$ -androsta-2,13-diene[3,2-d]pyridazines, androsta-2-ene[3,2-d]pyridazines and  $\Delta^{1,3,5(10)}$ -estratrieno[16,17-d]pyridazines were synthesized via two steps involving the Vilsmeier-Haack reaction of enolizable steroidal ketones giving chlorovinyl aldehydes followed by imination of the former with oxamic acid thiohydrazides. The antiproliferative activity of the synthesized compounds against breast and prostate cancer cell lines, along with lead compounds' in-depth characterization are included. The lead compounds were found to have potent selectivity and in some cases a significant effect on the signaling pathways in parental and 4-hydroxytamoxifen-resistant cells.

**Keywords:** steroids; heterocycles; thiohydrazides; anticancer activity

**Citation:** Volkova, Y.; Scherbakov, A.; Zavarzin, I. Thiohydrazides in the Synthesis of Functionalized Extranuclear Heterosteroids. *Chem. Proc.* **2023**, *14*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Name

Published: 15 November 2023



**Copyright:** © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction


The heterocyclic derivatives of steroids have gained significant attention in pharmaceutical research as a promising framework for drug discovery [1,2]. For example, abiraterone, 17-pyridine-bearing androsta-5,16-dien-3-ol, is currently used in clinic for the treatment of advanced prostate cancers [3]. Synthetic androgen danazol, also known as isoxazolethisterone, is marked for the treatment of endometriosis and premenstrual syndrome [4]. Dutasteride, bearing 5,6-dihydro-2H-pyridin-2-one motif, is a potent 5 $\alpha$ -reductase inhibitor used as a part of benign prostatic hyperplasia therapy [5]. The continually growing demand for the development of new therapeutic agents with improved selectivity and reduced side effects for the treatment of a variety of diseases

requires the synthesis of heterocyclic derivatives of steroids with high levels of molecular diversity.

A number of excellent reviews on synthetic strategies directed towards the synthesis of heterosteroids have emerged in recent years [6,7]. Some of them focused on heterocycles [8], including fused-thiazoles [9], pyrazolines [10], bicyclic pyridines [11] and pyrimidines [12]. Other highlight the advances of steroid D-ring modification [13,14] or advances in the synthesis of steroidal conjugates and dimers [15–17]. The use of a click chemistry [18] and transition-metal species [19] in the synthesis of heterosteroids is also considered. Here we summarize our achievements in the synthesis of heterocyclic derivatives of steroids based on oxamic acid thiohydrazides as polyheteroatom reagents. These reagents are readily available from  $\alpha$ -chloroacetamides [20] and enable various five- and six-membered *N,O*- and *S*-heterocycles construction [21–26]. The use of oxamic acid thiohydrazides as a reagent for the modification of steroids has never been systematically analyzed. The antiproliferative activity of the synthesized compounds against breast and prostate cancer cell lines, as well as a detailed characterization of the lead compounds, are included.

## 2. Results and Discussion


Our interest in the functionalized heterocyclic derivatives of steroids [27–33] resulted in based on oxamic acid thiohydrazides flexible strategy for the installation of heterocyclic pendant at steroid core initially containing a carbonyl group (Figure 1). Intermediate hydrazones derived from oxamic acid thiohydrazides and steroids due to tautomerization and *cis/trans* isomerization can undergo oxidative aromatization, electrocyclic reactions, SH- and NH-nucleophilic cyclizations discussed below.



**Figure 1.** Reactivity of hydrazones derived from oxamic acid thiohydrazides and steroids.

### 2.1. Steroidal 1,3,4-thiadiazoles

A frequently observed reaction of thiohydrazone-derived hydrazones when subjected to oxidative conditions is the occurrence of an intramolecular cyclization. This process involves the attack of a *S*-nucleophile on the electrophilic *C=N* bond, resulting in the formation of 1,3,4-thiadiazoles [24]. Having observed a seemingly general transformation pattern for hydrazones of oxamic acid thiohydrazides derived from steroids, we elaborated an efficient method for the synthesis of steroidal carbamoyl-1,3,4-thiadiazoles [34]. Reactions of thiohydrazides **2** with 16-hydroxymethylidene-5 $\alpha$ -androstan-17-one and 16-hydroxymethylidene- $\Delta^{1,3,5(10)}$ -estratrieno-17-one **1** furnished the corresponding 1,3,4-thiadiazoles derivatives **5**.




**Scheme 1.** Heterocyclization toward 1,3,4-thiadiazoles.

The reaction proceeds via an AdN-type cyclization of the thione isomer intermediate **3** to form cyclic thiadiazoline **4**, which subsequently undergoes aromatization (oxidative, if X = H) leading to the formation of thiadiazoles **5**. The efficiency of air as an oxidant in the reaction involving glacial acetic acid has been observed.

## 2.2. Steroidal Pyrazolines

This reaction can be further extended to steroidal  $\alpha,\beta$ -unsaturated ketones **6**, resulting in the selective formation of androst-5-ene-[17,16d]-pyrazolines **10** (Scheme 2) [35–37]. The reaction conditions employed are comparable to those documented in literature for the preparation of thiadiazoles [24] except for heating in pure glacial acetic acid. This slight alteration in circumstances promotes the development of pyrazolines **10**.




**Scheme 2.** Synthesis of androst-5-ene-[17,16d]-pyrazolines.

The proposed mechanism suggests that the initial step of the reaction involves the formation of hydrazones, as mentioned earlier. However, when these hydrazones are in their protonated form (referred to as form **7**), they undergo NH-nucleophilic addition to an activated double bond leading to the synthesis of pyrazolines **8**. After the 1,3[H] shift and deprotonation of intermediates **9**, pyrazolines **10** are formed. The heterocyclization process was observed to be more favorable for oxamic acid thiohydrazides that have electron-donating substituents at the aryl group (yields ranged from 75–82%).

## 2.3. Steroidal Pyridazines


Besides, it has been shown that the reaction of the oxamic acid thiohydrazides **2** with steroidal  $\beta$ -chlorovinyl aldehydes **11** under acidic conditions results in annulated steroidal pyridazines **14** (Scheme 3) [38]. The developed method was marked by significant efficiency in synthesizing derivatives of the androstene and estrane series, which incorporate pyridazine motifs fused to the A and D rings of the steroid core. The reaction is believed to occur via cascade imination/ $6\pi$ -electrocyclization of hydrazone thiol tautomers **13** to furnish the pyridazines **14** [39].



**Scheme 3.** Cyclization towards steroidal pyridazines.


#### 2.4. Steroidal 1,3,4-thiadiazines

Reactions of oxamic acid thiohydrazides **2** with steroids bearing  $\alpha$ -bromoketone moiety resulted in steroidal 1,3,4-thiadiazines with high chemoselectivity [40]. Namely reaction of 21-bromopregna-5,16-dien-20-one **15** with oxamic acid thiohydrazides **2** under mild basic conditions affords 17-(6'H-1',3',4'-thiadiazine-2'-carboxamide)androst-5,17-dienes **17** in virtually quantitative yields (Scheme 4).



**Scheme 4.** Synthesis of 1',3',4'-thiadiazine androst-5,17-dienes.


The spiro-androstene-17,6'[1',3',4']thiadiazines **20** were exclusively synthesized by reaction of 16 $\beta$ ,17 $\beta$ -epoxypregnenolone **18** with oxamic acid thiohydrazides **2** in the presence of sulfuric acid (Scheme 5) [40]. The relative spatial arrangement of the electrophilic reaction site and the carbonyl moiety in the steroid structure defines the reaction product. In both heterocyclizations key step is the nucleophilic attack of hydrazone SH-group at electrophilic reaction sites at the steroid core (see intermediates **16** and **19**).



**Scheme 5.** Synthesis of 16 $\beta$ -hydroxyspiro-androsteno-17,6'[1,3,4]thiadiazines.

#### 2.5. Antiproliferative Activity

The above-described heterocyclic derivatives of steroids showed significant antiproliferative activity. Overall lead compounds **I** and **II**-pyridazines annulated with 17 $\beta$ -hydroxy-5 $\alpha$ -androst-2-ene at A-ring and 3 $\beta$ -acetoxyandrost-5-ene at D-ring, exhibited selectivity against hormone-dependent human breast cancer cell line along with higher cytotoxicity than the cisplatin-reference drug (Figure 2) [38].



**Figure 2.** Cytotoxic effects of steroidal pyridazines and 1,3,4-thiadiazines.

17-[1',3',4']Thiadiazine-substituted androstenes with general formula **III** (Figure 2) were lead compounds against human androgen receptor-positive prostate cancer cells 22Rv1 [40]. They showed micromolar values of IC<sub>50</sub> with antiproliferative potency higher than those for bicalutamide-reference drug. Compounds **III**-induced 22Rv1 cell death was shown to be associated with modulation of the AR, ERK 1/2, NF-κB, and PARP pathways. Thus, compounds **I-III** are of great interest for more in-depth study and may be considered as a candidates for future anticancer drug design, in particular, against hormone-receptor-positive breast and prostate cancers.

### 3. Conclusions

The development of synthetic methods based on simple intermediates leading to a variety of structurally diverse heterocyclic derivatives of natural products has been a focus for medicinal and organic chemists for years. The use of oxamic acid thiohydrazides as precursors for the formation of hydrazones from steroids has led to a range of heterocyclic derivatives of steroids in a highly chemoselective manner. Carbamoyl-substituted steroidal thiadiazoles, thiadiazines, pyrazolines, and pyridazines were obtained from readily available starting materials under mild conditions by tuning the hydrazones reactivity profiles. Data on the biological activity of these compounds could enable the further exploration of the therapeutic potential of heterocyclic derivatives of steroids. First of all, it is relevant for developing novel and highly effective anticancer drugs for the treatment of breast and prostate cancers.

**Funding:** This work was financially supported by the Russian Science Foundation (project No. 22-13-00161, <https://rscf.ru/project/22-13-00161/> accessed on 6 September 2023).

**Conflicts of Interest:** The authors declare no conflict of interest.

### References

1. Yu, B.; Cai, Z.; Wang, S.; Liu, H. Recent advances on the synthesis and antitumor evaluation of exonuclear heterosteroids. *Chin. J. Org. Chem.* **2017**, *37*, 1952.
2. Singh, H.; Jindal, D.P.; Yadav, M.R.; Kumar, M. 5 Heterosteroids and Drug Research. In *Progress in Medicinal Chemistry*; Ellis, G.P., West, G.B., Eds.; Elsevier: Amsterdam, The Netherlands, 1991; Volume 28, pp. 233–300.
3. Abiraterone acetate. *Drugs R&D* **2010**, *10*, 261–269.
4. Hughes, E.; Brown, J.; Tiffin, G.; Vandekerckhove, P. Danazol for unexplained subfertility. *Cochrane Database Syst. Rev.* **2007**, Cd000069.
5. Arif, T.; Dorjay, K.; Adil, M.; Sami, M. Dutasteride in Androgenetic Alopecia: An Update. *Curr. Clin. Pharmacol.* **2017**, *12*, 31–35.
6. Farhan, A.; Alshamusi, Q.; Jebur, M. Synthesis of steroid bearing heterocyclic derivatives and biological activity. Review 2014–2020. In *Journal of Physics: Conference Series*; IOP Publishing: Bristol, UK, 2021; p. 012057.
7. Ibrahim-Ouali, M.; Dumur, F. Recent syntheses of steroidal derivatives containing heterocycles. *ARKIVOC-Online J. Org. Chem.* **2020**, *2019*, 304–339.
8. Sharma, K.; Kumar, H. Formation of nitrogen-containing six-membered heterocycles on steroidal ring system: A review. *Steroids* **2022**, *191*, 109171.
9. Pawar, S.; Kumar, K.; Gupta, M.K.; Rawal, R.K. Synthetic and medicinal perspective of fused-thiazoles as anticancer agents. *Anti-Cancer Agents Med. Chem. (Former. Curr. Med. Chem.-Anti-Cancer Agents)* **2021**, *21*, 1379–1402.
10. Matiadis, D.; Sagnou, M. Pyrazoline hybrids as promising anticancer agents: An up-to-date overview. *Int. J. Mol. Sci.* **2020**, *21*, 5507.
11. Hammouda, M.M.; Elattar, K.M.; Rashed, M.M.; Osman, A.M.A. Synthesis and biological activities of bicyclic pyridines integrated steroid hybrid. *Steroids* **2023**, *199*, 109287.
12. Elattar, K.M.; El-Mekabaty, A. Heterocyclic steroids: Synthetic routes and biological characteristics of steroidal fused bicyclic pyrimidines. *J. Heterocycl. Chem.* **2021**, *58*, 389–414.
13. Iqbal, A.; Siddiqui, T. A review on synthesis and biological activities of D-ring modified pregnenolone. *Steroids* **2021**, *170*, 108827.
14. Huo, H.; Li, G.; Shi, B.; Li, J. Recent advances on synthesis and biological activities of C-17 aza-heterocycle derived steroids. *Bioorganic Med. Chem.* **2022**, *69*, 116882.
15. Raičević, V.; Radulović, N.; Sakač, M. Toward Selective Anticancer Agents: Ferrocene-Steroid Conjugates. *Eur. J. Inorg. Chem.* **2022**, *2022*, e202100951.

16. Nahar, L.; Sarker, S.D. A review on steroid dimers: 2011–2019. *Steroids* **2020**, *164*, 108736.
17. Paquin, A.; Reyes-Moreno, C.; Bérubé, G. Recent Advances in the Use of the Dimerization Strategy as a Means to Increase the Biological Potential of Natural or Synthetic Molecules. *Molecules* **2021**, *26*, 2340.
18. Ibrahim-Ouali, M.; Hamze, K. A click chemistry approach to secosteroidal macrocycles. *Steroids* **2014**, *80*, 102–110.
19. Le Bideau, F.; Dagonne, S. Synthesis of transition-metal steroid derivatives. *Chem. Rev.* **2013**, *113*, 7793–7850.
20. Yarovenko, V.; Shirokov, A.; Krupinova, O.; Zavarzin, I.; Krayushkin, M. Synthesis of oxamic acids thiohydrazides and carbamoyl-1, 3, 4-thiadiazoles. *Russ. J. Org. Chem.* **2003**, *39*, 1133–1139.
21. Komendantova, A.S.; Lyssenko, K.A.; Zavarzin, I.V.; Volkova, Y.A. Iodine-promoted synthesis of pyrazoles from 1, 3-dicarbonyl compounds and oxamic acid thiohydrazides. *Org. Chem. Front.* **2020**, *7*, 1640–1646.
22. Komkov, A.V.; Komendantova, A.S.; Menchikov, L.G.; Chernoburova, E.I.; Volkova, Y.A.; Zavarzin, I.V. A straightforward approach toward multifunctionalized pyridazines via imination/electrocyclization. *Org. Lett.* **2015**, *17*, 3734–3737.
23. Komendantova, A.S.; Ivanova, K.A.; Lyssenko, K.V.; Volkova, Y.A.; Zavarzin, I.V. Facile Synthesis of Carboxamide-Substituted 1, 3, 4-Thiadiazines and 5, 6-Dihydro-4 H-1, 3, 4-Thiadiazin-5-Ols. *Chem. Heterocycl. Compd.* **2019**, *55*, 665–671.
24. Krayushkin, M.; Yarovenko, V.; Zavarzin, I. Synthesis and reactivity of monothiooxamides and thiohydrazides of oxamic acids. *Russ. Chem. Bull.* **2004**, *53*, 517–527.
25. Volkova, Y.A.; Chernoburova, E.I.; Petrova, A.S.; Shtil, A.A.; Zavarzin, I.V. Reactions of hydrazones derived from oxamic acid thiohydrazides. *Phosphorus Sulfur Silicon Relat. Elem.* **2017**, *192*, 237–240.
26. Krayushkin, M.; Yarovenko, V.; Zavarzin, I. Synthesis of heterocyclic compounds based on oxamic acid monothiooxamides and thiohydrazides. *Russ. Chem. Bull.* **2019**, *68*, 1143–1163.
27. Vorontsova, S.K.; Yadykov, A.V.; Scherbakov, A.M.; Minyaev, M.E.; Zavarzin, I.V.; Mikhaevich, E.I.; Volkova, Y.A.; Shirinian, V.Z. Novel d-Annulated Pentacyclic Steroids: Regioselective Synthesis and Biological Evaluation in Breast Cancer Cells. *Molecules* **2020**, *25*, 3499.
28. Samanta, S.; Ghosh, A.K.; Ghosh, S.; Ilina, A.A.; Volkova, Y.A.; Zavarzin, I.V.; Scherbakov, A.M.; Salnikova, D.I.; Dzichenka, Y.U.; Sachenko, A.B. Fe (iii)-Catalyzed synthesis of steroidal imidazoheterocycles as potent antiproliferative agents. *Org. Biomol. Chem.* **2020**, *18*, 5571–5576.
29. Volkova, Y.A.; Kozlov, A.S.; Kolokolova, M.K.; Uvarov, D.Y.; Gorbatov, S.A.; Andreeva, O.E.; Scherbakov, A.M.; Zavarzin, I.V. Steroidal N-Sulfonylimidates: Synthesis and biological evaluation in breast cancer cells. *Eur. J. Med. Chem.* **2019**, *179*, 694–706.
30. Scherbakov, A.M.; Zavarzin, I.V.; Vorontsova, S.K.; Hajra, A.; Andreeva, O.E.; Yadykov, A.V.; Levina, I.S.; Volkova, Y.A.; Shirinian, V.Z. Synthesis and evaluation of the antiproliferative activity of benzylidenes of 16-dehydropregesterone series. *Steroids* **2018**, *138*, 91–101.
31. Scherbakov, A.M.; Komkov, A.V.; Komendantova, A.S.; Yastrebova, M.A.; Andreeva, O.E.; Shirinian, V.Z.; Hajra, A.; Zavarzin, I.V.; Volkova, Y.A. Steroidal pyrimidines and dihydrotriazines as novel classes of anticancer agents against hormone-dependent breast cancer cells. *Front. Pharmacol.* **2018**, *8*, 979.
32. Rassokhina, I.V.; Volkova, Y.A.; Kozlov, A.S.; Scherbakov, A.M.; Andreeva, O.E.; Shirinian, V.Z.; Zavarzin, I.V. Synthesis and antiproliferative activity evaluation of steroidal imidazo [1, 2-a] pyridines. *Steroids* **2016**, *113*, 29–37.
33. Birukova, V.; Scherbakov, A.; Ilina, A.; Salnikova, D.; Andreeva, O.; Dzichenka, Y.; Zavarzin, I.; Volkova, Y. Discovery of highly potent proapoptotic antiestrogens in a series of androst-5, 16-dienes D-modified with imidazole-annulated pendants. *J. Steroid Biochem. Mol. Biol.* **2023**, *231*, 106309.
34. Zavarzin, I.; Antonov, Y.S.; Chernoburova, E.; Shchetinina, M.; Kolotyrykina, N.; Shashkov, A. Interaction of 16-hydroxymethylidene derivatives of androstane and estrone with thiohydrazides of oxamic acids. *Russ. Chem. Bull.* **2013**, *62*, 2603–2608.
35. Zavarzin, I.V.; Kamernitsky, A.; Chertkova, V.V.; Chernoburova, E.I.; Yarovenko, V.N.; Kraushkin, M.M.; Kachala, V.V. Synthesis of 1'-arylcarbamoylthiocarbonyl-3'-methyl-3-oxoandrost-4-eno [16 $\alpha$ , 17 $\alpha$ -d] pyrazolines. *Arkhivoc* **2008**, *4*, 62–70.
36. Kamernitsky, A.; Chernoburova, E.; Chertkova, V.; Zavarzin, I.; Yarovenko, V.; Krayushkin, M. Acylhydrazones of 20-keto steroids and their transformations: I. Synthesis and properties of 1'-acyl-substituted 3'-methylandrosteno [16, 17-d] pyrazolines. *Russ. J. Bioorganic Chem.* **2007**, *33*, 315–319.
37. Kamernitskii, A.; Chernoburova, E.; Chertkova, V.; Yarovenko, V.; Zavarzin, I.; Krayushkin, M. Effect of  $\omega$ -substituents in the hydrazones of conjugated pregnane 20-ketosteroids on their ability to cyclize to pyrazolines. *Russ. Chem. Bull.* **2006**, *55*, 2117–2118.
38. Volkova, Y.A.; Antonov, Y.S.; Komkov, A.V.; Scherbakov, A.M.; Shashkov, A.S.; Menchikov, L.G.; Chernoburova, E.I.; Zavarzin, I.V. Access to steroidal pyridazines via modified thiohydrazides. *RSC Adv.* **2016**, *6*, 42863–42868.
39. Komendantova, A.S.; Fakhrutdinov, A.N.; Menchikov, L.G.; Sukhorukov, A.Y.; Zavarzin, I.V.; Volkova, Y.A. Cyclization of  $\beta$ -Chlorovinyl Thiohydrazones into Pyridazines: A Mechanistic Study. *Eur. J. Org. Chem.* **2019**, *2019*, 527–536.
40. Komendantova, A.S.; Scherbakov, A.M.; Komkov, A.V.; Chertkova, V.V.; Gudovanniy, A.O.; Chernoburova, E.I.; Sorokin, D.V.; Dzichenka, Y.U.; Shirinian, V.Z.; Volkova, Y.A. Novel steroidal 1, 3, 4-thiadiazines: Synthesis and biological evaluation in androgen receptor-positive prostate cancer 22Rv1 cells. *Bioorganic Chem.* **2019**, *91*, 103142.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.