

Enhancing Vegetative Growth of Quinoa and Soil Properties Under Water Shortage through Targeted Organic Amendments

Muhammad Zubair Akram ^{1,2,*}, Angela Libutti ³ and Anna Rita Rivelli ²

¹ Ph.D. Program in Agricultural, Forest and Food Sciences, University of Basilicata, Via dell'Ateneo Lucano, 10, 85100 Potenza, Italy

² School of Agricultural, Forest, Food and Environmental Sciences, University of Basilicata, Via dell'Ateneo Lucano, 10, 85100 Potenza, Italy

³ Department of Science of Agriculture, Food, Natural Resources and Engineering, University of Foggia, Via Napoli, 25, 71122 Foggia, Italy

* Correspondence: muhammadzubair.akram@unibas.it

Abstract: The scarcity of water resources is considered a major threat and challenge for agriculture. The water limitations could strongly affect the growth and development of many crops including quinoa, a nutrition-rich, climate-resilient crop that is gaining attention globally. The organic amendment application is reported as a suitable option to mitigate the detrimental impacts of water shortage on soil and plant growth. In this context, two experiments were conducted on *Chenopodium quinoa* "Titicaca"; in the first one, we investigated the effect of different organic amendments i.e., woodchips biochar (Bw), vineyard pruning biochar (Bv) and vermicompost (V), applied (alone and mixed) at 2% soil dry weight, on soil properties and plant biomass of quinoa subjected to a water stress period during vegetative development. Among organic amendments tested, Bw and Bw+V increased plant biomass on average by 15%, while Bv and Bv+V reduced the plant biomass by 62% compared to non-amended soil (C). A significant reduction in soil pH was observed by Bw (7.61) while BV increased pH (8.04) compared to C (7.76). The Bw and Bv also reduced soil bulk density (BD) (1.19 g/m³ and 1.13 g/m³, respectively) compared to C (1.28 g/m³). As Bw performed better in the first experiment, the second one assess only Bw at different doses, i.e., 0%, 2% and 4% under water shortage by restoring only 50% evapotranspiration losses, when soil water content reached the 50% of available water content. Considering the Bw rates, the plants treated with Bw2% showed 34% and 19% more biomass, while 36% and 66% more panicles than Bw0% and Bw4%, respectively. The Bw2% decreases the soil pH (7.79 versus 7.85) and electrical conductivity (286 versus 307 μS/cm) compared to Bw0% that was not different from Bw4%. No differences were observed in BD between Bw0% and Bw2% (on average 1.28 g/m³), while decreased in Bw4% (1.06 g/m³). The findings of both experiments highlighted that the appropriate type and dose of biochar could improve soil properties and help the quinoa plants to grow better under water-limited conditions.

Keywords: *Chenopodium quinoa*; Titicaca genotype; water shortage; organic amendments; biochar rates; vegetative development; soil properties

Citation: Akram, M.Z.; Libutti, A.; Rivelli, A.R. Enhancing Vegetative Growth of Quinoa and Soil Properties Under Water Shortage through Targeted Organic Amendments.

2023, volume number, x.

<https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Last-name

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Quinoa (*Chenopodium quinoa* Willd.) is a resilient seed crop belonging to the Amaranthaceae family; native to the Andean region is nowadays cultivated in many parts of the world. The exceptional nutritional qualities and tolerance against various abiotic stresses are well-documented [1,2]. However, wide differences are reported among the germplasm collections of the species currently available [3]. The Danish genotype Titicaca exhibited varying morphological and physiological responses, revealed its sensitivity to drought [4]. As already known, drought significantly affects soil's physico-chemical properties, including soil fertility, soil aeration, decomposition of organic matter and microbial

activity [5]. Therefore, it is necessary to devise alternative methods for soil management, environmentally friendly and sustainable such as organic soil amendments, i.e. compost and biochar. Compost contains a large amount of labile organic matter, which stimulates soil biota, promotes plant growth by enhancing microbial activity and many soil properties [6], mitigating negative consequences of water limitations [5]; the biochar addition improves soil's structure, porosity and water-holding capacity, plant water status and photosynthetic efficiency [7]; in quinoa it enhances the vegetative development when facing early periods of severe water stress [8]. However, biochar's net benefits depend upon the feedstocks, pyrolysis conditions, dose, soil type and crop [9]. The present study aims to investigate the effects of different organic amendments on selected soil properties and growth of quinoa plants (Titicaca genotype), followed by testing the best performing amendment at different application doses, when the species is subjected to water limitations.

2. Material and Methods

2.1. Description of Treatments and Experimental Designs

Two experiments were carried out at the University of Basilicata, Potenza (PZ, 40°38'N–15°48' E, 819 m a.s.l.) in Southern Italy, during the spring-summer 2022 on quinoa (*Chenopodium quinoa* Willd), Titicaca genotype. The sandy loam texture soil (USDA classification) was used in the experiments. The biochar from woodchips (Bw) was purchased from Nera Biochar Company (Settimo Vittone, Italy), the biochar from vineyard prunings (Bv) was produced at the STAR*Facility Centre of Foggia University (South Italy), and the vermicompost from cattle manure (V) was purchased from a company located in Montescaglioso, Matera district (South Italy).

2.1.1. Experiment 1

Pots were filled with 5 kg of soil or soil treated with different organic amendments, i.e., the two types of biochar, Bw and Bv, and vermicompost (V), alone or in mixed form (Bw+V and Bv+V) at a rate of 2% of soil dry weight (dw). Unamended soil was considered as a control (C). Then, soil was brought to field capacity (FC), previously determined together with the permanent wilting point (PWP) at -0.03 and -1.5 MPa, respectively, for each treatment. The soil available water content (AWC, % dw) was calculated as the difference between FC and PWP. Following, 10 seeds were sown in each pot and thinned to one plant per pot once the emergence was completed. The seven experimental treatments were replicated three times and the 21 experimental units were arranged according to a completely randomized design. From the emergence until the twelve-leaf stage, the plants were kept well-irrigated by restoring 100% of evapotranspiration losses at the depletion of 40% of the AWC, then a period of water stress was applied until the soil reached the PWP. At the end of this period, the pots were re-watered to FC and kept well-watered until the end of experiment, i.e., initiation of flowering when plants were cut and parameters were recorded.

2.1.2. Experiment 2

The pots were filled with 6 kg of soil treated with three different rates (w/w) of woodchip biochar: 0% (Bw0% as control), 2% (Bw2%) and 4% (Bw4%) of dry soil weight (dw). Each experimental treatment was replicated three times and the 9 experimental units were arranged according to a completely randomized design. From seedling emergence to the beginning of flowering, plants were irrigated by restoring 50% of evapotranspiration losses at the depletion of 50% of the AWC.

2.2. Soil and Plant Growth-Related Parameters.

At the end of both experiments, soil-related parameters i.e., pH, electrical conductivity (EC) and bulk density (BD) were measured. Additionally, the total number of panicles

(TNP, n°) and total plant fresh weight (TFW, g) were recorded. All the experimental data were processed by one-way analysis of variance (ANOVA) by using “Statistix 8.1” software. When significant differences among means were detected, the latter were compared by Tukey’s honest significance difference post hoc test at the 5% probability level.

3. Results

3.1. Soil- and plant growth-related parameters of experiment 1.

The changes in soil-related parameters due to the application of different organic amendments at 2% rate, are shown in Table 1. The pH variations ranged between 7.6 and 8.1. The woodchip biochar treatment (Bw) resulted in a significant decrease ($p < 0.001$) of soil pH (0.2 unit), while vineyard pruning biochar alone (Bv) and mixed with vermicompost (Bv+V) increased ($p < 0.001$) soil pH (0.3 unit) compared to the control (C). The electrical conductivity (EC) was not found significantly affected by treatments. Bulk density (BD) significantly ($p < 0.001$) declined in all treatments compared to C, with greater reduction in Bv+V (18%). The biochar produced from woody material has a larger surface area and lower ash content as compared to the biochar produced from crop residues or grasses [10]. In our case, the higher ash content in Bv (9.9 %) is the driver for the higher pH than Bw (4.4 %), which could reduced the availability of micronutrients, according to previous study [11]. Moreover, the higher porosity of Bv [12] than Bw, reduced bulk density and likely increased nutrient retention [12].

The significant effect of the organic amendments tested on plant growth under period of water stress is presented in Table 1. The Bw and Bw+V treated plants produced more ($p < 0.001$) panicles (on average 17) contrary to Bv and Bv+V (on average 9.5). Likewise, the Bw and Bw+V treatments yielded higher ($p < 0.001$) biomass (on average 35.9 g), while the Bv and Bv+V yielded the lower one (on average 19.3 g). Using biochar boosted the nutritional availability and source-sink relationship under drought, increasing panicle length and grain yield also in rapeseed [13], and in quinoa as reported by Kammann et al. [14] under similar biochar application. Probably, the higher porosity of Bv [12] than Bw, although reducing the bulk density, increased nutrient retention, especially N [15], lowering their availability for plant growth.

Table 1. pH, Electrical Conductivity (EC, $\mu\text{S}/\text{cm}$), Bulk Density (BD, g/m^3) of soil along with Total Number of Panicles (TNP), and Total Fresh Weight (TFW, g) of the quinoa plants grown under different organic amendments.

Experimental Factors	Soil related parameters			Plant Growth	
	pH	EC	BD	TNP	TFW
C	7.8 B	268	1.28 A	13 B	31.4 B
Bw	7.6 C	297	1.19 B	18 A	34.6 A
Bv	8.0 A	279	1.13 BC	7 C	18.6 C
V	7.8 B	291	1.20 AB	15 AB	31.1 B
Bw+V	7.7 BC	274	1.14 BC	16 AB	37.1 A
Bv+V	8.1 A	302	1.05 C	12 BC	20.1 C
Significance	***	ns	***	***	***

Values are means ($n = 3$). In each column, means followed by different letters are significantly different ($p \leq 0.05$; Tukey’s test). ***, F test significant at $p \leq 0.001$, respectively. C, control; Bw, woodchip biochar; Bv, vineyard pruning biochar; V, vermicompost.

3.2. Soil- and plant growth-related parameters of experiment 2.

Applying different woodchip biochar rates significantly affected the soil properties under water shortage conditions (Table 2). More specifically, the addition of 2% biochar (Bw2%) decreased ($p < 0.001$) pH (7.79) than 4% rate (Bw4%) (7.82) as compared to the non amended soil (Bw0%) (7.9). The EC was also reduced ($p < 0.05$) in Bw2% treated soil (286 $\mu\text{S}/\text{cm}$), while increased in Bw4% (313 $\mu\text{S}/\text{cm}$) compared to Bw0% (307 $\mu\text{S}/\text{cm}$). Both

Bw4% and Bw2% treatments reduced bulk density (BD) by 20% and 8%, respectively, compared to Bw0%. Several studies reported changes (increase or decrease) in pH level by application different biochar rates [16]. Consistent with our study, a decrease in soil pH after biochar addition was observed by Qiang et al. [17]. Soil pH variation after biochar application, influence the microbial activity and enzyme production, carbon and nutrient availability [18]. The higher biochar application rate may increase soil salinity and EC, subsequently leading to undesirable impacts on the plant growth [19]. Moreover, biochar addition, by lowering soil bulk density increases the total soil porosity, thereby positively impacting root development [20].

Significant variations of plant growth were observed among the different biochar rates under water shortage conditions (Table 2). In particular, the Bw2% treated plants showed more ($p < 0.05$) panicles (15) compared to Bw0% and Bw4% (on average 10). Similarly, the Bw2% plants produced higher ($p < 0.05$) biomass (19.5 g) than Bw0% and Bw4% (on average 15.6 g). The significant enhancement of biomass and yield was reported in quinoa under water limited conditions by Kammann et al. [14]. However, some non-significant or negative impacts of biochar have also been reported [21,22], leading to consider appropriate biochar type and dose for desired responses. In the present study, Bw4% negatively affects plant growth, as reported by several studies due to detrimental effects including reduced soil nutrient availability under altered pH [15,23]. The higher biochar rate increased negative charges on soil surface, leading to intensified electrostatic interactions with cations [24], as also reported by Rees et al. [15] under woody biochar application, resulting in detrimental effect on plant growth and yield due to reduced nutrient availability [25].

Table 2. pH, Electrical Conductivity (EC, $\mu\text{S}/\text{cm}$), Bulk Density (BD, g/m^3) of soil along with Total Number of Panicles (TNP) and Total Fresh Weight (TFW, g) of the quinoa plants grown under different rates of woodchips biochar.

Experimental Factors	Soil related parameters			Plant Growth	
	pH	EC	BD	TNP	TFW
Bw0%	7.85 A	307 A	1.32 A	11 B	14.6 B
Bw2%	7.79 B	286 B	1.23 A	15 A	19.7 A
Bw4%	7.82 A	313 A	1.06 B	9 B	16.5 B
Significance	***	*	**	*	*

Values are means ($n = 3$). In each column, means followed by different letters are significantly different ($p \leq 0.05$; Tukey's test). *, **, ***, F test significant at $p \leq 0.05$, $p \leq 0.01$ and $p \leq 0.001$, respectively. Bw0%, no biochar; Bw2%, 2% biochar, Bw4%, 4% biochar.

4. Conclusions

Among the organic amendments tested, woodchip biochar application, alone or mixed with vermicompost, was shown to be the better option to mitigate short-term water stress conditions arising during the vegetative growing cycle of quinoa, Titicaca genotype. Considering the different woodchip biochar rates, 2% application determined greater growth response of Titicaca under water limitations compared to 4% biochar, probably due to more favorable soil conditions. Both experiments pointed out that the proper type and dose of biochar could help the quinoa plants to grow better and cope with water-limited conditions. Further investigations should be extended to the evaluation of different quinoa genotypes to better understand the specific response to biochar application, in controlled and open field conditions.

Author Contributions: Conceptualization A.L. and A.R.R.; methodology, M.Z.A. A.L. and A.R.R.; software, M.Z.A.; validation, A.L. and A.R.R.; formal analysis, M.Z.A.; investigation, M.Z.A., A.L. and A.R.R.; resources, A.R.R.; data curation, M.Z.A., A.L. and A.R.R.; writing—original draft preparation, M.Z.A., A.L. and A.R.R.; writing—review and editing, M.Z.A., A.R.R. and A.L.;

visualization, M.Z.A.; supervision, A.L. and A.R.R.; project administration, A.R.R.; funding acquisition, A.L. and A.R.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available upon request from the corresponding authors.

Acknowledgments: We would like to thank Simone Orlandini and Leonardo Verdi for the quinoa seeds for the current experiments. The authors are grateful to Giuseppe Mercurio for his technical assistance in the greenhouse experiment and the laboratory analyses.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Akram, Z.; Maqsood Ahmed Basra, S.; Bilal Hafeez, M.; Khan, S.; Nazeer, S.; Iqbal, S.; Sohail Saddiq, M.; Zahra, N. Adaptability and yield potential of new quinoa lines under agro-ecological conditions of Faisalabad-Pakistan. *Asian Journal of Agriculture and Biology* **2021**, *2*, 2005301.
2. Yadav, R.; Gore, P.G.; Gupta, V.; Siddique, K.H. Quinoa (*Chenopodium quinoa* Willd.)—a smart crop for food and nutritional security. In *Neglected and Underutilized Crops*; Elsevier: 2023; pp. 23–43.
3. Hafeez, M.B.; Iqbal, S.; Li, Y.; Saddiq, M.S.; Basra, S.M.; Zhang, H.; Zahra, N.; Akram, M.Z.; Bertero, D.; Curti, R.N. Assessment of phenotypic diversity in the USDA collection of quinoa links genotypic adaptation to germplasm origin. *Plants* **2022**, *11*, 738.
4. Issa Ali, O.; Fghire, R.; Anaya, F.; Benlhabib, O.; Wahbi, S. Physiological and Morphological Responses of two Quinoa Cultivars (*Chenopodium quinoa* Willd.) to Drought Stress. *Gesunde Pflanzen* **2019**, *71*.
5. Quintana, J.R.; Martín-Sanz, J.P.; Valverde-Asenjo, I.; Molina, J.A. Drought differently destabilizes soil structure in a chronosequence of abandoned agricultural lands. *Catena* **2023**, *222*, 106871.
6. Chen, Y.; Lv, X.; Qin, Y.; Zhang, D.; Zhang, C.; Song, Z.; Liu, D.; Jiang, L.; Huang, B.; Wang, J. Effects of different botanical oil meal mixed with cow manure organic fertilizers on soil microbial community and function and tobacco yield and quality. *Frontiers in Microbiology* **2023**, *14*, 1191059.
7. Abideen, Z.; Koyro, H.-W.; Huchzermeyer, B.; Bilquees, G.; Khan, M.A. Impact of a biochar or a biochar-compost mixture on water relation, nutrient uptake and photosynthesis of *Phragmites karka*. *Pedosphere* **2020**, *30*, 466–477.
8. Akram, M.Z.; Libutti, A.; Rivelli, A.R. Evaluation of Vegetative Development of Quinoa under Water Stress by Applying Different Organic Amendments. *Agronomy* **2023**, *13*, 1412.
9. Manikandan, S.; Vickram, S.; Subbaiya, R.; Karmegam, N.; Chang, S.W.; Ravindran, B.; Awasthi, M.K. Comprehensive review on recent production trends and applications of biochar for greener environment. *Bioresource Technology* **2023**, 129725.
10. Li, Q.; Zhang, X.; Mao, M.; Wang, X.; Shang, J. Carbon content determines the aggregation of biochar colloids from various feedstocks. *Science of The Total Environment* **2023**, *880*, 163313.
11. Kishimoto, S. Charcoal as a soil conditioner. In *Proceedings of the Symposium on Forest Product Research, International Achievements for the Future, 1985, 1985*; pp. 12–23.
12. Azuara, M.; Sáiz, E.; Manso, J.A.; García-Ramos, F.J.; Manyà, J.J. Study on the effects of using a carbon dioxide atmosphere on the properties of vine shoots-derived biochar. *Journal of Analytical and Applied Pyrolysis* **2017**, *124*, 719–725.
13. Khan, Z.; Khan, M.N.; Zhang, K.; Luo, T.; Zhu, K.; Hu, L. The application of biochar alleviated the adverse effects of drought on the growth, physiology, yield and quality of rapeseed through regulation of soil status and nutrients availability. *Industrial Crops and Products* **2021**, *171*, 113878.

14. Kammann, C.I.; Schmidt, H.-P.; Messerschmidt, N.; Linsel, S.; Steffens, D.; Müller, C.; Koyro, H.-W.; Conte, P.; Joseph, S. Plant growth improvement mediated by nitrate capture in co-composted biochar. *Scientific reports* **2015**, *5*, 11080.
15. Rees, F.; Germain, C.; Sterckeman, T.; Morel, J.-L. Plant growth and metal uptake by a non-hyperaccumulating species (*Lolium perenne*) and a Cd-Zn hyperaccumulator (*Noccaea caerulescens*) in contaminated soils amended with biochar. *Plant and Soil* **2015**, *395*, 57-73.
16. Palansooriya, K.N.; Ok, Y.S.; Awad, Y.M.; Lee, S.S.; Sung, J.-K.; Koutsospyros, A.; Moon, D.H. Impacts of biochar application on upland agriculture: A review. *Journal of environmental management* **2019**, *234*, 52–64.
17. Qiang, M.; Gao, J.e.; Han, J.; Zhang, H.; Lin, T.; Long, S. How adding biochar improves loessal soil fertility and sunflower yield on consolidation project land on the Chinese loess plateau. *Polish Journal of Environmental Studies* **2020**, *29*, 3759-3769.
18. Stark, S.; Männistö, M.K.; Eskelinen, A. Nutrient availability and pH jointly constrain microbial extracellular enzyme activities in nutrient-poor tundra soils. *Plant and Soil* **2014**, *383*, 373–385.
19. Tag, A.T.; Duman, G.; Ucar, S.; Yanik, J. Effects of feedstock type and pyrolysis temperature on potential applications of biochar. *Journal of analytical and applied pyrolysis* **2016**, *120*, 200–206.
20. Chang, Y.; Rossi, L.; Zotarelli, L.; Gao, B.; Shahid, M.A.; Sarkhosh, A. Biochar improves soil physical characteristics and strengthens root architecture in Muscadine grape (*Vitis rotundifolia* L.). *Chemical and Biological Technologies in Agriculture* **2021**, *8*, 1–11.
21. Haider, G.; Koyro, H.-W.; Azam, F.; Steffens, D.; Müller, C.; Kammann, C. Biochar but not humic acid product amendment affected maize yields via improving plant-soil moisture relations. *Plant and soil* **2015**, *395*, 141–157.
22. Ramlow, M.; Foster, E.; Del Grosso, S.; Cotrufo, M. Broadcast woody biochar provides limited benefits to deficit irrigation maize in Colorado. *Agriculture, Ecosystems & Environment* **2019**, *269*, 71–81.
23. Jeffery, S.; Verheijen, F.G.; van der Velde, M.; Bastos, A.C. A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agriculture, ecosystems & environment* **2011**, *144*, 175–187.
24. Rees, F.; Simonnot, M.-O.; Morel, J.-L. Short-term effects of biochar on soil heavy metal mobility are controlled by intra-particle diffusion and soil pH increase. *European Journal of Soil Science* **2014**, *65*, 149–161.
25. Wang, Y.; Zhang, L.; Yang, H.; Yan, G.; Xu, Z.; Chen, C.; Zhang, D. Biochar nutrient availability rather than its water holding capacity governs the growth of both C3 and C4 plants. *Journal of Soils and Sediments* **2016**, *16*, 801–810.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.