

Proceeding Paper

A Study on the People of North-East Part of Bangladesh with the Effect of Flood Disaster that Occurred in 2022

Asraf Ibna Helal*

Department of Statistics, Shahjalal University of Science & Technology, Sylhet, Bangladesh

* Correspondence: asraf16@student.sust.edu;

† Presented at the 4th International Electronic Conference on Applied Sciences, 27 October – 10 November 2023; Available online: <https://asec2023.sciforum.net>.

Abstract: Bangladesh is characterized by its tropical climate and low elevation. Every year, this region is affected by flooding. The proximity of India adds another significant layer of complexity to this issue. Because India opens its switch gate during the rainy season, Bangladesh sees an increase in the amount of water that flows through the country. The area often has issues with flooding. In the year 2022, the residents of the north-east part of Bangladesh were put in a terrible position due to flooding. Therefore, one of our goals is to find factors associated with this flood disaster. In addition to this, one of our goals is to mitigate the negative impact that the flood had on the physical and emotional health of the people living in the north-east part of Bangladesh. This study also includes some demographic and socio-economic factors associated with this environmental disaster.

Keywords: environment; flood; disaster; physical health; mental health

1. Introduction

In South Asia, floods occur frequently, particularly in Bangladesh, where these cause a variety of losses [1]. Bangladesh is ranked sixth in the world as the nation most susceptible to flooding due to its topography and the negative effects of climate change [2]. Bangladesh has already experienced bad floods during the floods of 1988, 1998, 2004, and 2007 and is gradually becoming more susceptible to floods, not only due to changes in the environment but also due to the high-speed increase in residents in the coastal plains and the prevalent poverty that forces individuals to live in floodplains [3]. The tropical climate and low elevation of Bangladesh are its defining features [4]. The proximity of India contributes to the region's annual floods [5]. The north-east part of Bangladesh, which includes Sylhet, Sunamganj, and Netrokona Districts, is the main sufferer of flood disasters [6]. The amount of water flowing through Bangladesh increases during the rainy season as a result of India opening its switch gate [7]. Floods in 2022 put the people of Sylhet in a difficult situation. Poverty is a major factor in people's susceptibility to flooding, and frequent flood impacts contribute to an increase in poverty and, consequently, susceptibility [8]. In Bangladesh, floods in 1974, 1988, 1998, and 2017 killed approximately 4,000 people and destroyed approximately 18 million kilograms of crops, according to previous research [9]. This overall condition highlights the vulnerability of Bangladesh's human sector to flooding. In densely populated regions with compact urban structures, flooding can also have devastating effects [10]. Bangladesh, a country with a low per capita income, suffers severe economic damage from floods. Human vulnerability refers to the physical, economic, social, and natural circumstances that make those who are exposed to hazards more vulnerable [11]. The process of identifying human vulnerability of flooding is not a novel topic of discussion due to the close relationship between location and people. Bangladesh is susceptible to four kinds of destructive flooding: river floods, precipitation floods, flash floods, and cyclone floods [12]. Heavy monsoon precipitation and thawing

Citation: To be added by editorial staff during production.

Academic Editor: Firstname Last-name

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

snow in the upper catchment areas of Bangladesh's main rivers cause river inundation [13]. Heavy precipitation in the eastern and northern hill streams is the primary cause of flash flooding [14]. In April and May, the most frequent climatic shocks in the region, a flash flood devastates the region and causes the collapse of flood protection infrastructures such as earthen dams and embankments, as well as inundating and destroying the developing paddy [15]. Bari et al., the quantity of rain that falls in the Sylhet region during the months of March and April (i.e., during the time of early flash flooding) is approximately three times the national average for the same time period. Geomorphologically, the region is situated between the northern Indian state of Meghalaya and the eastern Indian states of Assam and Tripura [16]. The terrain of the Haor region is significantly lower than that of the adjacent Indian region [17]. Thus, the haor belt not only obtains its own rainfall but also the runoff water from the aforementioned Indian catchments upstream [18]. Cherrapunji, a region of Indian Meghalaya, is one of the highest rainfall regions in the globe and is situated directly above the Haor region of Bangladesh that is only 30 kilometers from the nearest Bangladeshi frontier, as seen from the air [19]. The temporal and spatial variability as well as the total rainfall that occurs in the northeastern part of Bangladesh and its upstream Indian areas are crucial for analyzing the flood characteristics in the haor regions, but this issue has received no attention whatsoever [20]. The northeastern region of Bangladesh is prone to excessive rainfall. This region in Bangladesh is separate from others due to its hilly terrain and propensity for inundation [21]. Agriculture, fisheries, and the local economy were adversely affected by the floods caused by sudden rainstorms [22]. The objectives of this study are to lessen the harm that the floods do to the mental and physical health of the residents of this area and to understand the current economic environment and the parameters of the damage rate. The goals of this study are to determine the impact of the recent flood tragedy on the residents of the north-east and the economic loss experienced by locals.

2. Materials and Methods

Information was collected from various districts in the north-east of Bangladesh that were affected by the 2022 disaster. Approximately 416 data were collected for this study. We used a Google Form and a manual document to collect the data.

Then the data were cleaned and processed for analysis using SPSS version 25. I performed descriptive statistics to see the actual scenario of flood-affected people. Exploratory data analysis and a cross-sectional study were performed to find factors that are associated with flood disasters. The chi-square test from a cross-sectional study was used to fulfill our aims. This study also executed binary logistic regression to determine the factors that affected the flood disaster.

3. Result and Discussion

Table 1. Descriptive Analysis of variables related to the flood effected peoples.

Variables	Category	Frequency	Percentage
Age of respondents	Less than 42	339	81.5
	Greater than or equal to 42	77	18.5
Gender	Male	320	76.9
	Female	96	23.1
Marital Status	Married	238	57.2
	Unmarried	174	41.8
	Others	4	1.0
Area of Resident	Rural	166	39.9
	Urban	250	60.1

Monthly Family Income	0-10000	84	20.2
	10000-15000	104	25
	15000-20000	68	16.3
	20000-30000	58	13.9
	30000-50000	70	16.8
	50000+	32	7.7
Flood Affected/not	Yes	358	86.1
	No	58	13.9
Damaged by flood	Yes	324	77.9
	No	92	22.1
Moved to safe place/shelter	Yes	202	48.6
	No	214	51.4
Suffered from food	Yes	244	58.7
	No	172	41.3
Received Relief Assistance	Yes	144	34.6
	No	272	65.4
Got any disease	Yes	126	30.3
	No	290	69.7
Type of disease	No disease	290	69.7
	Watery Diarrhea	20	4.8
	Others (fever, cough, cold etc.)	106	25.5
Mentally pressured	Yes	384	92.3
	No	32	7.7
Having children in family	Yes	204	49
	No	212	51
Got children disease	Yes	82	19.7
	No	334	80.3
Type of child disease	No disease	334	80.3
	Watery Diarrhea	20	4.8
	Others (fever, cough, cold etc.)	62	14.9
Suffered from pure drinking water	Yes	260	62.5
	No	156	37.5
Got water purification tablet	Yes	104	25
	No	312	75

Table 1 shows the descriptive study of flood-affected people in the north-east part of Bangladesh. It gives an overview of the area affected by the flood in 2022. It shows that 81.5% of respondent's ages are less than 42, and 18.5% of respondents ages are greater than or equal to 42. The gender-wise percentages of male and female respondents are 76.9 and 23.1, respectively. The study results that 39.9% of respondents live in rural areas, and the rest live in urban areas. The percentage of people who live in urban areas is 60.1, which means a huge portion of people live in urban societies. Additionally, it has been seen that floods affect 86.1% of people and cause 77.9% of property damage. Approximately half of the respondents could move to safe places during the flood. The percentage of people who suffered from hunger and received relief assistance is 58.7 and 34.6, respectively. There are various diseases that affect about 30.3% of people, and the proportions of people who have watery diarrhea and those who don't are 4.8% and 25.5%, respectively. This flood disaster put a lot of respondents under mental strain. The percentage of people who suffered from pure drinking water was 62.5. 25% of people got the water purification tablet.

Table 2. Chi-square test result.

Factors		Flood affected/not		Chi-square value	P-value
		Yes	No		
Damaged by flood	Yes	320	38	197.180 ^a	0.000
	No	4	54		
Moved to safe places/shelter	Yes	194	164	32.609 ^a	0.000
	No	8	50		
Suffered from food	Yes	234	124	47.662 ^a	0.000
	No	10	48		
Received relief assistance	Yes	138	220	17.541 ^a	0.000
	No	6	52		
Got any disease	Yes	114	244	2.941 ^a	0.086
	No	12	46		
Mental pressure	Yes	350	8	107.713 ^a	0.000
	No	34	24		
Got children disease	Yes	80	278	11.264 ^a	0.001
	No	2	56		
Suffer from pure drinking water	Yes	260	98	112.328 ^a	0.000
	No	0	58		
Got water purification tablet	Yes	96	264	4.514 ^a	0.034
	No	8	50		

Table 2 shows the cross-sectional study result, which is the chi-square test result. This interprets the associated factors with the flood disaster. The chi-square test revealed that eight variables have p-values that are less than 0.05. The variables are: damaged by flood, moved to safe places or shelters, suffered from food, received relief assistance, experienced mental pressure, had children with diseases, suffered from pure drinking water, and got a water purification tablet. These variables have a significant association with flood disasters. This means that these variables have a significant effect on flooding. But unfortunately, any disease has no association with flooding because the p-value (0.084) of the variable is greater than 0.05. Further, I have performed a binary logistic regression model with these associated variables. The result is given in the below tables.

Table 3. Omnibus Tests result of Binary Logistic Regression Model.

Omnibus Tests of Model Coefficient				
		Chi-square	df	Sig.
Step 1	Step	236.444	8	0.000
	Block	236.444	8	0.000
	Model	236.444	8	0.000

Table 4. Hosmer and Lemeshow Test result of Binary Logistic Regression Model.

Hosmer and Lemeshow Test			
Step	Chi-square	df	Sig.
1	6.402	8	.602

Tables 3 and 4 show the results of the binary logistic regression model. Table 3 has demonstrated the omnibus test results. Omnibus tests of the model coefficient are used to test the model fit. If the model is significant, this shows that there is a significant improvement in fit as compared to the null model. Hence, the model is showing good fit.

Table 4 has revealed the Hosmer and Lemeshow test results, a test of model fit. The Hosmer and Lemeshow statistics indicate a poor fit if the significance value is less than 0.05. The model adequately fits the data. There is no difference between the observed and

predicted models. Finally, the model interprets that these associated variables have an impact on flood-affected people.

Then the study has gone through one way ANOVA to find area wise variation of flood affected peoples. The result is given below,

Table 5. ANOVA table for one way ANOVA.

ANOVA					
	Sum of squares	df	Mean square	F	Sig
Between groups	1.732	1	1.732	14.881	0.000
Within groups	48.182	414	.116		
Total	49.913	415			

Table 6. Descriptive analysis table for one way ANOVA.

	N	Mean	Std. Deviation	Std. Error	95% CI for Mean	
					Lower Bound	Upper Bound
Rural	166	.06	.239	.019	.02	.10
Urban	250	.19	.395	.025	.14	.24
Total	416	.14	.347	.017	.11	.17

Tables 5 and 6 show the results of a one-way ANOVA between flood-affected and non-flood-affected areas of residence, which is used to find the area-based variation in living of flood-affected people. ANOVA (Table 5) shows that the test is statistically significant (P-value 0.05). Table 6 reveals there is a significant mean difference between rural and urban flood disasters. Where the mean of rural areas' flood disasters (0.06) is greater than the mean of urban areas' flood disasters (0.19). This means that urban areas are most affected by flooding.

4. Conclusion

In the conclusion, the findings show precise descriptive studies about the peoples of the north-east part of Bangladesh who were affected by the flood in 2022. It has given an overview of the flood disaster and flood-affected people. Most of the flood-affected people lived in urban areas in those districts of Bangladesh. The study includes eight significant variables associated with the flood disaster. Those who were damaged by floods moved to safe places or shelters, suffered from food, received relief assistance, suffered from mental pressure, got children's diseases, suffered from pure drinking water, and got water purification tablets. The outcome of this research has shown these variables have a significant impact on flood-affected people and the urban area is more affected by flooding than the rural area.

References

1. T. Hasnat, M.A. Kabir, M.A. Hossain, Biodiversity Survey of Madhupur National Park sub-project under Biodiversity Conservation and Eco-tourism Development in Bangladesh project View project Recolonization of Threatened Tree Species View project, (n.d.).
2. S. Agrawala, T. Ota, A.U. Ahmed, J. Smith, M. Van Aalst, ENVIRONMENT DIRECTORATE DEVELOPMENT CO-OPERATION DIRECTORATE Working Party on Global and Structural Policies Working Party on Development Co-operation and Environment DEVELOPMENT AND CLIMATE CHANGE IN BANGLADESH: FOCUS ON COASTAL FLOODING AND THE SUNDARBANS, (n.d.).
3. M. Alam, G. Rabbani, Vulnerabilities and responses to climate change for Dhaka, IIED), 19 (2007) 81–97.
4. G. Mcgranahan, D. Balk, B. Anderson, The rising tide: assessing the risks of climate change and human settlements in low elevation coastal zones, IIED), 19 (2007) 17–37.
5. Gerd Marmulla Fishery Resources Officer Inland Water Resources and Aquaculture Service Fishery Resources Division FAO Fisheries Department Dams, fish and fisheries Opportunities, challenges and conflict resolution, (2001).
6. S. Hossain, Md., Flood damage and risk assessment model in the haor basin of Bangladesh, (2013).

7. M. Mainuddin, F. Karim, D.S. Gaydon, J.M. Kirby, Impact of climate change and management strategies on water and salt balance of the polders and islands in the Ganges delta, *Scientific Reports* 1, 11 (123AD) 7041.
8. D. Philip, M.I. Rayhan, Vulnerability and Poverty: What are the causes and how are they related?, (2004).
9. S.H. Jahangir Masum, Climatic Hazards in Bangladesh-A Literature Review Safeguarding Coastal Ecosystem from Irresponsible Commercial Shrimp farming (SECO) View project Strengthening People's Action on Climate Risk Reduction & Energy Efficiency (SPACE) View project, (2019).
10. K./ Arnold, / Carlin, M. Arnold, A. Carlin, T.W. Bank, Building Safer Cities The Future of Disaster Risk, (2003).
11. N. Brooks, Vulnerability, risk and adaptation: A conceptual framework, (2003).
12. Floods in a Changing Climate: Risk Management - Slobodan P Simonović - Google Books, (n.d.).
13. T.A. Waseem Ghazi Anm Muniruzzaman Singh, Climate Change & Security in South Asia Cooperating for Peace, (2016).
14. G. El Afandi, M. Morsy, F. El Hussieny, Heavy Rainfall Simulation over Sinai Peninsula Using the Weather Research and Forecasting Model, *International Journal of Atmospheric Sciences*, 2013 (2013) 11.
15. T. Islam, Performance of ecological revetment in Haor areas of Bangladesh, (2021).
16. S.J. Al -Hussain, "Identification of Drought and Flood Induced Critical Moments and Coping Strategies in Hazard Prone Lower Teesta River Basin" SUBMITTED BY, (2017).
17. M.T. Sikder, K.M. Elahi, Environmental Degradation and Global Warming-Consequences of Himalayan Mega Dams: A Review, *American Journal of Environmental Protection*, 2 (2013) 1–9.
18. M. Rahman, C. Ningsheng, G.I. Mahmud, M.M. Islam, H.R. Pourghasemi, H. Ahmad, J.M. Habumugisha, R.M.A. Washakh, M. Alam, E. Liu, Z. Han, H. Ni, T. Shufeng, A. Dewan, Flooding and its relationship with land cover change, population growth, and road density, *Geoscience Frontiers*, 12 (2021) 101224.
19. V.R. Sinha, Benefit sharing opportunities in the Meghna Basin Community management of protected areas in Bangladesh View project Community-based Fisheries Management Project Phase II View project, (n.d.).
20. N.C. Dey, M. Parvez, M.R. Islam, A study on the impact of the 2017 early monsoon flash flood: Potential measures to safeguard livelihoods from extreme climate events in the haor area of Bangladesh, *International Journal of Disaster Risk Reduction*, 59 (2021) 102247.
21. M. Khalequzzaman, Recent floods in Bangladesh: Possible causes and solutions, *Natural Hazards*, 9 (1994) 65–80.
22. A. Poncet, F. Gemenne, M. Martiniello, H. Bousetta, A country made for disasters: Environmental vulnerability and forced migration in Bangladesh, *Environment, Forced Migration and Social Vulnerability*, (2010) 211–222.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.