

# Kumquat (*Fortunella margarita*): A Good alternative for the Ingestion of Nutrients and Bioactive Compounds <sup>†</sup>

Clarice Silva e Souza <sup>1,\*</sup>, Pamella Cristine Anunciação <sup>1</sup>, Ceres Mattos Della Lucia <sup>1</sup>, Rosana Gonçalves Rodrigues das Dôres <sup>2</sup>, Regina Célia Rodrigues de Miranda Milagres <sup>1</sup> and Helena Maria Pinheiro Sant'Ana <sup>1,\*</sup>

<sup>1</sup> Department of Nutrition and Health, Federal University of Viçosa, Avenida P.H. Rolfs, s/n, Viçosa 36571-000, MG, Brazil; nutripamella@gmail.com (P.C.A.); ceresnut@yahoo.com.br (C.M.D.L.); rcrmmilagres@gmail.com (R.C.R.d.M.M.)

<sup>2</sup> Health Center, Federal University of Ouro Preto, University Campus, Morro do Cruzeiro. s/n, Ouro Preto 35400-000, MG, Brazil; rosanagr@gmail.com

\* Correspondence: cla\_souzabio@yahoo.com.br (C.S.eS.); helenapinheirosantana@gmail.com (H.M.P.S.)

<sup>†</sup> Presented at the 1st International Electronic Conference on Food Science and Functional Foods, 10–25 November 2020; Available online: [https://foods\\_2020.sciforum.net/](https://foods_2020.sciforum.net/).

Submitted: 10 July 2020; Accepted: date; Published: 9 November 2020

**Abstract:** Citrus fruit is preferred in the choice of consumers. Kumquat (*F. margarita*) is an unconventional citrus of increasing consumer interest because of its exotic flavor, and its functional potential that offers health benefits to consumers. It is a fruit traditionally consumed by whole fruit (peel and pulp), giving this fruit a distinctive flavor. For this reason, this study analyzed physical, chemical, and nutritional characteristics of kumquat (peel + pulp). The physicochemical analysis was performed according to the Adolfo Lutz Institute. Analysis of moisture, ashes, macronutrients, and total dietary fiber was according to AOAC. Minerals were analyzed by inductively coupled plasma optical emission spectrometry. Vitamins C and E, carotenoids and flavonoids were analyzed by HPLC. Phenolic compounds (Folin-Ciocalteu) and antioxidant capacity (DPPH) were determined by spectrophotometry. The kumquat had low pH, soluble solids content and low caloric value. It was a source of dietary fiber, minerals (K, Ca, P, Mg) and carotenoids, the most expressive was  $\alpha$ -carotene ( $661.81 \mu\text{g}\cdot 100 \text{g}^{-1}$ ). The ascorbic acid concentration was  $2326.24 \mu\text{g}\cdot 100 \text{g}^{-1}$ .  $\alpha$ -tocopherol ( $569.00 \mu\text{g}\cdot 100 \text{g}^{-1}$ ) was the most expressive component of vitamin E. There was a presence of apigenin and eriodictyol. The fruit (peel+pulp) has a high concentration of total phenolic compounds ( $98.55 \pm 1.93 \text{ mg GAE}\cdot 100 \text{g}^{-1}$ ) and good antioxidant capacity (62%) was found. Kumquat is a good source of fiber and vitamin A and due to its antioxidant capacity and the presence of other essential and beneficial nutrients for a diet, consumption of kumquat can be suggested to complement the diet. This fruit is a viable food alternative, and its consumption should be encouraged, contributing as a source of income, sovereignty, and food security.

**Keywords:** antioxidant; alternative food; citrus; exotic fruit; phenolics; vitamins

## 1. Introduction

Citrus fruits are among the largest producers and traders in the world [1]. Kumquat is the smallest existing Citrus, of Asian origin, specifically in China and India and belongs to the Rutaceae family [2]. The presence of this fruit is reported in rural backyards, in agroecological markets, as well as in the preparation of value-added foods such as jams, jellies, sweets and liqueurs [3,4]. However, it is still a little known and consumed fruit in Brazil, different from China, where the fruit is native

and popularly consumed in nature (peel + pulp). The commercialization of kumquat in Brazil is incipient in some states, being marketed or in large supermarkets as exotic fruits of high added value or in free markets present in small cities be it fruit or jellies, cakes, and other products. The kumquat is consumed preferably in natura, whole peel and pulp. However, few studies are described in the literature, so far, with the species *F. margarita* [5–8]. The fruit pulp showed antioxidant effects against the prevention of non-transmissible chronic diseases [8], activity against the inhibition of prostate cancer cells [9] and antimicrobial activity [5]. Due to its functional potential, it has increased the interest in the consumption of this fruit by Brazilians.

## 2. Materials and Methods

The preparation and analysis in kumquat (peel + pulp) were performed in four replicates. The fruits of each replicate were randomly selected for the measurement of length (cm), diameter (cm) and fruit weight (g) using analytical balance (Gehaka, AG200) and pachymeter (Disma, 150 mm). Analysis of pH, soluble solids and titratable acidity were performed according to Adolfo Lutz Institute [10], the lipids by the Soxhlet and the total ashes were quantified using a muffle (Quimis, model Q320 M, Brazil). The protein was determined by the Kjeldahl [11]. The carbohydrates content was calculated following the equation:  $[100 - (\% \text{ moisture} + \% \text{ lipids} + \% \text{ protein} + \% \text{ total fiber} + \% \text{ ash})]$ . The chemical elements (the minerals calcium-Ca, potassium-K, phosphorus-Pb, magnesium-Mg, sulfur-S, copper-Cu, iron-Fe, zinc-Zn, manganese-Mn, sodium-Na, chrome-Cr and the heavy metals cadmium-Cd, aluminum-Al, nickel-Ni and lead-Pb) were determined by inductively coupled plasma optical emission spectrometry (ICP-OES). The vitamin C, carotenoids and flavonoids were performed in a high performance liquid chromatography system (HPLC) (Shimadzu, SCL 10AT VP) detector (DAD) (Shimadzu, SPD-M10A); vitamin E used HPLC system (Shimadzu, SPD-M10A); fluorescence detector (290 nm excitation and 330 nm emission) (RF-10A XL). For each extraction and analysis followed an actor with modifications. Vitamin C [12]. Vitamin E components ( $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$ -tocopherols and tocotrienols) [13]. The  $\alpha$ -carotene,  $\beta$ -carotene and lutein followed the extraction methodology [14]. The 3-desoxyanthocyanidins (3-DXAs – luteolinidin, apigeninidin, 7-methoxyapigeninidin and 5-methoxy-luteolinidin), flavones (luteolin and apigenin) and flavanones (naringenin and eriodictiol) [15]. The total phenolic compounds were determined using the Folin-Ciocalteu reagent [16]. For antioxidant capacity was determination by methanolic DPPH solution (1.1-diphenyl-2-picrylhydrazyl).

## 3. Results and Discussion

The biometric analysis of the fruit shown in the (Figure 1). Study with kumquat reported values for fruit length and diameter from 2.0 and 3.0 cm and weight ranging from 5 to 20 g [17]. The kumquat of the present study had low pH. [7] found a pH of 4.22, higher than in the present study. The authors [7] analyzing different citrus fruits, reported the value of soluble solids in kumquat of 21.1 Brix and 8.35 Brix in 'Tahiti' lemon, being the concentration found in kumquat of the present study (16.41 Brix) smaller than that reported by the authors. Our result for soluble solids in kumquat was higher than [18] the titratable acidity of 2.43 mg of citric acid.100 g<sup>-1</sup> higher than the values found in citrus varieties by [19]. The centesimal composition shown in the (Table 1).



**Figure 1.** Images of kumquat fruits collected in Brazil. A: longitudinal and cross-section; B: weighing in semi analytical balance; C: processing. Source: Personal collection.

**Table 1.** Centesimal composition of kumquat (*F. margarita*) (peel + pulp) collected in Brazil.

Centesimal composition				
Moisture <sup>1</sup> (g·100 g <sup>-1</sup> )	Lipids <sup>1</sup> (g·100 g <sup>-1</sup> )	Total Ash <sup>1</sup> (g·100 g <sup>-1</sup> )	Protein <sup>1</sup> (g·100 g <sup>-1</sup> )	Carbohydrates <sup>1</sup> (g·100 g <sup>-1</sup> )
76.79 ± 0.98	1.18 ± 0.06	3.66 ± 0.18	7.38 ± 0.39	5.23 ± 0.30
Total fiber <sup>2</sup> (g·100 g <sup>-1</sup> )	Insoluble fiber <sup>2</sup> (g·100 g <sup>-1</sup> )	Soluble fiber <sup>2</sup> (g·100 g <sup>-1</sup> )	TEV <sup>3</sup> (kcal·100 g <sup>-1</sup> )	
5.31 ± 0.06	3.28 ± 0.15	2.03 ± 0.09	61.06	

<sup>1</sup> Data expressed as fresh basis, as mean ± standard deviation, <sup>2</sup> Data expressed as fresh basis, as mean ± standard deviation, <sup>3</sup> TEV–Total energy value.

The moisture was lower than that found for orange (*Citrus sinensis*) [1]. The ash, lipids and fiber were higher than orange (*C. sinensis*) [1]. Among the chemical elements, the most expressive in kumquat were K, Ca, Pb and Mg (Table 2). The concentration of AA in kumquat (Table 3) was lower than that observed in *C. sinensis* [1]. The concentration of vitamin C in Citrus is quite varied according to the species and maturation time, and the method of analysis used may also influence the total value of vitamin C. The α-tocopherol was the major vitamin E component found in kumquat.

**Table 2.** Composition of chemical elements present in kumquat (*F. margarita*) (peel + pulp) collected in Brazil.

Chemical Elements	Concentration (mg·100 g <sup>-1</sup> )
Phosphor	16.94 ± 0.23
Potassium	163.16 ± 3.29
Calcium	64.99 ± 1.41
Magnesium	16.71 ± 0.40
Sulfur	13.92 ± 0.23
Copper	0.07 ± 0.01
Iron	0.30 ± 0.06
Zinc	0.09 ± 0.00
Manganese	0.10 ± 0.00
Sodium	2.63 ± 0.00
Chrome	0.01 ± 0.33
Cadmium	0.00 ± 0.00
Aluminum	0.57 ± 0.33
Nickel	0.00 ± 0.00
Lead	0.00 ± 0.00

Data expressed as fresh basis, as mean of triplicates ± standard deviation.


(569.00 µg·100 g<sup>-1</sup>), followed by β-tocopherol (66.89 µg·100 g<sup>-1</sup>) (Table 3) (Figure 2). The total vitamin E concentration found in kumquat was almost three times higher than that found in orange (*C. sinensis*) [20]. There are few studies in the scientific literature relating the concentration of vitamin E in Citrus. The α-carotene was the most expressive carotenoid in kumquat (661.81 µg·100 g<sup>-1</sup>) (Table 3). Among the analyzed flavonoids, apigenin (38,157.30 µg·100 g<sup>-1</sup>) was detected in higher concentration in the present study (Table 3), one old study identified eight different classes of flavonoids in *F. margarita* but did not find apigenin in *Fortunella* sp. [21]. Other studies have demonstrated the presence of flavonoids [22, 23]. The concentration of total phenolic compounds in kumquat (98.55 ± 1.93 mg GAE·100 g<sup>-1</sup>) was much lower than the *F. crassifolia* collected in Taiwan, China (229 mg GAE·100g<sup>-1</sup>) [8]. The kumquat presented good antioxidant capacity (62%) (Table 3), In general, phytochemicals are present in different parts of the same fruit. As the peel is the part that gives protection to the fruit, it tends to have more amount of phenolic compounds and other bioactive

compounds in relation to the pulp [24]. (Thus, kumquat is a good alternative for planting, marketing, and consumption, which can contribute to food and nutritional sovereignty and security and provide a source of income for farming families.

**Table 3.** Occurrence and concentration of vitamins, carotenoids and bioactive compounds in kumquat (*F. margarita*) (peel+pulp) collected in Brazil.

<b>Components.</b>	<b>Concentration</b>
<b>Vitamin C</b> (mg·100 g <sup>-1</sup> )	
Ascorbic acid	2.32 ± 44.24
<b>Vitamin E</b> (µg·100 g <sup>-1</sup> )	
α-tocopherol	569.00 ± 10.20
α-tocotrienol	35.76 ± 4.03
β-tocopherol	nd
β-tocotrienol	66.89 ± 39.93
γ-tocopherol	4.22 ± 0.13
γ-tocotrienol	nd
δ-tocopherol	nd
δ-tocotrienol	nd
Total Vitamin E	675.87 ± 54.29
<b>Carotenoids</b> (µg·100 g <sup>-1</sup> )	-
α-carotene	661.81 ± 22.76
β-carotene	447.74 ± 19.90
Lutein	173.60 ± 33.61
Sum of carotenoids	1283.15
Vitamin A value (RAE 100 g <sup>-1</sup> ) <sup>1</sup>	129.77
<b>3-desoxyanthocianidins</b> (µg·100 g <sup>-1</sup> )	
Luteolinidin	nd
Apigeninidin	nd
5-methoxy-luteolinidin	nd
7-methoxy-apigeninidin	nd
<b>Flavones</b> (µg·100 g <sup>-1</sup> )	
Apigenin	38,157.30 ± 531.00
Luteolin	nd
Sum of flavones	38,157.30 ± 531.00
<b>Flavanones</b> (µg·100 g <sup>-1</sup> )	
Eriodictiol	36,880.95 ± 384.02
Naringenin	nd
Sum of flavanones	36,880.95 ± 384.02
<b>Total phenolics</b> (mg GAE·100 g <sup>-1</sup> )	98.55 ± 1.93
<b>Antioxidant capacity</b> (%)	62.01 ± 3.41

Data expressed as fresh basis, as an average ± standard deviation. <sup>1</sup> Equivalent of retinol activity, Nd: not detected.



**Figure 2.** Analysis by HPLC in kumquat (peel+pulp) collected in Brazil. Vitamin C (A); vitamin E (B); carotenoids (C); eriodictiol (D) and apigenin (E).

**Acknowledgments:** This work was conducted during a scholarship supported by the International Cooperation Program CAPES at the Universidade Federal de Viçosa, Minas Gerais, Brazil. Financed by CAPES, Brazilian Federal Agency for Support and Evaluation of Graduate Education within the Ministry of Education of Brazil and FAPEMIG, Fundação de Amparo à Pesquisa do Estado de Minas Gerais, Brazil.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. USDA. Food Composition Databases. National Nutrient Database for Standard, 2017. Available online: <https://ndb.nal.usda.gov/ndb/search> (accessed on 15 March 2020).
2. Donadio, L.C.; Mourão-Filho, F.A.A.; Moreira, C.S. Centros de origem, distribuição geográfica das plantas cítricas e histórico na citricultura no Brasil. *Citros* **2005**, *2*, 2–18.
3. Koller, O.L.; Soprano, E. Principais cultivares cítricos 1. Ed. In *Citricultura catarinense*, 1st ed.; Epagri: Florianópolis, Brazil, 2013; pp. 57–119.
4. Lazaroto, J.; Raiher, A.P. Determinantes da renda e pobreza dos agricultores do Vale do Ribeira. *Agric. Policy Rev.* **2013**, *22*, 5–25.
5. Wang, Y.C.; Chuang, Y.C.; Ku, Y.H. Quantitation of bioactive compounds in citrus fruits cultivated in Taiwan. *Food Chem.* **2007**, *102*, 1163–1171, doi:10.1016/j.foodchem.2006.06.057.

6. Schirra, M.; Palma, A.; D'aquino, S.; Angioni, A.; Minello, E.V.; Melis, M.; Cabras, P. Influence of postharvest hot water treatment on nutritional and functional properties of kumquat (*Fortunella japonica* Lour. Swingle Cv. Ovale) fruit. *J. Agric. Food Chem.* **2008**, *56*, 455–460, doi:10.1021/jf071416.
7. Oliveira, D.R.; Diniz, A.B. Composição química da laranja kinkan e de frutas cítricas. *DEMETRA Food Nutr. Health* **2015**, *10*, 835–844, doi:10.12957/demetra.2015.16726.
8. Chen, M.H.; Yang, K.M.; Huang, T.C.; Wu, M.L. Traditional small-size citrus from Taiwan: Essential oils, bioactive compounds and antioxidant capacity. *Medicines* **2017**, *4*, 28.
9. Jayaprakasha, G.K.; Murthy, K.C.; Etlinger, M.; Mantur, S.M.; Patil, B.S. Radical scavenging capacities and inhibition of human prostate (LNCaP) cell proliferation by *Fortunella margarita*. *Food Chem.* **2012**, *131*, 184–191.
10. Instituto Adolfo Lutz. *Normas Analíticas do Instituto Adolfo Lutz: Métodos Químicos e Físicos para Análises de Alimentos*, 4th ed.; Brasília, Brasil, 2005; Volume 1, 1018p.
11. Official Methods of Analysis. *Official Methods of Analysis of the AOAC International*, 19th ed.; Association of Official Analytical Chemists: Gaithersburg, MD, USA, 2012.
12. Campos, F.M.; Ribeiro, S.M.R.; Della Lucia, C.M.; Pinheiro-Sant'ana, H.M.; Stringheta, P.C. Optimization of methodology to analyze ascorbic and dehydroascorbic acid in vegetables. *Química Nova* **2009**, *32*, 87–91, doi:10.1590/S0100-40422009000100017.
13. Pinheiro-Sant'ana, H.M.; Guinazi, M.; Da Silva Oliveira, D.; Della Lucia, C.M.; De Lazzari Reis, B.; Brandão, S.C.C. Method for simultaneous analysis of eight vitamin E isomers in various foods by high performance liquid chromatography and fluorescence detection. *J. Chromatogr. A* **2011**, *1218*, 8496–8502, doi:10.1016/j.chroma.2011.09.067.
14. Rodriguez-Amaya, D.B. Assessment of the provitamin A contents of foods—The Brazilian experience. *J. Food Compos. Anal.* **1996**, *9*, 196–230, doi:10.1006/jfca.1996.0028.
15. Dykes, L.; Seitz, L.M.; Rooney, W.L.; Rooney, L.W. Flavonoid composition of red sorghum genotypes. *Food Chem.* **2009**, *116*, 313–317, doi:10.1016/j.foodchem.2009.02.052.
16. Singleton, V.L.; Orthofer, R.; Lamuela-Raventós, R.M. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods Enzymol.* **1999**, *299*, 152–178, doi:10.1016/S0076-6879(99)99017-1.
17. Illescas, J.L.; Bacho, O.; Ferrer, S. Análisis de los principales frutos tropicales comercializados. *Distrib. Y Consumo* **2007**, *95*, 33–85.
18. Ramful, D.; Tarnus, E.; Aruoma, O.I.; Bourdon, E.; Baborun, T. Polyphenol composition, vitamin C content and antioxidant capacity of Mauritian citrus fruit pulps. *Food Res. Int.* **2011**, *7*, 2088–2099.
19. Couto, M.A.L.; Guidolin Canniatti-Brazaca, S. Quantificação de vitamina C e capacidade antioxidante de variedades cítricas. *Ciência E Tecnol. De Aliment.* **2010**, *30*, 15–19.
20. Oliveira, R.P.; Schroder, E.C.; Souza, E.L.S.; Scivittaro, W.B.; Castro, L.A.S.; Rocha, P.S.G. Laranjeiras sem acidez. Pelotas: Embrapa Clima Temperado. *Embrapa Clima Temperado Documentos*, **2010**, *298*, 23.
21. Kawaii, S.; Tomono, Y.; Katase, E.; Ogawa, K.; Yano, M. Quantitation of flavonoid constituents in Citrus fruits. *J. Agric. Food Chem.* **1999**, *47*, 3565–3571.
22. Pereira, R.M.; López, B.G.C.; Diniz, S.N.; Antunes, A.A.; Garcia, D.M.; Oliveira, C.R.; Marcucci, M.C. Quantification of flavonoids in brazilian orange peels and industrial orange juice processing wastes. *Agric. Sci.* **2017**, *8*, 631, doi:10.4236/as.2017.87048.
23. Lou, S.N.; Ho, C.T. Phenolic compounds and biological activities of small-size citrus: Kumquat and calamondin. *J. Food Drug Anal.* **2017**, *25*, 162–175, doi:10.1016/j.jfda.2016.10.024.
24. Manzoor, M.; Anwar, F.; Saari, N.; Ashraf, M. Variations of antioxidant characteristics and mineral contents in pulp and peel of different apple (*Malus domestica* Borkh.) cultivars from Pakistan. *Molecules* **2012**, *17*, 390–407, doi:10.3390/molecules17010390.

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).