

Selected Thermodynamic Parameters of Antioxidant Activity of Coumarin Based Heterocyclic Compounds [†]

Ana Amić * and Elena Gotal

Department of Chemistry, Josip Juraj Strossmayer University of Osijek, Ulica cara Hadrijana 8A, 31000 Osijek, Croatia; gotal551@icloud.com

* Correspondence: aamic@kemija.unios.hr

† Presented at the 24th International Electronic Conference on Synthetic Organic Chemistry, 15 November–15 December 2020; Available online: <https://ecsoc-24.sciforum.net/>.

Published: date

Abstract: Coumarin and coumarin derivatives are bioactive compounds that have an important role in medicinal chemistry, for example in the development of anti-inflammatory, anticancer and antiviral drugs. These compounds are also very powerful antioxidants that successfully scavenge free radicals and prevent or alleviate oxidative stress. The antioxidant potential of selected heterocyclic compounds containing coumarin core was investigated theoretically, the focus of this study was on hydrogen atom transfer mechanism (HAT) and single electron transfer followed by proton transfer mechanism (SET-PT). Using MOPAC2012 PM7, reaction enthalpies related to the cleavage of O–H, N–H and C–H bonds *via* selected mechanisms of free radical scavenging were studied and calculated. The effect of the position of the hydroxyl group, as well as other functional groups, on the antioxidant activity was examined. Based on obtained results, Schiff bases, thiosemicarbazides, oxadiazoles and 4-thiazolidinones containing one or more OH– groups exhibit higher radical scavenging properties.

Keywords: coumarin; antioxidants; PM7; HAT; SET-PT

1. Introduction

Coumarin and coumarin compounds are a group of phenolic compounds that was first isolated from the beans of the Tonka plant, *Dipteryx odorata* (Aubl.) Wild, in 1820. These heterocyclic compounds containing oxygen are widely distributed in plant kingdom, where some are present in free form while others are present in the form of glycosides, e.g. bound with carbohydrate (for example glucose). These phytochemicals can be described as colorless crystalline compounds soluble in organic solvents [1,2]. Hundreds have been isolated from plants, the most widespread being umbelliferon, esculetin and scopoletin. Some plant species, such as dates, strawberries, apricots, cherries and cinnamon contain a very high amount of these compounds [3–7].

Due to their relatively simple structure, coumarins and their derivatives have various uses, for example in the perfume industry (due to the unique and intense scent similar to vanilla), as tobacco aroma enhancers, in some alcoholic beverages [8], in insecticides, as additives in soaps and toothpaste, as sweeteners, as additives to rubber and plastic materials, in aerosols, etc. [7,9]. Additionally, bioactivity of these compounds is especially diverse – coumarin and coumarin derivatives have shown anticancer, antitumor, antibacterial, antifungal and anti-inflammatory activity, some may inhibit activity of some enzymes and reduce concentration of triglycerides in blood. Hydroxycoumarins are very powerful antioxidants and have the ability to eliminate excess free radicals and reduce oxidative stress. Free radicals are constantly produced in our body for

metabolic purposes, but we are exposed to various additional sources of free radicals as well (ionizing radiation, UV light, pollution, etc.). Excess free radicals may attack biological macromolecules giving rise to membrane and DNA damage, and protein modifications [10]. This type of oxidative damage may be involved in pathogenesis of various chronic diseases, so to fight oxidative stress various exogenous antioxidants may be of use. In this way, some coumarin derivatives may play an important role in the prevention of chronic and age related diseases such as neurodegenerative and cardiovascular diseases [4].

2. Methods

MOPAC2012™ is a semiempirical quantum-chemical program based on Dewar and Thiel NDDO approximation. In this paper it was used to calculate reaction enthalpies of selected compounds with coumarin core in gas phase in order to study the thermodynamics of O–H, N–H and C–H bond cleavage necessary for antioxidant activity. Though application of other commercially available computational programs (such as Gaussian) may result in more accurate calculations of thermodynamic parameters, their disadvantage is their high price and long calculation time, depending on the computer, complexity of the compound structure and computational details (level of theory, basis set, functional, solvation model, etc.). Thermodynamic calculations run *via* PM7 method are less accurate, but are faster obtained and are similar to the results obtained *via* other computational programs. Hence MOPAC was used in this preliminary study, primarily because of very fast calculations and fast obtained results.

In this preliminary study, two mechanisms of antioxidant activity based on hydrogen atom abstraction were studied. Those mechanisms were hydrogen atom abstraction (HAT mechanism) and single electron transfer followed by proton transfer (SET-PT) mechanism. As can be seen on Figure 1, HAT mechanism is characterized by translocation of a hydrogen atom from the antioxidant molecule (A–OH) to the free radical (RO•). This mechanism occurs in one step and is characterized by thermodynamic parameter named bond dissociation enthalpy, or BDE [11].



Figure 1. HAT mechanism.

In HAT mechanism homolytic bond cleavage occurs, there is no charge separation, so it is preferred in gas phase and non-polar medium. Equation (1) shows how BDE can be calculated:

$$\text{BDE} = H(\text{A-O}\cdot) + H(\text{H}\cdot) - H(\text{A-OH}) \quad (1)$$

where $H(\text{A-O}\cdot)$, $H(\text{H}\cdot)$ and $H(\text{A-OH})$ are enthalpies of formation of antioxidant radical, transferred hydrogen atom, and antioxidant molecule. Lower H_{BDE} indicates a better ability of an antioxidant in donating hydrogen atom to the free radical [11,12].

SET-PT mechanism occurs in two steps as shown in Figure 2. In the first step of this mechanism, electron transfer takes place and neutral antioxidant molecule is modified into radical cation ($\text{AOH}^{\bullet+}$), while in the second step radical cation loses a proton and antioxidant radical is formed ($\text{AO}\cdot$) [11].




Figure 2. SET-PT mechanism.

In case of SET-PT, heterolytic bond cleavage occurs, there is charge separation and this mechanism is preferred in polar medium. This mechanism is characterized with thermodynamic parameters shown in Equations (2) and (3):

$$IP = H(A-O^{\cdot+}) + H(e^-) - H(A-OH) \quad (2)$$

$$PDE = H(A-O^{\cdot}) + H(H^+) - H(A-O^{\cdot+}) \quad (3)$$

Ionization potential, IP, is the energy necessary to release an electron and this value characterizes the first step of SET-PT mechanism. It can be calculated by subtracting the sum of the enthalpy of formation of radical cation and enthalpy of formation of an electron from the enthalpy of formation of the antioxidant molecule. The second step, in which antioxidant radical is formed, is characterized by proton dissociation enthalpy, or PDE, and can be calculated by subtracting the enthalpy of formation of radical cation from the sum of the enthalpy of formation of antioxidant radical and proton [11,13,14].

All above mentioned mechanisms of antioxidant activity have the same products – neutralized radical and a stable antioxidant radical. Accordingly, thermodynamic balance is presented with Equation (4), in which can be seen that reaction enthalpy BDE matches the sum of reaction enthalpies IP and PDE [11]:

$$\Delta_r H_{BDE} = \Delta_r H_{IP} + \Delta_r H_{PDE} \quad (4)$$


It is correct to say that there is a competitive relationship between these reactions and mechanisms, and there is also the possibility for these mechanisms to occur parallel [11].









3. Results and Discussion








Coumarins may reduce production of free radicals in the organism and stimulate scavenging of reactive oxygen species (ROS), meaning they may behave as antioxidants and act to diminish biological damage caused by oxidative stress. In order to determine the ability of compounds, namely phenolic antioxidants, to scavenge ROSs, O–H bond cleavage and potential to donate electron is studied. This implies calculation of BDE and IP. Lower BDE and IP lead to better antioxidant activity. For example, Mohajeri et al. (2009) found that BDE values in gas phase are in the range 69.9–104.0 kcal/mol for vitamins and 71.2–85.1 for phenolic acids. The authors also discovered that comparing BDE of vitamin E and C, the reduction of heterocyclic ring enhances antioxidant activity. Determined IP values for vitamins and phenolic acids were in the range 167.38–205.59 kcal/mol, leading to conclusion that both BDE and IP values were lower for phenolic acids than for vitamins, indicating phenolic acids as better antioxidants [15].









In this preliminary study, BDE and IP values for four groups of compounds with coumarin core were studied. Structures of compounds (1–51) and results of thermodynamic calculations of studied mechanisms of antioxidant activity are presented in Tables 1–3. Thermodynamic calculations refer to gas phase.

Table 1. Structures of studied Schiff bases (compounds 1–26) and results of thermodynamic calculations (kcal/mol).

No.	STRUCTURE	BDE	IP
1		341.6	772.2

2		375.5	831.8
3		336.6	778.1
4		334.5	785.7
5		346.3	773.0
6		335.1	755.1
7		311.1	747.3
8		335.8	848.5
9		319.4	761.7

10		299.1	787.1
11		316.9	749.8
12		315.3	787.4
13		319.6	736.2
14		350.9	750.6
15		352.9	852.2
16		285.5	684.5

17		331.2	797.8
18		314.9	793.8
19		354.5	820.0
20		338.5	824.6
21		338.4	819.2
22		351.3	816.3
23		324.7	795.6
24		341.8	784.5









25		333.9	763.6
26		301.2	668.0

Table 2. Structures of studied thiosemicarbazides and 4-thiazolidinones (compounds 27–36) and results of thermodynamic calculations (kcal/mol).

No.	STRUCTURE	BDE	IP
27		291.1	766.0
28		285.3	753.7
29		324.2	740.2
30		275.5	753.7
31		305.2	731.3
32		358.4	804.3

















33		314.0	761.3
34		359.7	689.3
35		359.3	690.4
36		358.9	668.0

Table 3. Structures of studied oxadiazoles (compounds 37–51) and results of thermodynamic calculations (kcal/mol).

No.	STRUCTURE	BDE	IP
37		294.3	782.1
38		258.5	799.4

39		269.5	801.3
40		300.9	801.5
41		296.9	801.0
42		299.5	802.0
43		254.4	807.7

44		298.1	804.7
45		296.8	798.7
46		293.5	797.5
47		297.7	804.3
48		261.7	793.9




49		299.1	796.1
50		298.5	797.7
51		282.3	759.0

Table 4 gives an overview of number of OH- and NH- groups and the number of vicinal OH-groups in the structure of studied compounds.

Table 4. Summary of main studied structural characteristics of analyzed compounds.

Compound	n-OH	n-Vicinal OH	n-NH
1	0	0	1
2	1	0	1
3	1	0	1
4	1	0	1
5	0	0	1
6	0	0	1
7	0	0	1
8	2	1	1
9	2	0	1
10	2	0	1
11	2	1	1
12	2	0	1
13	1	0	1
14	0	0	1
15	1	0	1
16	0	0	1
17	0	0	1
18	0	0	1

19	0	0	1
20	0	0	1
21	0	0	1
22	0	0	1
23	0	0	1
24	0	0	1
25	0	0	1
26	0	0	1
27	0	0	3
28	0	0	3
29	0	0	3
30	0	0	3
31	0	0	3
32	0	0	1
33	0	0	1
34	0	0	1
35	0	0	1
36	0	0	1
37	0	0	0
38	0	0	0
39	0	0	0
40	0	0	0
41	0	0	0
42	0	0	0
43	0	0	0
44	0	0	0
45	0	0	0
46	0	0	0
47	0	0	0
48	0	0	0
49	0	0	0
50	0	0	0
51	0	0	0

Table 5 gives a comparison of thermodynamic parameters for compounds that have OH- and NH- group (Schiff bases, compounds 2-4, 8-13, and 15).

Table 5. Comparison of BDE values for compounds containing OH- and NH- group.

COMPOUND	N-H BDE	O-H BDE
2	457.5	375.5
3	336.6	342.8
4	347.0	334.5
8	379.7	335.8
9	321.3	319.4
10	322.3	299.1
11	336.6	316.9
12	315.3	332.6
13	335.8	319.6
15	352.9	394.7

As can be seen in Table 4, some studied compounds have only one OH- group (2-14, 13, 15), some have two (8-12), and some have vicinal OH- groups (8 and 11). All these compounds are Schiff bases and all show better results of thermodynamic calculations. Some compounds contain one (1-26, 32-36) or more (27-31) NH- groups, but calculated BDE and IP values for these compounds were not better compared to the ones obtained for Schiff bases. Some compounds contain OH- and NH-

group (Tables 4 and 7, compounds 2–4, 8–13, and 15). These compounds are Schiff bases, and as can be seen from Table 7, the BDE of O–H bond is slightly more favorable compared to the BDE of N–H bond cleavage. Compounds 37–51 (oxadiazoles) do not have OH– or NH– group, hence for these compounds BDE and IP values for C–H bond was calculated. Obtained results do not show a better antioxidant potential for oxadiazoles, but they are not “much worse” antioxidants than compounds with OH– and NH– group.

BDE values for Schiff bases were determined to be in the range 285.5–373.5 kcal/mol, BDE values for thiosemicarbazides and 4-thiazolidinones were in the range 275.5–358.9 kcal/mol, and for oxadiazoles in the range 258.5–300.9 kcal/mol. All obtained results are almost twice higher than results obtained by Mohhajeri et al. (2009). These results indicate studied compounds as weaker antioxidants. Their inefficiency may be due to the fact that studied thiosemicarbazides, 4-thiazolidinones and oxadiazoles do not possess OH– group, while five Schiff bases have OH– group, five have two OH– groups, and only two have vicinal OH– groups.

Our study determined following IP values: 684.5–852.2 kcal/mol for Schiff bases, 668.0–804.3 kcal/mol for thiosemicarbazides and 4-thiazolidinones, and 782.1–807.7 kcal/mol for oxadiazoles. These results are in some cases almost four times higher than results obtained by Mohajeri et al. (2009) and clearly indicates lower antioxidant activity of studied compounds [15]. Overview of scientific literature related to the experimental determination of antioxidant activity (DPPH essay) [16–21] also indicates this group of compounds as not significant antioxidants and relates observed antioxidant activity with the number of OH– groups.

4. Conclusions

Coumarins are heterocyclic compounds widely used in medicinal chemistry and drug development, due to their anticancer, antibacterial and anti-inflammatory activity. Some of these compounds have a very powerful antioxidant activity that makes them useful in removal of reactive oxygen species and minimizing of oxidative damage of biological macromolecules. In this preliminary study, MOPAC and PM7 were implemented in order to determine thermodynamic parameters BDE and IP for heterocyclic compounds containing coumarin core (Schiff bases, thiosemicarbazides, oxadiazoles and 4-thiazolidinones). Two mechanism of antioxidant activity were studied, HAT and SET-PT. They are competitive mechanisms, with HAT being preferred in non-polar medium and SET-PT in polar medium. Calculations were done in gas phase, for which preferred mechanism is HAT. Thermodynamics of O–H, C–H and N–H bond cleavage was studied, as well as the effect of number and position of OH– and other functional groups (if there are any) on antioxidant activity. Obtained results indicate a very unfavorable BDE value and at least twice as big IP value of studied compounds. Results thus indicate that these compounds do not exhibit a very high antioxidant activity, what is probably due to the lack of structural motifs characteristic for superior antioxidant activity, such as OH– groups. It could also be concluded that antioxidant activity probably does not occur via studied mechanisms. Hence, in order to determine operative mechanism of antioxidant activity of these compounds, additional research is necessary, which would focus on other locations of antioxidant activity and other mechanisms by implementing other computational program packages.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Jain, P.K.; Joshi, H. Coumarin: Chemical and pharmacological profile. *J. Appl. Pharm. Sci.* **2012**, *2*, 236–240.
2. Bruneton, J. *Pharmacognosy, Phytochemistry, Medical Plants*, 2nd ed.; Intercept: Hampshire, UK, 1999.
3. Matilla, P.; Hellström, J.; Törrönen, R. Phenolic acids in berries, fruits, and beverages. *J. Agric. Food Chem.* **2006**, *54*, 7193–7199.
4. Wildman, R.E. *Handbook of Nutraceuticals and Functional Foods*, 2nd ed.; CRC Press: Boca Raton, FL, USA, 2016.
5. Aslam, K.K.; Khosa, M.K.; Jahan, N.; Nosheen, S. Short communication: synthesis and applications of Coumarin. *Pak. J. Pharm. Sci.* **2010**, *23*, 449–454.

6. Abraham, K.; Wöhrlich, F.; Lindtner, O.; Heinemeyer, G.; Lampen, A. Toxicology and risk assessment of coumarin: focus on human data. *Mol. Nutr. Food Res.* **2010**, *54*, 228–239.
7. Liu, B.; Raeth, T.; Beuerle, T.; Beerhues, L. A novel 4-hydroxycoumarin biosynthetic pathway. *Plant. Mol. Biol.* **2010**, *72*, 17–25.
8. Matos, M.J.; Santana, L.; Uriarte, E.; Abreu, O.A.; Molina, E.; Yordi, E.G. Coumarins—An Important Class of Phytochemicals. In *Isolation, Characterisation and Role in Human Health*, 1st ed.; Rao, V., Rao, L., Eds.; IntechOpen: London, UK, 2015; pp. 113–140.
9. Lake, B.G. Coumarin metabolism, toxicity and carcinogenicity: relevance for human risk assessment. *Food. Chem. Toxicol.* **1999**, *37*, 423–453.
10. Gamulin, S.; Kovač, Z.; Marušić, M. *Patofiziologija*, 7th ed.; Medicinska naklada: Zagreb, Croatia, 2011.
11. Jeremić, S.; Amić, A.; Stanojević-Pirković, M.; Marković, Z. Selected anthraquinones as potential free radical scavengers and P-glycoprotein inhibitors. *Org. Biomol. Chem.* **2018**, *16*, 1890–1902.
12. Prior, R.L.; Wu, X.; Schaich, K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J. Agric. Food Chem.* **2005**, *53*, 4290–4302.
13. Marković, Z. Study of the mechanisms of antioxidative action of different antioxidants. *J. Serbian Soc. Comput. Mech.* **2016**, *10*, 135–150.
14. Dimić, D.S. Eksperimentalno i teorijsko ispitivanje odnosa struktura-antiradikalna aktivnost odabranih neurotransmitera, njihovih prekursora i metabolita. Ph.D. Thesis, Sveučilište u Beogradu, Fakultet za fizikalnu kemiju, Beograd, Serbia, 2018.
15. Mohajer, A.; Somayeh Asemiani, S. Theoretical investigation on antioxidant activity of vitamins and phenolic acids for designing a novel antioxidant. *J. Mol. Struct.* **2009**, *930*, 15–20.
16. Čačić, M.; Molnar, M. Design, Synthesis and Characterization of Some Novel 3-Coumarinyl-5-arylidene-1,3-thiazolidine-2,4-diones and Their Antioxidant Activity. *Z. Naturforsch.* **2011**, *66b*, 177–183.
17. Čačić, M.; Molnar, M.; Šarkanj, B.; Has-Schon, E.; Rajković, V. Synthesis and antioxidant activity of some new coumarinyl-1, 3-thiazolidine-4-ones. *Molecules* **2010**, *15*, 6795–6809.
18. Čačić, M.; Molnar, M.; Strelec, I. Synthesis and biological evaluation of a novel series of 1, 3-dicoumarinyl-5-aryl-2-pyrazolines. *Heterocycles* **2011**, *83*, 1553–1566.
19. Šarkanj, B.; Čačić, M.; Molnar, M.; Grille, L. 4-Methyl-7-hydroxycoumarin antifungal and antioxidant activity enhancement by substitution with thiosemicarbazide and thiazolidinone moieties. *Food Chem.* **2013**, *139*, 488–495.
20. Al-Amiery, A.A.; Al-Majedy, Y.K.; Kadhum, A.A.H.; Mohamad, A.B. Hydrogen Peroxide Scavenging Activity of Novel Coumarins Synthesized Using Different Approaches. *PLoS ONE* **2015**, *10*, e0132175.
21. Molnar, M.; Amić, A.; Pavić, V.; Kovač, T.; Kovač, M.; Has-Schon, E. Biological study on novel coumarinyl 1, 3, 4-oxadiazoles. *Turk. J. Chem.* **2018**, *42*, 146–157.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).