

Lignocellulosic-based nanoparticles loaded with essential oils against *Staphylococcus aureus* and *Pseudomonas aeruginosa*-mediated infections

Joana Domingues¹, Maria Olívia Pereira², Helena P. Felgueiras¹, Joana C. Antunes^{1*}

¹Centre for Textile Science and Technology (2C2T), University of Minho, Portugal

²Centre of Biological Engineering (CEB), University of Minho, Portugal

*joana.antunes@2c2t.uminho.pt


Infected Diabetic Foot Ulcers (DFU's)

DFUs are a frequent and costly complication of diabetes, with limb amputation being highly prevalent worldwide. Persistent pathogens such as *Staphylococcus aureus* and *Pseudomonas aeruginosa* are the main microbial inhabitants of infected DFUs, often gaining antimicrobial-resistance to treatment. Nanoparticle (NP)-mediated therapies may overcome this problem, as they are able to carry and protect loads from biodegradation, be internalized by the cell, and release the load(s) in a controlled manner. As payloads, plant-derived essential oils (EOs) exert quick and strong bactericidal action. This work proposes cinnamon leaf oil (CLO) and clove oil (CO)-encapsulation into polyelectrolyte complexed (PEC) NPs fabricated with natural, renewable, and bactericidal polymers [quaternized cellulose (QC) and carboxymethyl lignin (CML)].¹



Polymer derivatives

QC was obtained through reaction with glycidyltrimethylammonium chloride (GTMAC), and the incorporation of the quaternized moieties was confirmed by the presence of a peak at 1482 cm^{-1} in FTIR spectra which corresponds to the methyl groups of cationic quaternary amines grafted onto the cellulose chain. CML was obtained via reaction with monochloroacetic acid, with the FTIR spectra showing two absorption bands at 1710 cm^{-1} and 1417 cm^{-1} corresponding to the introduction of the negatively charged carboxyl groups.


Quaternized cellulose



Carboxymethyl lignin




FTIR spectrum of QC




FTIR spectrum of CML

Polyelectrolyte complexes (PECs)

PECs were produced through electrostatic interaction between the opposite electrical charges of the polymers' derivatives: QC is positively charged and CML is negatively charged. EOs were blended with CML solution before complexation.




1635,6 µg of CLO entrapped into NPs (83% loading efficiency)



Morphology of QC/CLO/CML NP's (40x)


234,2 µg of CO entrapped into NPs (12% loading efficiency)



Morphology of QC/CO/CML NP's (40x)

Antibacterial potential

NPs functionalized with EOs have significant antibacterial potential against *S. aureus* and *P. aeruginosa*.¹



Conclusions

It was possible to produce PEC NPs with QC and CML through electrostatic interactions. CLO and CO were successfully encapsulated into QC/CML NPs.

References

¹Miranda, C. S., Antunes, J. C., Homem, N. C. & Felgueiras, H. P. Controlled Release of Cinnamon Leaf Oil from Chitosan Microcapsules Embedded within a Sodium Alginate/Gelatin Hydrogel-Like Film for *Pseudomonas aeruginosa* Elimination. Proceedings 69, 39 (2020).

Acknowledgements

Authors acknowledge FCT for supporting the projects with references PTDC/CTM-TEX/28074/2017(POCI-01-0145-FEDER-028074) and UID/CTM/00264/2021.

Joana Domingues is also grateful for the FCT PhD grant 2020.07387.BD.