

Characterization musts, wines and sparkling wines based on their elemental composition determined by ICP-OES and ICP-MS

Biel Granell^a, Anaïs Izquierdo-Llopart^a, Àngels Sahuquillo^a, J. Fermín López-Sánchez^a, Javier Saurina^{a,b}

^a Department of Chemical Engineering and Analytical Chemistry, University of Barcelona, Martí i Franquès 1-11, E08028 Barcelona, Spain.

^b Research Institute in Food Nutrition and Food Safety, University of Barcelona, Av. Prat de la Riba 171, Edifici Recerca (Gaudí), E08921 Santa Coloma de Gramenet, Spain.

The control of the composition of food products with Protected Designation of Origin, such as wines and sparkling wines, is essential to prevent fraudulent practices and adulterations. A wide range of compounds can be used as tentative biomarkers for characterization and authentication purposes, being elemental composition one of the most successful sources of information, especially for dealing with geographical origin and varietal issues. Currently, Inductively Coupled Plasma with Optical Emission Spectrometry (ICP-OES) and Inductively Coupled Plasma Mass Spectrometry (ICP-MS) are the techniques of choice to carry out multi-elemental analysis of this kind of beverages in a rapid and simple way.

In our study, samples from the different processing stages in the elaboration of sparkling wine (*cava*)—including must, base wine and sparkling wine— of Pinot Noir and Xarel·lo grape varieties have been analyzed by ICP techniques to determine the elemental composition. The resulting data has been used to classify these products according to oenological practices and product qualities. For this purpose, Principal Components Analysis, box plot diagrams and bar charts have been used. Different markers and sample patterns have been found dealing with changes resulting from the different steps of the production process *cava* wines.

Results have revealed the relevance of some elements as descriptors of winemaking processes. For instance, Cu and K are abundant in musts and their concentrations progressively decreases through the *cava* production process. S levels suddenly increase at the base wine step (and further decay) of the addition of sulphites as preserving species. Finally, concentrations of Na, Ca, Fe and Mg increase from the first fermentation due to the addition of clarifying agents such as bentonite.