

# Assessment of the Relationship between the Presence of Vitamin D-Containing Foods in the Diet of Young People and the Development of Allergic Diseases <sup>†</sup>

Elena Aksonova <sup>1</sup>, Dmitry Torianik <sup>1</sup>, Natalia Otroshko <sup>2</sup> and Sergey Gubsky <sup>1,\*</sup>

<sup>1</sup> Department of Chemistry, Biochemistry, Microbiology and Hygiene of Nutrition, State Biotechnological University, 61051 Kharkiv, Ukraine; eaksonova@gmail.com (E.A.); datory@gmail.com (D.T.)

<sup>2</sup> Derzhavin Tambov State University, Russia; otroshko@mail.ru

\* Correspondence: sergey.m.gubsky@gmail.com

<sup>†</sup> Presented at the 2nd International Electronic Conference on Foods, 15–30 October 2021; Available online: <https://foods2021.sciforum.net/>.

**Abstract:** A significant part of vitamin D, which is necessary for important vital processes, a person receives with food. Vitamin D affects the proper functioning of the lungs and the immune system, which can be important in the development, severity and course of allergic diseases (asthma, eczema and food allergies). The purpose of this study was to investigate the relationship between the presence of vitamin D-containing foods in the diets of young people and the appearance of allergic diseases symptoms. The study was a cross-sectional descriptive survey of 223 first year student of the Kharkiv State University of Food Technology and Trade (Ukraine). Using questionnaires, the frequency of consumption and food preferences were assessed in relation to 22 vitamin D-containing foods and for the screening of allergic diseases. Analysis of the results showed that the consumption of foods rich in vitamin D is at a lower level than the respondents' desire to consume them. It was revealed that the main food source of vitamin D is chicken eggs. This fact speaks in favor of the fact that young people in Ukraine experience a certain lack of vitamin D. Analysis of questionnaires for screening allergic diseases showed that, depending on the year of the survey, about 69% of the respondents had symptoms of allergic diseases. The survey suggests that foods rich in vitamin D are not among the preferences of young people. As a consequence, there is a definite relationship between the presence of vitamin D deficiency in young people and a high percentage of respondents with allergic diseases symptoms.

**Keywords:** vitamin D-containing foods; intake of vitamin D; descriptive survey; allergic diseases; food preferences

**Citation:** Aksonova, E.; Torianik, D.; Otroshko, N.; Gubsky, S.

Assessment of the Relationship between the Presence of Vitamin D-Containing Foods in the Diet of Young People and the Development of Allergic Diseases. *Biol. Life Sci. Forum* **2021**, *1*, x.

<https://doi.org/10.3390/xxxxx>

Published: date

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Vitamin D is a fat-soluble vitamin, which is necessary for important vital processes. Vitamin D affects the proper functioning of the lungs and the immune system, which can be important in the development, severity and course of allergic diseases (asthma, eczema and food allergies). The COVID-19 pandemic has further increased interest in vitamin D as studies have emerged that link the severity of the disease to vitamin D levels in the human body [1]. Therefore, it is suggested that increasing intake of vitamin D in different forms may reduce the risk and severity of COVID-19 infection [2,3].

The uniqueness of this vitamin is associated with the duality of its intake into the human body. The first path is associated with the synthesis of vitamin D by the human body under the influence of sunlight. And the second way is associated with the consumption of foods containing this vitamin. However, the diet of a modern person is characterized by the presence in it of highly processed food, as a rule, deficient in terms of vitamin D. Research data suggests that, vitamin D insufficiency affects almost 50% of the

population worldwide. An estimated one billion people worldwide, across all ethnicities and age groups, have a vitamin D deficiency [4].

The duality of vitamin D intake in the human body is a significant obstacle to the formulation of diets with adequate intake of this vitamin. At the same time, the analysis of the diet of the population is a rather serious tool for determining the level of vitamin D intake and other aspects of nutritional status and nutritional behavior. There is experience in conducting such studies in Czech [5] and Germany [6] populations. Studies have shown that even people do not reach the current recommended vitamin D intake level. In Ukraine, in recent years, there is practically no reliable open scientific information regarding research in the field of analysis of the diets of the population, including young people. In this regard, an attempt was made to pilot study on a small sample of respondents for an assessment of the relationship between the presence of vitamin D-containing foods in the diet of young people and the development of allergic diseases. The original hypothesis assumed the existence of such a relationship.

## 2. Methods

### 2.1. Desing of Questionnaire

Data collection was carried out by conducting two parallel questionnaires, using questionnaires A and B. Using questionnaire A, the frequency of consumption and food preferences were assessed for 22 vitamin D-containing foods (Table 1).

**Table 1.** Vitamin D-containing foods of research.

Cluster	Sample Code	Food	Amount of Vitamin D, $\mu\text{g}/100\text{ g}$ [7]	Consumption Frequency	Preferences	Cluster Average	
						Consumption Frequency	Preferences
1	10	Chicken egg	2.2	3.13	2.94	3.05	3.09
	22	Cheese	1	2.96	3.23		
2	4	Sprats in oil (canned)	20.5	1.90	2.44	1.91	2.54
	6	Mackerel	16.1	2.10	2.62		
	7	Atlantic salmon	11	1.58	2.54		
	15	Caviar red granular	2.9	1.53	2.65		
	20	Butter	1.3	2.42	2.45		
3	3	Salty fat herring	30	1.54	1.87	1.32	1.95
	11	Tuna	5.7	1.25	2.03		
	12	Chanterelles	5.3	1.30	2.08		
	13	Quail egg	1.4	1.23	1.96		
	14	River perch	3	1.46	1.94		
4	17	Pike	2.5	1.13	1.84	0.97	1.60
	2	Codfish liver (canned)	100	0.95	1.51		
	8	Humpback salmon	10.9	1.00	1.61		
	9	Caviar black granular	8	0.77	1.83		
	16	Flounder	2.8	0.87	1.56		
5	18	Sea bass	2.3	0.98	1.58	0.52	0.74
	19	Ghee	1.8	1.27	1.50		
	1	Fish Oil	250	0.46	0.44		
5	5	Dog-salmon	16.3	0.39	0.79	0.52	0.74
	21	Goat's milk	1.3	0.71	1.00		

The assessment of the taste preferences to the above-mentioned products was carried out on a 5-point rating scale: "very much" (4), "like" (3), "I have nothing against" (2),

“rather not like than like” (1), “not at all like” (0). The evaluation of the frequency of consumption in food also occurred on a 5-point rating scale: with anchors “I eat very often” (4), “I eat, but not often” (3), “sometimes I eat” (2), “very rarely I eat” (1), “I don’t eat at all” (0). Using questionnaire B, the identify allergic diseases according method [8] than developed by the Ministry of Health of Ukraine were assessed.

## 2.2. Questionnaire Distribution and Data Collection

The study was a cross-sectional descriptive survey. The respondents included in the study were 223 students of the Kharkiv State University of Food Technology and Trade (Ukraine) aged 17 to 21, of which 30% were women and 70% were men. The assessment procedure was carried out in 2017–2021 among first-year students who voluntarily agreed to take part in the study. The overwhelming majority of participants in the study lived before admission to university in the Kharkiv region.

## 2.3. Data Analysis

Before analysis of data the reliability was carried out. Reliability within the study was considered as the degree of consistency between two independently obtained series of indicators within the framework of a statistical approach using Pearson correlation coefficients. The assessment of reliability as a measure of the internal consistency of the results was carried out using the Cronbach alpha method [9], within which the spread of each element was compared with the overall scale range. The obtained coefficients of 0.924 and 0.928 testify to the reliability of the data obtained on consumption frequency and preferences of vitamin D-containing foods.

The hierarchical cluster analysis (HCA) was performed to classify the food samples based on the similarities of their frequency of consumption and preferences. The linkage method, which was applied here, was Ward’s method. Clusters were calculated using the Euclidian distance. For this purpose, the data were centered and divided by their standard deviation to ensure that all the elements had equal weight in the results. Moreover, the statistical significance of the obtained clusters was tested by various formal and empirical statistical criteria: ratios of agglomeration and cross-tabulation and adjusted  $R^2_{adj}$  calculated by (1) the expression as in research [10]:

$$R^2_{adj} = 1 - WSS(n - 1)/TWSS(n - k), \quad (1)$$

where WSS is within the cluster the sum of squares of distances; TWSS is within sampling the total sum of the squares of distances; n is total samples; k is number of clusters. The HCA analysis was performed by using and Minitab ver. 18 (Minitab Inc., USA) software.


## 3. Results and Discussions

### 3.1. Data Analysis of Frequency of Consumption and Preferences of Vitamin D-Containing Foods

Based on the data from the questionnaires, the respondents’ average consumption and preferences were calculated for all these 22 vitamin D-containing foods (Table 1). The obtained values of consumption and preferences correlate with each other with Pearson correlation  $r = 0.887$  (Figure 1a). Data analysis allows us to trace several patterns:

1. Consumption of products containing vitamin D is below the possible level of consumption, which should arise based on the preferences values of the respondents;
2. Only for three products chicken eggs (10), fish oil (1) and butter (20) is there a correspondence between consumption and preferences;
3. Products Fish Oil (1), Dog-salmon (5), Humpback salmon (8), Caviar black granular (9), Goat’s milk (21) have the lowest score according to both Food preferences and Frequency of consumption;
4. The leading product in terms of frequency of consumption is chicken eggs (on average 3.13 points, which corresponds to the definition “I eat, but not often”).

- The leading product in Food preferences is cheese (3.1 points on average, which corresponds to the definition of “like”).
- The same high scores, both in the Frequency of consumption category and in the Food preferences category, have products 4 (Sprats in oil (canned)), 6 (Mackerel), 7 (Atlantic salmon), 15 (Caviar red granular), 20 (Butter).



**Figure 1.** (a) Pearson correlation between consumption frequency and preferences of vitamin D-containing foods; (b) Hierarchical cluster analysis dendrogram with four distinguished vitamin D-containing foods.

HCA analysis was performed to classify samples based on the similarity of consumption frequency and food preferences for foods containing vitamin D (Figure 1b). Based on this analysis, all 22 products were divided into clusters (Table 1). The resulting clusters actually reflect the ratio of Preferences and Consumption of the studied foodstuffs containing vitamin D. A retrospective review of the obtained clustering allows us to highlight some facts:

- The distribution of 22 vitamin D-containing food by clusters is obtained as a ratio 2:5:6:6:3;
- Foods containing the highest amount of vitamin D belong to clusters 4 and 5, ie. and less attractive to students, and less consumed;
- The most useful product Fish oil (1) is included in the cluster of the least consumed foods and with the least preference;
- Cluster number 1 with the highest values of consumption and preference is made up of traditional products in the diets of the population of Ukraine, but with a low content of vitamin D;
- Cluster number 1 is the only one in which the consumption meets the preference, which partly reflects the specifics of the student’s diet.

### 3.2. Results of Processing the Data of Questionnaire B

Questionnaire B showed that, on average, about 70% of all respondents have signs of various types of allergic diseases caused by substances of different origins: natural allergens (for example, seasonal flowering of plants that cause allergic reactions), the presence of food allergens in their diets, as well as allergens anthropogenic origin (including household chemicals, pharmaceuticals). The average value for the time period 2017–2021 of the number of respondents using vitamins—10.3% (Table 2).

**Table 2.** Results of the analysis of the data of questionnaire B.

Percent of Respondents		Consumption Frequency (Average Score)	
With Symptoms of Allergic Diseases	Using Vitamin D Supplements	Respondents without Symptoms of Allergic Diseases	Respondents with Symptoms of Allergic Diseases
68.8	10.3	1.59	1.49

However, during this period, there was a positive trend regarding an increase in the percentage of young people using vitamin D supplements, which correlates with a decrease in the number of allergic manifestations. The increased consumption of vitamin D supplements may be related to information in the media about its positive effect on immunity and the severity of COVID-19, as interest in vitamin D has increased since 2019 among the population.

Analysis of the B survey data also showed that there is a certain relationship between the low frequency of consumption of foods containing high concentrations of vitamin D and the presence of allergic diseases in the respondents. The respondents who, as a result of the questionnaire, did not reveal signs of allergic diseases have a higher average score for the frequency of consumption of foods rich in vitamin D (Table 2). This is especially true in 2017 and 2018.

In general, the tendency revealed as a result of the study is quite alarming, since in the studied cohort of young people, on the one hand, there is a low level of consumption of foods rich in vitamin D, and on the other hand, a high percentage of respondents with allergic diseases. At the same time, the percentage of consumption of food supplements with vitamin D is low. In addition, it was shown that in this sample of respondents, the frequency of consumption of foods rich in vitamin D in those with symptoms of allergic diseases is lower 1.49 than in those who do not have these symptoms. The latter fact may speak of the beneficial effect of vitamin D on the scenario of the development of allergic diseases.

#### 4. Conclusions

The survey suggests that foods rich in vitamin D are not among the preferences of young people, with the exception of foods such as eggs and cheese. Different types of fish and seafood containing the highest concentrations of vitamin D have rather low scores both in terms of frequency of consumption and food preferences. It should also be noted that the scores are quite high both in relation to the frequency of consumption and in relation to food preferences for butter. Despite the fact that Ukraine has a very poorly developed tendency to fortification of products with vitamin D, however, this product could be very convenient for these purposes. Since, as studies have shown, it is well accepted by young people, is traditional for Ukraine and is excellent for fortification with fat-soluble vitamins, including vitamin D. In addition, when introducing vitamin D fortification technology in Ukraine, the rich world experience of oil fortification can be used. cheese and dairy products.

Undoubtedly, it is impossible to draw an unambiguous conclusion about the dependence of the presence or development of allergic diseases and the frequency of consumption of foods rich in vitamin D on the basis of the above data. However, it should be recognized that a certain dependence exists. These studies require continuation and conduct of the questionnaire on a wider scale. In addition, in our opinion, in future studies, in order to assess the correlation between the presence of allergic diseases and the consumption of foods rich in vitamin D, the analysis of diets should also be taken into account, indicating the exact amounts of food consumed. Also in the same study, the questionnaires should include questions about the duration of exposure to the sun, depending on the season.

**Supplementary Materials:** The following are available online at [www.mdpi.com/xxx/s1](http://www.mdpi.com/xxx/s1), Figure S1: title, Table S1: title, Video S1: title.

**Author Contributions:** Conceptualization, E.A., S.G., S.B. and F.F.S.; methodology, E.A. and S.G.; software, D.T. and S.G.; validation, D.T.; formal analysis, E.A. and N.O.; writing—original draft preparation, E.A. and S.G.; writing—review and editing, E.A., N.O. and S.G. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** data presented in this study are available on request from the 108 corresponding author.

**Acknowledgments:** In this section, you can acknowledge any support given which is not covered by the author contribution or funding sections. This may include administrative and technical support, or donations in kind (e.g., materials used for experiments).

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Cui, C.; Xu, P.; Li, G.; Qiao, Y.; Han, W.; Geng, C.; Liao, D.; Yang, M.; Chen, D.; Jiang, P. Vitamin D Receptor Activation Regulates Microglia Polarization and Oxidative Stress in Spontaneously Hypertensive Rats and Angiotensin II-Exposed Microglial Cells: Role of Renin-Angiotensin System. *Redox Biol.* **2019**, *26*, 101295, <https://doi.org/10.1016/j.redox.2019.101295>.
2. Grant, W.; Lahore, H.; McDonnell, S.; Baggerly, C.; French, C.; Aliano, J.; Bhattoa, H. Evidence That Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths. *Nutrients* **2020**, *12*, 988, <https://doi.org/10.3390/nu12040988>.
3. Mendy, A.; Apewokin, S.; Wells, A.A.; Morrow, A.L. Factors Associated with Hospitalization and Disease Severity in a Racially and Ethnically Diverse Population of COVID-19 Patients. *medRxiv Prepr. Serv. Heal. Sci.* **2020**, <https://doi.org/10.1101/2020.06.25.20137323>.
4. Nair, R.; Maseeh, A. Vitamin D: The Sunshine Vitamin. *J. Pharmacol. Pharmacother.* **2012**, *3*, 118, <https://doi.org/10.4103/0976-500X.95506>.
5. Bischofova, S.; Dofkova, M.; Blahova, J.; Kavrik, R.; Nevrla, J.; Rehurkova, I.; Ruprich, J. Dietary Intake of Vitamin D in the Czech Population: A Comparison with Dietary Reference Values, Main Food Sources Identified by a Total Diet Study. *Nutrients* **2018**, *10*, 1452. <https://doi.org/10.3390/nu10101452>.
6. Jungert, A.; Spinneker, A.; Nagel, A.; Neuhäuser-Berthold, M. Dietary Intake and Main Food Sources of Vitamin D as a Function of Age, Sex, Vitamin D Status, Body Composition, and Income in an Elderly German Cohort. *Food Nutr. Res.* **2014**, *58*, 23632, <https://doi.org/10.3402/fnr.v58.23632>.
7. Skurikhin, I.M.; Volgarev, M.N. (Eds.) *Chemical Composition of Food Products. Book 2: Reference Tables of Amino Acids, Fatty Acids, Vitamins, Macro- and Micronutrients, Organic Acids and Carbohydrates*; Agropromizdat: Moskow, Russia, 1987.
8. *About Organizational Measures for Introduction of Modern Technologies of Diagnostics and Treatment of Allergic Diseases*; Order N 127/18; The Ministry of Health of Ukraine: Kyiv, Ukraine, 2002.
9. Cronbach, L.J. A Case Study of the Split-Half Reliability Coefficient. *Psychometrika* **1951**, *16*.
10. Goralchuk, A.; Gubsky, S.; Omel'chenko, S.; Riabets, O.; Grinchenko, O.; Fedak, N.; Kotlyar, O.; Cheremskaya, T.; Skrynnik, V. Impact of Added Food Ingredients on Foaming and Texture of the Whipped Toppings: A Chemometric Analysis. *Eur. Food Res. Technol.* **2020**, *246*, 1955–1970, <https://doi.org/10.1007/s00217-020-03547-3>.