

**ASEC
2021**

The 2nd International Electronic Conference on Applied Sciences

15-31 OCTOBER 2021 | ONLINE

Detection of SARS-CoV-2 by plasmonic optical fibers and molecularly imprinted polymers

Chiara Perri^{1*}, Nunzio Cennamo¹, Girolamo D'Agostino², Francesco Arcadio¹, Guido Chiaretti², Eva Maria Parisio³,
Giulio Camarlinghi³, Chiara Vettori³, Francesco Di Marzo⁴, Rosario Cennamo⁴, Giovanni Porto² and Luigi Zeni¹

¹ Department of Engineering, University of Campania *L. Vanvitelli*, Aversa, Italy

² Moresense srl, Milan, Italy

³ Operative Unit of Chemical-Clinical and Microbiological Analysis, San Luca Hospital, Usl Toscana Nord Ovest, Lucca, 55100, Italy

⁴ UOC Chirurgia Generale, Ospedale Valtiberina, Usl Toscana Sud-Est, Sansepolcro, 52037, Italy

*Email: chiara.perri@unicampania.it




Detection of SARS-CoV-2 by plasmonic optical fibers and molecularly imprinted polymers


Outline

- Plasmonic sensor based on a D-shaped Plastic Optical Fiber (POF)
- Molecularly Imprinted Polymer (MIP) for SARS-CoV-2 detection
- Experimental setup
- Experimental results
- Conclusions


Plasmonic sensor based on a D-shaped Plastic Optical Fiber (POF)



Molecularly Imprinted Polymer (MIP) for SARS-CoV-2 detection in aqueous solution



Outline of the Sensor for SARS-CoV-2 detection in different aqueous matrices.



Schematic representation of the MIP synthesis process.




Software

Spectrum analyzer
(200 ÷ 850 nm or 300 ÷ 1050 nm)



Experimental setup




White Light Source
(360 ÷ 2000 nm)



SMA connectors

Experimental results


Preliminary binding test with S1 subunit of SARS-CoV-2 spike protein




Response curves of different concentrations of SARS-Cov-2 Spike S1 subunit.

Experimental results

SARS-CoV-2 detection in real nasopharyngeal swabs



Response curves of different dilutions of a SARS-CoV-2 Positive swab (36th RT-PCR cycle) collected in UTM, tested with SARS-CoV-2 MIP-sensor. Physiological solution was used to make dilutions.



Response curves of different dilutions of a SARS-CoV-2 Positive swab (36th RT-PCR cycle) collected in physiological medium, tested with SARS-CoV-2 MIP-sensor. Physiological solution was used to make dilutions.

Prospects and Conclusions

A plasmonic sensor based on D-shaped POF and MIP for SARS-CoV-2 detection has been reported.

Tests on real nasopharyngeal swabs in two different matrices (UTM and physiological solution) in order to investigate the sensitivity in both media, were reported.

These preliminary results demonstrated the effectiveness of the proposed sensing approach in detecting the virus in real samples within few minutes, making this kind of sensor a promising tool for a more rapid, cost-effective detection system in diagnostics.

Thank you for your attention



Chiara Perri

Department of Engineering,
University of Campania *Luigi Vanvitelli*,
Via Roma, 29
81031 Aversa, Italy
E-mail: chiara.perri@unicampania.it