

IECPS
2021

The 2nd International Electronic Conference on Plant Science

01-15 DECEMBER 2021 | ONLINE



Chaired by **DR. ADRIANO SOFO**



Mineral quantification of *Triticum aestivum* L. enriched in Zinc – correlation between minerals in soils and whole wheat flours

Inês Carmo Luís ^{1,2,*}, Cláudia Campos Pessoa ^{1,2}, Diana Daccak ^{1,2}, Ana Coelho Marques ^{1,2}, Ana Rita F. Coelho ^{1,2}, Manuel Patanita ^{2,3}, José Dôres ³, Ana Sofia Almeida ^{2,4}, Maria Manuela Silva ^{2,5}, Maria Fernanda Pessoa ^{1,2}, Fernando H. Reboredo ^{1,2}, Manuela Simões ^{1,2}, Paulo Legoinha ^{1,2}, Carlos Galhano ^{1,2}, Isabel P. Pais ^{2,6}, Paula Scotti Campos ^{2,6}, José C. Ramalho ^{2,7} and Fernando C. Lidon ^{1,2}

1 Earth Sciences Department, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Campus Caparica, 2829-516 Caparica, Portugal

2 GeoBioTec Research Center, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Campus Caparica, 2829-516 Caparica, Portugal

3 Escola Superior Agrária, Instituto Politécnico de Beja, R. Pedro Soares S/N, 7800-295 Beja, Portugal

4 Instituto Nacional de Investigação Agrária e Veterinária, I.P. (INIAV), Estrada de Gil Vaz 6, 7351-901 Elvas, Portugal

5 ESEAG-COFAC, Avenida do Campo Grande 376, 1749-024 Lisboa, Portugal

6 Instituto Nacional de Investigação Agrária e Veterinária, I.P. (INIAV), Avenida da República, Quinta do Marquês, 2780-157 Oeiras, Portugal

7 PlantStress & Biodiversity Lab, Centro de Estudos Florestais (CEF), Instituto Superior Agronomia (ISA), Universidade de Lisboa (ULisboa), Quinta do Marquês, Av.

República, 2784-505 Oeiras, Portugal

* Correspondence: idc.rodrigues@campus.fct.unl.pt



geobiotec
Geobiociências, Geoengenharias
e Geotecnologias



INSTITUTO
SUPERIOR D
AGRONOMIA 1
Universidade de Lisboa

Mineral quantification of *Triticum aestivum* L. enriched in Zinc – correlation between minerals in soils and whole wheat flours



ANALYZES

- pH
- Electrical conductivity
- Organic Matter Content
- Moisture Content



Soils

Zn K
 Ca



Whole bread
wheat flours

Cu P Mg

Fe

Triticum aestivum L.
(cvs. Paiva and Roxo)

Abstract: *Triticum aestivum* L. is one of the most produced staple crops worldwide in which its zinc biofortification is of the utmost importance to diminish malnutrition. In addition, the pronounced increase of human population demands a higher food production within quality standards. Zinc plays an important role not only in promoting the maintenance of human health, but it is also linked with plant growth. Under this framework, a zinc agronomic biofortification of *Triticum aestivum* L. was implemented in an experimental field with two varieties (Paiva and Roxo) in Beja, Portugal. This itinerary comprised two ZnSO₄ foliar spraying along the plant cycle with three different concentrations (control – 0; 8.1 and 18.2 kg ha⁻¹). Soil analyses (moisture, organic matter, pH, electrochemical conductivity and mineral quantification) and atomic absorption with the mineral quantification (Ca, K, Mg, P, Fe, Cu and Zn) of whole wheat flours were carried out. Zinc foliar spraying enhanced Zinc content in both varieties in the flours in which was not observed significant differences between ZnSO₄ treatments. P and K presented higher values in the flours contrasting with Ca and Mg. In general, there was no significant differences between the soil samples in the respective analyses. It was concluded that wheat flour biofortified in zinc can be a product to help overcome malnutrition.

Keywords: agronomic biofortification; mineral interactions; soil analyses; *Triticum aestivum* L; zinc foliar spraying

Introduction

The world's population is expected to increase to more than 8 billion by 2030 (United Nations, 2019). Thereby, according to (FAO, 2021) the food production will have to increase approximately 60 % by 2050, in a sustainable way and keeping quality standards. *Triticum aestivum* L. is one of the most produced staple crops worldwide. Thus, it is estimated to reach a production of 776.7 million tons by 2021/2022 (FAO, 2021). Zinc (Zn) plays an important role (at the function, structure and regulation level) not only in promoting the maintenance of human health, but it is also linked with plant growth (Begum *et al.* 2016; Cakmak and Kutman, 2018). Biofortification is likely to diminish malnutrition figures, provided that an essential nutrient in the edible part of staple crops is enhanced and becomes bioavailable (Bouis and Saltzman, 2017; Beaudreault, 2019). This paper aims to analyze the correlations between the minerals present in the sample soils collected from the experimental field which was subjected to a Zn biofortification workflow. A further study was also conducted as a means of investigating the interactions between the minerals present in the whole bread wheat flours of two varieties of *Triticum aestivum* L.

Results and Discussion

Samples	pH (H ₂ O)	Electrical Conductivity	Organic Matter	Moisture	K	Ca	Fe	Cu	Zn	Mg	P
		$\mu\text{S.cm}^{-1}$	%			mg.kg^{-1}					
A	7.70 ± 0.05 a	543 ± 44 a	6.92 ± 0.02 a	22.52 ± 0.17 a	0.652 ± 0.008 a	1.26 ± 0.11 a	38136 ± 278 a	43.45 ± 2.31 b	59.11 ± 1.20 a		
B	7.57 ± 0.10 a	412 ± 19 a	6.71 ± 0.07 a	22.62 ± 0.16 a	0.643 ± 0.029 ab	1.01 ± 0.05 a	39390 ± 257 a	44.56 ± 1.057 ab	60.36 ± 1.00 a	< 1500	< 200
C	7.73 ± 0.08 a	568 ± 47 a	6.95 ± 0.12 a	22.29 ± 0.38 a	0.572 ± 0.020 b	1.23 ± 0.12 a	38806 ± 1065 a	52.70 ± 3.591 a	59.96 ± 1.78 a		

Soil analyzes pH, electrical conductivity, organic matter and moisture contents and the values of calcium (Ca), iron (Fe) and Zn did not show significant differences among the different soil samples. The values of pH were slightly above 7 and the electrical conductivity varied between 412 and 568 $\mu\text{S.cm}^{-1}$. Potassium (K) showed lower values when compared to Ca (almost twice the values of K). The minerals quantification demonstrated higher levels of Fe, followed by Zn and copper (Cu) (Cu and Zn showed similar values). The minerals Mg and P presented values lower than 1500 and 200 mg.kg^{-1} , respectively.

Results and Discussion

There was a strong and positive correlation between the minerals: K and Fe for samples A and B; K and Zn for samples A and C; and Fe and Zn for the sample C. By contrast, there was a strong and negative correlation between the minerals: Cu and Zn for samples A and C; Fe and Ca for sample A; and Fe and Cu for the sample C.

a)						b)						c)					
A	Ca	K	Fe	Cu	Zn	B	Ca	K	Fe	Cu	Zn	C	Ca	K	Fe	Cu	Zn
Ca	1	-0,033	-0,45	0,217	-0,2	Ca	1	0,517	0,517	-0,1	0,517	Ca	1	0,517	-0,517	0,367	-0,467
K	-0,364	1	0,85	-0,583	0,717	K	0,397	1	0,917	-0,367	0,4	K	0,31	1	0,35	-0,45	0,4
Fe	-0,837	0,781	1	-0,667	0,633	Fe	0,668	0,932	1	-0,317	0,267	Fe	-0,51	0,652	1	-0,75	0,95
Cu	0,345	-0,671	-0,653	1	-0,8	Cu	-0,238	-0,349	-0,352	1	0,317	Cu	0,574	-0,518	-0,914	1	-0,817
Zn	-0,289	0,742	0,612	-0,913	1	Zn	0,332	0,093	0,235	0,499	1	Zn	-0,385	0,71	0,967	-0,839	1

Results and Discussion

No significant differences were observed between ZnSO₄ treatments for both varieties in the minerals magnesium (Mg), phosphor (P), Ca and Cu. Relatively to the minerals, P and K presented higher values in the flours contrasting with Mg and Ca. While assessing values of the microelements, Cu presented lower values than Zn and Fe. When comparing control samples (T0), it was found that Paiva variety presented higher mineral content for all the minerals. Zinc foliar spraying enhanced Zn content in Paiva and Roxo varieties in the flours. After applying Zn fertilizer in Paiva a decrease of the mineral content for Fe, K, Ca and Mg was observed. In contrast, considering Roxo, an increase of the mineral content for the minerals P, K, Mg, Fe, Zn and Cu regarding control samples, was observed.

Variety	Treatment	Mg	P	K	Ca	Fe	Cu	Zn	
		%				mg/kg			
Paiva (P)	T0	8.87 ± 0.03 a	115 ± 3.00 a	65.54 ± 5.32 b	1.87 ± 0.13 a	5.71 ± 0.10 a	0.241 ± 0.002 a	0.653 ± 0.013 b	
	T1	8.78 ± 0.30 a	138 ± 6.50 a	63.47 ± 1.39 a	1.04 ± 0.12 a	0.58 ± 0.17 ab	0.276 ± 0.013 a	0.739 ± 0.035 a	
	T2	8.32 ± 0.44 a	118 ± 11.6 a	63.89 ± 4.88 b	1.19 ± 0.30 a	1.44 ± 0.99 c	0.239 ± 0.006 a	1.143 ± 0.099 a	
Roxo (R)	T0	8.56 ± 0.13 a	98.1 ± 6.20 a	59.41 ± 5.97 b	1.31 ± 0.26 a	3.33 ± 0.06 bc	0.225 ± 0.021 a	0.638 ± 0.145 b	
	T1	9.04 ± 0.14 a	111 ± 3.20 a	69.73 ± 1.14 ab	1.58 ± 0.055 a	4.13 ± 0.10 ab	0.263 ± 0.003 a	1.177 ± 0.018 a	
	T2	8.92 ± 0.03 a	118 ± 2.80 a	60.92 ± 4.95 b	1.24 ± 0.10 a	3.970 ± 0.10 ab	0.257 ± 0.001 a	1.175 ± 0.011 a	

Results and Discussion

A strong and positive correlation was presented between the minerals: Zn and Ca, and K and Zn for samples PT0, RT0 and RT1; K and P for samples PT0, PT1 and RT1; K and Mg, Mg and Zn and Mg and Ca for samples PT0, PT1, RT0 and RT1; P and Mg for samples PT0, PT1, PT2 and RT1; K and Ca for samples PT0, PT1, PT2, RT0, RT1 and RT2; Zn and Cu, K and Cu, and Mg and Cu for samples PT1, RT0, and RT1; Zn and P and Cu and P for samples PT1 and RT1; Fe and Ca and Fe and K for samples PT1 and PT2; Fe and Mg and P and Fe for samples PT1 and RT2; Fe and Cu for sample PT1; Cu and Ca for samples RT0 and RT1; and P and Ca for samples RT1 and RT2.

Results and Discussion

Conversely, there was a strong and negative correlation between the minerals: Fe and Cu for sample PT0; Zn and P for samples PT2, RT0 and RT2; Cu and P for samples RT0 and RT2; K and Mg for sample PT2 and RT2; K and P and P and Ca for samples PT2 and RT0; P and Ca, and P and Mg for sample RT0; P and Fe, Fe and Mg, and Mg and Ca for sample PT2; and Zn and Fe, Cu and Ca, Zn and Ca, and K and Cu for the sample RT2.

a)

Paiva T0	Zn	Cu	Fe	Ca	K	P	Mg
Zn	1	-0,5	0,5	1	0,5	0,5	0,5
Cu	-0,514	1	-1	-0,5	0,5	0,5	0,5
Fe	0,645	-0,987	1	0,5	-0,5	-0,5	-0,5
Ca	0,982	-0,342	0,487	1	0,5	0,5	0,5
K	0,831	0,05	0,111	0,922	1	1	1
P	0,547	0,436	-0,287	0,697	0,92	1	1
Mg	0,823	0,064	0,096	0,916	1	0,926	1

b)

Paiva T1	Zn	Cu	Fe	Ca	K	P	Mg
Zn	1	1	0,5	0,5	0,5	1	0,5
Cu	0,968	1	0,5	0,5	0,5	1	0,5
Fe	0,521	0,719	1	1	1	0,5	1
Ca	0,203	0,443	0,941	1	1	0,5	1
K	0,668	0,834	0,983	0,864	1	0,5	1
P	0,927	0,992	0,803	0,556	0,899	1	0,5
Mg	0,895	0,978	0,847	0,619	0,93	0,997	1

c)

Paiva T2	Zn	Cu	Fe	Ca	K	P	Mg
Zn	1	0,5	0,5	0,5	0,5	-0,5	-0,5
Cu	0,343	1	-0,5	-0,5	-0,5	0,5	0,5
Fe	0,486	-0,654	1	1	1	-1	-1
Ca	0,635	-0,508	0,984	1	1	-1	-1
K	0,695	-0,437	0,966	0,997	1	-1	-1
P	-0,721	0,403	-0,956	-0,993	-0,999	1	1
Mg	-0,697	0,434	-0,965	-0,997	-1	0,999	1

d)

Roxo T0	Zn	Cu	Fe	Ca	K	P	Mg
Zn	1	0,5	0,5	0,5	1	-1	0,5
Cu	0,971	1	-0,5	1	0,5	-0,5	1
Fe	0,452	0,225	1	-0,5	0,5	-0,5	-0,5
Ca	0,784	0,91	-0,199	1	0,5	-0,5	1
K	1	0,973	0,443	0,79	1	-1	0,5
P	-0,999	-0,96	-0,488	-0,758	-0,999	1	-0,5
Mg	0,8	0,92	-0,174	1	0,806	-0,775	1

e)

Roxo T1	Zn	Cu	Fe	Ca	K	P	Mg
Zn	1	0,5	-0,5	0,5	0,5	1	0,5
Cu	0,873	1	0,5	1	1	0,5	1
Fe	-0,462	0,03	1	0,5	0,5	-0,5	0,5
Ca	0,942	0,986	-0,138	1	1	0,5	1
K	0,935	0,989	-0,119	1	1	0,5	1
P	0,995	0,821	-0,547	0,905	0,896	1	0,5
Mg	0,817	0,994	0,135	0,963	0,968	0,756	1

f)

Roxo T2	Zn	Cu	Fe	Ca	K	P	Mg
Zn	1	0,5	-1	-0,5	0,5	-0,5	-0,5
Cu	0,699	1	-0,5	-1	-0,5	-1	0,5
Fe	-0,835	-0,192	1	0,5	-0,5	0,5	0,5
Ca	-0,761	-0,996	0,279	1	0,5	1	-0,5
K	-0,29	-0,887	-0,283	0,842	1	0,5	-1
P	-0,994	-0,774	0,77	0,828	0,394	1	-0,5
Mg	-0,454	0,319	0,869	-0,232	-0,72	0,353	1

Results and Discussion

Since all the soil analyzes, except for the minerals K and Cu, did not present significant differences, we can presume that the experimental field is homogeneous. The mineral K plays an important roll in plants metabolism, for example in the regulation of the opening and closing of stomates, activation of some enzymes and balancing the use of N. This mineral has an antagonist effect on absorb Ca, Mg and P (Fageria, 2001). According to (Pagani *et al.* 2013; El-Ramady *et al.* 2014), K presents a synergetic interaction with Fe. Besides, K moves in soils by diffusion and is mobile in the plant (Pagani *et al.* 2013; El-Ramady *et al.* 2014). The minerals Mg, Ca, Fe, Cu and Zn move in the soil by mass flow, whereas the minerals P and Fe, move in soil by diffusion (El-Ramady *et al.* 2014). Regarding the mineral's mobility in plant, the minerals are P, mobile and the minerals are Mg (relatively immobile), Ca, Fe, Cu and Zn immobile (Pagani *et al.* 2013; El-Ramady *et al.* 2014).

Results and Discussion

One of the functions of Ca is to be a cofactor of various enzymes of ATP. The mineral has an antagonist interaction with Cu, Fe and Zn, but also interacts with Cu and Zn in a synergetic way (Pessoa *et al.* 2021; El-Ramady *et al.* 2014). Magnesium is a component of chlorophyll and functions as an enzyme activator in plants. This mineral interacts with Cu, Fe and Zn in an antagonist way, however, presents a synergetic interaction with Zn (Pagani *et al.* 2013; El-Ramady *et al.* 2014). The mineral P is an important constituent of nucleic acids, proteins, metabolic substrates and coen-zymes. This mineral has both antagonist and synergetic interactions with Cu, Fe and Zn (Pagani *et al.* 2013; El-Ramady *et al.* 2014). Iron plays an important role in plants in chlorophyll synthesis and also in enzyme electron transfer. Iron presents an antagonist interaction with Ca, Mg and P, whereas it only interacts with P in a synergetic way (El-Ramady *et al.* 2014). The mineral Cu is part of a diversity of enzymes and works as a catalyst for respiration. Copper interacts with Ca and P in both antagonist and synergetic ways, and only presents an antagonist interaction with Mg (El-Ramady *et al.* 2014). Zinc has a myriad of functions in plants like being part of enzymes from regulation and has both antagonist and synergetic interactions with Ca, Mg and P (Pagani *et al.* 2013; El-Ramady *et al.* 2014). Taking everything into account, most of the results obtained were not in line with what was said by the authors (El-Ramady *et al.* 2014) has most of the minerals presented strong positive correlations in the whole wheat flours, so the majority of the minerals showed a synergetic interaction.

Conclusions

In general, there were no significant differences between the soil samples in the various parameters analyzed. Considering macroelements, Ca presented higher values in the soils. Conversely, Fe was the dominant microelement. In the soil samples, it was observed that only the minerals K, Fe and Zn were strongly and positively correlated, however, the minerals Fe, Cu, Ca (only with Fe) and Zn (only with Cu) had a strong and negative correlation. When compared to Roxo, Paiva variety presented higher mineral content for all the minerals in the flours of the control samples (P0 and R0). When applying Zn fertilizer in Paiva a decrease of the mineral content for Fe, K, Ca and Mg, was observed. Nevertheless, an increase of the mineral content for the minerals P, K, Mg, Fe, Zn and Cu regarding control samples was observed in Roxo. Zn foliar spraying enhanced Zn content in both varieties. Thus, wheat flour biofortified in zinc can be a product to help overcome malnutrition. Regarding whole bread wheat flours, it was observed that, in general, the minerals were strongly and positively correlated, although in some cases the minerals also had a strong and negative correlation.

Acknowledgments

The authors thanks to Sociedade Agrícola Saramago de Brito, Instituto Politécnico de Beja and Associação de Agricultores do Baixo Alentejo for technical assistance and for facilities regarding *Triticum aestivum* L. experimental field. Furthermore, we also thank to the research center (GeoBioTec) UIDB/04035/2020 for lab facilities.

