

Proceeding Paper

Zn Nutrition of *Vitis Vinifera* White Grapes: Characterization of Antagonistic and Synergistic Interactions by μ EDXRF Tissue Analyses [†]

Diana Daccak ^{1,2,*}, Cláudia Campos Pessoa ^{1,2}, Inês Carmo Luís ^{1,2}, Ana Coelho Marques ^{1,2}, Ana Rita F. Coelho ^{1,2}, José C. Ramalho ^{2,3}, José N. Semedo ^{2,4}, Mauro Guerra ⁵, Roberta G. Leitão ⁵, Paula Scotti Campos ^{2,4}, Isabel P. Pais ^{2,4}, Maria Manuela Silva ^{2,6}, Fernando H. Reboredo ^{1,2}, Manuela Simões ^{1,2}, Maria Fernanda Pessoa ^{1,2}, Paulo Legoinha ^{1,2} and Fernando C. Lidon ^{1,2}

- ¹ Earth Sciences Department, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa. Campus da Caparica, 2829-516 Caparica, Portugal
 - ² GeoBioTec Research Center, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa. Campus da Caparica, 2829-516 Caparica, Portugal
 - ³ PlantStress & Biodiversity Lab, Centro de Estudos Florestais (CEF), Instituto Superior Agronomia (ISA), Universidade de Lisboa (ULisboa), Quinta do Marquês, Av. República, 2784-505 Oeiras, and Tapada da Ajuda, 1349-017 Lisboa, Portugal
 - ⁴ Instituto Nacional de Investigação Agrária e Veterinária, I.P. (INIAV), Quinta do Marquês, Av. República, 2780-157 Oeiras, Portugal
 - ⁵ LIBPhys-UNL, Physics Department, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal
 - ⁶ ESEAG-COFAC, Avenida do Campo Grande 376, 1749-024 Lisboa, Portugal
- * Correspondence: d.daccak@campus.fct.unl.pt

[†] Presented at the 1st International Online Conference on Agriculture—Advances in Agricultural Science and Technology (IOCAG2022), 10–25 February 2022; Available online: <https://iocag2022.sciforum.net/>.

Citation: Daccak, D.; Pessoa, C.C.; Luís, I.C.; Marques, A.C.; Coelho, A.R.F.; Ramalho, J.C.; Semedo, J.N.; Guerra, M.; Leitão, R.G.; Campos, P.S.; et al. Zn Nutrition of *Vitis Vinifera* White Grapes: Characterization of Antagonistic and Synergistic Interactions by μ EDXRF Tissue Analyses. *Chem. Proc.* **2022**, *4*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s):

Published: 10 February 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Nutritional status performs an essential role in agriculture, affecting productivity and keeping crops functioning properly. Despite being required in smaller amounts, micronutrients are also needed for adequate growth, namely Zn (zinc) with regulatory, catalytic and structural functions. Fertilization with Zn is used to ameliorate the deficits of this micronutrient in more susceptible crops such as grapes. Yet, the management of this application must consider the antagonistic and synergistic interactions among nutrients, as it affects their uptake and translocation rates. Therefore, a workflow with three ZnO foliar applications (30% and 60%, 450 and 900 g ha⁻¹ respectively) in the variety *Vitis Vinifera* cv. Fernão Pires, was implemented in a field located in Palmela, Portugal. The concentration of Zn in the tissues was therefore evaluated by microenergy X-ray dispersion fluorescence (μ EDXRF), showing an increase of 1.82 and 2.54 times in the seed and flesh of grapes fertilized with a concentration of 60%, compared to control grapes, respectively. Using the same method, a synergistic relationship was observed for macronutrients such as Ca and K, and micronutrients such as Fe, P and Mn. In addition, a complementary analysis of grapes' density was carried out to verify changes in quality, in which no negative impact was observed due to Zn application. This study allows us to verify that the concentration of the applied Zn fertilizer brings benefits in the amount of nutrients that are important for development and crops quality.

Keywords: antagonism; grapes; synergism; *Vitis vinifera*; Zn fertilizer

1. Introduction

Micronutrient's deficiencies or hidden hunger is affecting more than 2 billion people worldwide, leading to the development of strategies to mitigate this problem [1]. In this context, agronomic biofortification has become a tool to improve the mineral content in

the edible parts of food crops faster, more reliably, more sustainably and economically [2]. Through foliar fertilization, it becomes possible to mitigate crop deficiencies, selecting target minerals [3], without limiting the fixation and retention of nutrients in the soil [4].

Fertilizing fruit trees, namely the vineyard, is considered an important tool to meet nutritional needs and optimize the yield and quality of the grapes [5,6]. Zinc (Zn) deficits are common worldwide, affecting the vine [7] with visible symptoms such as chlorosis, necrotic spots and small leaves [6]. In this regard, Zn is an important micronutrient for plant growth, with catalytic, structural and regulatory functions [8], as a co-factor in the auxin metabolism, enzymatic activation, chlorophyll and nucleotide synthesis, and genes expression and regulation [9]. According to research in grapefruit, Zn agronomic biofortification has been shown to be efficient and, in addition enhancing growth and development [10].

Likewise, phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), sulphur (S), iron (Fe) and manganese (Mn) are also essential, affecting crops productivity [11]. Yet, through application of one nutrient, absorption and utilization of other nutrients can be affected, in a positive or negative way, in a synergistic and antagonistic relationship respectively [12]. These interactions become visible through growth and changes in nutrients amounts [13] affecting several metabolic processes [14].

2. Materials and Methods

2.1. Experimental Field

The experimental work was carried out in a *Vitis vinifera* cv. Fernão Pires, at a field located in Palmela, Portugal (N 38° 35'41.467" O 8° 50'44.535" W). Grapes were pulverized three times, during the reproductive cycle, with ZnO at concentrations of 30% and 60% (corresponding to 450 and 900 g ha⁻¹). Harvest was performed at 17th of September.

2.2. Quantification of Nutrients in Grapes and Accumulation Level in Grape Tissues

The nutrient content in the grapes was determined in the tissues (flesh and seeds) at harvest, with a Micro-energy X-ray Dispersion Fluorescence (μ -EDXRF) (M4 Tornado™, Bruker, Berlin, Germany) system, according to [15], with an adaptation of the assessment of nutrients such as Zn, Fe, Mn, P, Ca, K and Mg in grapes.

2.3. Physicochemical Parameter of Grapes

Determination of density (kg m⁻³) was performed considering three randomized grapes per treatment.

2.4. Statistical Analysis

Data were statistically treated applying one-way ANOVA ($p \leq 0.05$) to determined differences, and before a Tukey's test for mean comparison (95% confidence level) was performed.

3. Results

3.1. Quantification of Nutrients in Grapes Flesh and Seed

Zn seed concentration reached an increase of 1.51 and 1.82 times more than control, with ZnO 30% and 60% respectively (both significantly different). Regarding Mn and P the same trend is observed with ZnO 60% treatment demonstrating the highest value, followed by ZnO 30%, being both significantly different compared to control. For Mn and P, the range of values varied between 8.22–35.9 and 0–3403 ppm respectively. Iron amount didn't showed significative differences between all samples presenting values between 91.6–106 ppm.

The flesh of the grapes showed a significant increase in Zn concentration of 2.32 and 2.54 times more with ZnO 30% and 60% respectively, compared to the control. Also, for

the nutrient Fe and P, significant differences were observed with both treatments (i.e., ZnO 30% and 60%) compared to the control, but contrary to what was observed in the seeds (i.e., Zn, Mn and P), the lower concentration of Zn fertilizer showed the highest value for these nutrients. On the other hand, for Mn, the same tendency of the seeds is observed, where the concentration of ZnO 60% presented a higher amount, with the two treatments being significantly different in relation to the control.

Seed Ca levels were higher with the 30% treatment (2.98%), followed by the 60% treatment (1.88%), both being significantly different compared to the control. In seed, the nutrient K, with the application of Zn fertilizer did not result in significant differences in relation to the control, showing values between 5.16–6.40%. The concentration of Mg in seeds was significantly different and higher for control compared to grapes treated with ZnO.

In relation to the grape flesh, the nutrients Ca and K submitted to the treatment increased significantly in relation to the control, with the highest values for the ZnO 30% treatment (1.89 and 15.4% for Ca and K respectively). Likewise, for the relationship observed in the seeds, for the pulp, Mg significantly decreased with the application of the two treatments.

3.2. Density

Fertilized grapes showed a higher value, although not demonstrating significant differences regarding to control. Values for density in Fernão Pires grapes showed a range between 1150–2185 kg m⁻³.

4. Discussion

Nowadays, to meet the food necessities according to population growth, different techniques are being explored to increase agricultural crops yields [16]. Accordingly, mineral nutrition is among the most used agricultural techniques to obtain higher yields and quality [17], namely Zn fertilization through soil and / or foliar applications [18]. Most studies were carried out with soil fertilization, implying the influence of physicochemical and biological soil properties, such as pH, water, structure, cation exchange capacity, redox conditions, and activity of microorganisms [19]. As mentioned in the literature, Zn fertilization in soil has an antagonistic response with most nutrients such as Fe, Ca, Mg, K, Mn, and P, related to similar absorption mechanisms that lead to competition (i.e., except for P which is a specific interaction) [20].

In this context, the present study carried out foliar applications with ZnO at different concentrations (i.e., 450 and 900 g ha⁻¹) in vines, proving to be efficient in increasing the Zn content in grapes, reaching 2.54 times more in the flesh and 1.82 in seed than without fertilization (Table 1). Furthermore, the other nutrients accessed, showed mainly an interaction of synergism in the case of micronutrients and macronutrients such as P, Mn, Ca, Fe and K (i.e., except Mg) (Tables 1 and 2). The results obtained, showed that increase of Zn concentration in the leaves, enhanced the uptake of other nutrients, being in agreement with [21], suggesting that micronutrients as Zn stimulate the plant metabolism resulting in an intensified uptake of nutrients through the roots. Also, Zn fertilization is related to positive effects on photosynthesis and chlorophyll synthesis, which also facilitates the absorption and accumulation of nutrients in the leaves [22]. Likewise, a study with foliar application of Zn in a fruit tree showed the same tendency of increasing the amount of Zn in the leaves to promote nutrients such as P and Ca and demonstrated to be antagonistic with Mn and Fe [22]. In the case of Fe, our results showed a different response, with Zn fertilization at concentrations applied (i.e., 450 and 900 g ha⁻¹), showing a relation of synergism in grapes flesh (Table 1). As for Potassium, it presented the same response as Fe, with a synergism relation, only in the flesh, with the 30% (i.e., 450 g ha⁻¹) treatment being the one with the greatest increase in the amount of this element (Table 2). A greater increase was noticed in most results with the lowest ZnO 30% treatment, namely for the

macronutrients Ca and K and for the micronutrients Fe and P (i.e., in the flesh) (Table 1 and 2). As for the antagonistic interactions, in this study, they only occurred in Mg, where the greatest amount of treatment with Zn significantly reduced the Mg concentration. Magnesium has a competitive relationship with other nutrients, which according to [23], the nutrient Ca, due to a greater affinity with the binding sites of the root plasma membrane, decreases the amount of Mg [23]. In fact, in the present study this condition is present with an increase in Ca content being observed with Zn fertilization.

Grape density is also important as it depends on important biophysical parameters (i.e., grape diameter, length, volume, and mass) and biochemical composition [24]. According to literature, higher densities (i.e., $\geq 1088 \text{ kg m}^{-3}$) demonstrated benefits, being more advantageous for health and sensorially [25]. Which, this experimental work with Fernão Pires grapes presented greater values ranging between 1150–2185 kg m^{-3} (Figure 1), where fertilization didn't negatively affect the quality of grapes, being this important for consumer acceptance. In fact, fertilized grapes showed higher values than control, although not significant.

Table 1. Average content ($n = 3$) of Zn, Fe, Mn and P (ppm) in grape flesh and seeds (after dehydration) in Fernão Pires grapes at harvest, and the respective degree of uncertainty. Letters a, b, c indicate significant differences between treatments ($p < 0.05$). Treatments ZnO with concentrations of 0%, 30%, 60%. (i.e., 0, 450, and 900 g ha^{-1}).

Micronutrient (Seed)				
Sample	Zn	Fe	Mn	P
Control	24.6 ± 1.23 c	91.6 ± 4.58 a	8.22 ± 0.41 c	0.00 ± 0.00 c
ZnO 30%	37.1 ± 1.85 b	106 ± 5.31 a	14.8 ± 0.74 b	1148 ± 57.4 b
ZnO 60%	44.7 ± 2.24 a	105 ± 5.24 a	35.9 ± 1.79 a	3403 ± 170 a
Micronutrients (Flesh)				
Control	23.4 ± 1.17 b	91.2 ± 4.56 c	8.66 ± 0.43 c	0.00 ± 0.00 c
ZnO 30%	54.4 ± 2.72 a	278 ± 13.9 a	27.4 ± 1.37 b	2662 ± 133 a
ZnO 60%	59.4 ± 2.97 a	228 ± 11.4 b	51.8 ± 2.59 a	2203 ± 110 b

Table 2. Average content ($n = 3$) of Ca, K and Mg (%) in grape flesh and seeds (after dehydration) in Fernão Pires grapes at harvest, and the respective degree of uncertainty. Letters a, b, c indicate significant differences between treatments ($p < 0.05$). Treatments ZnO with concentrations of 0%, 30%, 60%. (i.e., 0, 450, and 900 g ha^{-1}).

Macronutrient (Seed)			
Sample	Ca	K	Mg
Control	0.82 ± 0.04 c	6.11 ± 0.31 ab	12.4 ± 0.62 a
ZnO 30%	2.98 ± 0.15 a	6.40 ± 0.32 a	0.39 ± 0.02 c
ZnO 60%	1.88 ± 0.09 b	5.16 ± 0.26 b	2.78 ± 0.14 b
Macronutrients (Flesh)			
Control	0.35 ± 0.02 c	5.44 ± 0.27 c	10.0 ± 0.50 a
ZnO 30%	1.89 ± 0.09 a	15.4 ± 0.77 a	5.84 ± 0.29 b
ZnO 60%	1.24 ± 0.06 b	10.9 ± 0.55 b	5.51 ± 0.28 b




Figure 1. Average content ($n = 3$) of density (kg m^{-3}) in Fernão Pires grapes at harvest, and the respective degree of uncertainty. Letter a indicate the absence of significant differences between treatments ($p < 0.05$). Treatments ZnO with concentrations of 0%, 30%, 60%. (i.e., 0, 450, and 900 g ha^{-1}).

5. Conclusions

Zinc foliar fertilization with ZnO is efficient, increasing the amount of this micronutrient in the flesh and seeds of the grape cv. Fernão Pires, and presenting a synergistic relationship with P, Fe, Mn, Ca and K. Thus, this technique is an important tool to improve the nutritional status and quality of the grape. An antagonistic relationship with the macronutrient Mg was also observed.

Additionally, fertilization did not affect density, with Fernão Pires grapes presenting values greater than 1088 kg m^{-3} being important for health and sensory.

6. Patents

Author Contributions: Conceptualization, M.G. and F.C.L.; methodology, M.G. and R.G.L.; formal analysis, D.D., C.C.P., I.C.L., A.C.M. and A.R.F.C.; resources, J.C.R., J.N.S., M.G., R.G.L., P.S.C., I.P.P., M.M.S., F.H.R., M.S., M.F.P. and P.L.; writing—original draft preparation, D.D.; writing—review and editing, D.D. and F.C.L.; supervision, F.C.L.; project administration, F.C.L.; funding acquisition, F.C.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by PDR2020, grant number 101-030727.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors thanks to Engenier Luís Silva (Adega Cooperativa de Palmela-Casa Agrícola Nunes Oliveira da Silva Lda) for technical assistance to project PDR2020-101-030727—for the financial support. We also thanks to the Research centres (GeoBioTec) UIDB/04035/2020. This work was further supported in part by the research center Grant N^o. UID/FIS/04559/2013 to LIBPhys-UNL, from the FCT/MCTES/PIDDAC and by the project PDR2020-101-030727.

Conflicts of Interest: The authors declare no conflict of interest. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

References

- Gödecke, T.; Stein, A.J.; Qaim, M. The global burden of chronic and hidden hunger: Trends and determinants. *Glob. Food Secur.* **2018**, *17*, 21–29. <https://doi.org/10.1016/j.gfs.2018.03.004>.
- Coelho, A.R.F.; Lidon, F.C.; Pessoa, C.C.; Marques, A.C.; Luís, I.C.; Caleiro, J.; Simões, M.; Kullberg, J.; Legoinha, P.; Brito, M.; et al. Can Foliar Pulverization with CaCl_2 and $\text{Ca}(\text{NO}_3)_2$ Trigger Ca Enrichment in *Solanum tuberosum* L. Tubers?. *Plants* **2021**, *10*, 245. <https://doi.org/10.3390/plants10020245>.
- Pessoa, C.C.; Coelho, A.R.F.; Luís, I.C.; Marques, A.C.; Daccak, D.; Simões, M.; Reboredo, F.; Silva, M.M.; Pessoa, M.F.; Galhano, C.; et al. A Technological workflow for Ca enrichment in rocha pears: Implication in quality. In Proceedings of the 1st International Conference on Water Energy Food and Sustainability (ICoWEFS 2021), Leiria, Portugal, 10–12 May 2021; da Costa Sanches Galvão J.R., Galvão, J., Brito, P., dos Santos, F., Craveiro, F., Almeida, H., Vasco, J., Neves, L., Gomes, R., Mourato, S.; et al., Eds.; Springer Nature: Cham, Switzerland, 2021. https://doi.org/10.1007/978-3-030-75315-3_21.

4. Sathishkumar, A.; Sakthivel, N.; Subramanian, E.; Rajesh, P. Productivity of field crops as influenced by foliar spray of nutrients: A review. *Agric. Rev.* **2021**, *41*, 146–152. <https://doi.org/10.18805/ag.R-1991>.
5. Brunetto, G.; Melo, G.W.B.D.; Toselli, M.; Quartieri, M.; Tagliavini, M. The role of mineral nutrition on yields and fruit quality in grapevine, pear and apple. *Rev. Bras. Frutic.* **2015**, *37*, 1089–1104. <https://doi.org/10.1590/0100-2945-103/15>.
6. Song, C.Z.; Liu, M.Y.; Meng, J.F.; Chi, M.; Xi, Z.M.; Zhang, Z.W. Promoting effect of foliage sprayed zinc sulfate on accumulation of sugar and phenolics in berries of *Vitis vinifera* cv. Merlot growing on zinc deficient soil. *Molecules* **2015**, *20*, 2536–2554. <https://doi.org/10.3390/molecules20022536>.
7. Sabir, A.; Sari, G. Zinc pulverization alleviates the adverse effect of water deficit on plant growth, yield and nutrient acquisition in grapevines (*Vitis vinifera* L.). *Sci. Hort.* **2019**, *244*, 61–67. <https://doi.org/10.1016/j.scienta.2018.09.035>.
8. Jurowski, K.; Szewczyk, B.; Nowak, G.; Piekoszewski, W. Biological consequences of zinc deficiency in the pathomechanisms of selected diseases. *J. Biol. Inorg. Chem.* **2014**, *19*, 1069–1079. <https://doi.org/10.1007/s00775-014-1139-0>.
9. Luís, I.C.; Lidon, F.C.; Pessoa, C.C.; Marques, A.C.; Coelho, A.R.F.; Simões, M.; Patanita, M.; Dôres, J.; Ramalho, J.C.; Silva, M.M.; et al. Zinc enrichment in two contrasting genotypes of *Triticum aestivum* L. grains: Interactions between edaphic conditions and foliar fertilizers. *Plants* **2021**, *10*, 204. <https://doi.org/10.3390/plants10020204>.
10. Fu, X.Z.; Xing, F.; Cao, L.; Chun, C.P.; Ling, L.L.; Jiang, C.L.; Peng, L.Z. Effects of foliar application of various zinc fertilizers with organosilicone on correcting citrus zinc deficiency. *HortScience* **2016**, *51*, 422–426. <https://doi.org/10.21273/HORTSCI.51.4.422>.
11. Toor, M.D.; Adnan, M.; Rehman, F.; Tahir, R.; Saeed, M.S.; Khan, A.U.; Pareek, V. Nutrients and Their Importance in Agriculture Crop Production; A Review. *Ind. J. Pure Appl. Biosci.* **2021**, *9*, 1–6. <https://doi.org/10.18782/2582-2845.8527>.
12. Palani, V.; Raju, I. (2019). Synergistic and Antagonistic Interactions of Calcium with Other Nutrients in Soil and Plants. *SSRN* **2019**. <https://doi.org/10.2139/ssrn.3503225>.
13. Fageria, V.D. Nutrient interactions in crop plants. *J Plant Nutr.* **2001**, *24*, 1269–1290. <https://doi.org/10.1081/PLN-100106981>.
14. Kumar, S.; Kumar, S.; Mohapatra, T. (2021). Interaction Between Macro-and Micro-Nutrients in Plants. *Front. Plant Sci.* **2021**, *12*, 753. <https://doi.org/10.3389/fpls.2021.665583>.
15. Marques, A.C.; Lidon, F.C.; Coelho, A.R.F.; Pessoa, C.C.; Luís, I.C.; Campos, P.S.; Simões, M.; Almeida, A.S.; Pessoa, M.F.; Galhano, C.; et al. Effect of rice grain (*Oryza sativa* L.) enrichment with selenium on foliar leaf gas exchanges and accumulation of nutrients. *Plants* **2021**, *10*, 288. <https://doi.org/10.3390/plants10020288>.
16. Mousavi, S.R.; Galavi, M.; Rezaei, M. The interaction of zinc with other elements in plants: A review. *Intl J Agri Crop Sci.* **2012**, *4*, 1881–1884.
17. de Souza Júnior, J.O.; da Silveira, F.G.F.; dos Santos, R.O.; Neves, J.C.L. Zinc fertilizers and additives for foliar fertilization of cocoa seedlings. *J. Agric. Sci.* **2019**, *11*, 471–478. <https://doi.org/10.5539/jas.v11n1p471>.
18. Wang, S.; Li, M.; Liu, K.; Tian, X.; Li, S.; Chen, Y.; Jia, Z. Effects of Zn, macronutrients, and their interactions through foliar applications on winter wheat grain nutritional quality. *PLoS ONE* **2017**, *12*, e0181276. <https://doi.org/10.1371/journal.pone.0181276>.
19. Frossard, E.; Bucher, M.; Mächler, F.; Mozafar, A.; Hurrell, R. Potential for increasing the content and bioavailability of Fe, Zn and Ca in plants for human nutrition. *J. Sci. Food Agric.* **2000**, *80*, 861–879. [https://doi.org/10.1002/\(SICI\)1097-0010\(20000515\)80:7<861::AID-JSFA601>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1097-0010(20000515)80:7<861::AID-JSFA601>3.0.CO;2-P).
20. Rietra, R.P.; Heinen, M.; Dimkpa, C.O.; Bindraban, P.S. Effects of nutrient antagonism and synergism on yield and fertilizer use efficiency. *Commun. Soil Sci. Plant Anal.* **2017**, *48*, 1895–1920. <https://doi.org/10.1080/00103624.2017.1407429>.
21. Stepien, A.; Wojtkowiak, K. Effect of foliar application of Cu, Zn, and Mn on yield and quality indicators of winter wheat grain. *Chilean J. Agric. Res.* **2016**, *76*, 220–227. <https://doi.org/10.4067/S0718-58392016000200012>.
22. Xie, R.; Zhao, J.; Lu, L.; Brown, P.; Guo, J.; Tian, S. Penetration of foliar-applied Zn and its impact on apple plant nutrition status: In vivo evaluation by synchrotron-based X-ray fluorescence microscopy. *Hortic. Res.* **2020**, *7*, 1–12. <https://doi.org/10.1038/s41438-020-00369-y>.
23. Njira, K.O.; Nabwami, J. A review of effects of nutrient elements on crop quality. *African J. Food Agric. Nutr. Dev.* **2015**, *15*, 9777–9793.
24. Letchov, G.; Roychev, V. Growth kinetics of grape berry density (*Vitis vinifera* L. 'Black Corinth'). *Vitis* **2017**, *56*, 155–159. <https://doi.org/10.5073/vitis.2017.56.155-159>.
25. Rolle, L.; Torchio, F.; Giacosa, S.; Segade, S.R. Berry density and size as factors related to the physicochemical characteristics of Muscat Hamburg table grapes (*Vitis vinifera* L.). *Food Chem.* **2015**, *173*, 105–113. <https://doi.org/10.1016/j.foodchem.2014.10.033>.