

Proceedings

Identification of Pathogens in Seedlings of Indian Sandalwood and Screening of Fungal Endophytes Against the Plant Pathogenic Fungi [†]

Swetha Purushotham ^{1,*}, Sundararaj Ramachandran ², Kandan Aravindaram ³, Ananda Kulal ⁴, and Kirana Mugaranja P ⁵¹ PhD research scholar, Institute of wood science and technology² Scientist-G and Head, Forest Protection division, Institute of wood science and technology; rsundari-wst@gmail.com³ Principle scientist, Division of Germplasm Conservation and Utilization, ICAR-National Bureau of Agricultural Insect Resources; genekannz@gmail.com⁴ Associate Professor, Biological Sciences, Poornaprajna Institute of Scientific Research; ananda@poornaprajna.org⁵ PhD scholar, Biological Sciences, Poornaprajna Institute of Scientific Research; ananda@poornaprajna.org

* Correspondence: mpkirana@poornaprajna.org

[†] Presented at the 3rd International Electronic Conference on Forests — Exploring New Discoveries and New Directions in Forests, 15 to 31 October 2022. Available online: <https://iecf2022.sciforum.net>

Citation: Burzyńska, I.; Burzyńska, I.; Sztabkowski, K. TITLE. *Environ. Sci. Proc.*, **2022**, *4*, x. <https://doi.org/10.3390/xxxxx>.

Academic Editor: Rodolfo Picchio

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Indian sandalwood (*Santalum album* Linn.), an evergreen tree, indigenous to the Indian peninsula is known for its fragrant heartwood worldwide. Sandalwood plantations are gaining importance throughout the Indian subcontinent demanding large-scale production and the establishment of nurseries with Quality Planting Material (QPM). However, sandalwood seedlings succumb to devastating diseases at nurseries leading to high mortality of the planting stock. Therefore, there is a dire need for the effective management of these diseases. In our study, we isolated and identified phytopathogenic fungi such as *Fusarium solani* causing wilt disease with seedling mortality of 25 % and *Colletotrichum siamense* causing anthracnose disease with a disease incidence of 75 %. We identified and characterized a total of 90 fungal endophytic isolates from leaf, stem, and root tissues of disease escaped or apparently healthy seedlings of sandalwood. Total fungal endophytes isolated from the disease escaped sandalwood seedlings comprised 33.3 % *Colletotrichum siamense*, 26.6% *Diaporthe melonis*, 13.3% *Aspergillus sclerotiorum*, 13.3% *Fusarium oxysporum*, 13.3 *Paraphoma radicina*, 6.6% *Alternaria alternata* and 6.6% *Pestalotiopsis microspora*. Molecular identification using the nuclear ribosomal DNA internal transcribed spacer (ITS) sequences were performed and those species which could not be resolved with ITS sequence were subjected to multi-locus gene (beta-tubulin (TUB2), glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene, chitin synthase 1 gene (CHS-1), actin gene (ACT) and glutamine synthetase (GS) genes) analysis and the sequences were deposited to GenBank. Dual culture test assay revealed that the fungal endophytes *Aspergillus sclerotiorum* and *Diaporthe melonis* showed the highest percent inhibition of 63.08% and 61.54%, respectively against *Fusarium solani* and *Diaporthe melonis* and *Fusarium oxysporum* showed highest percent inhibition of 55.38 % and 67.69 % in case of the pathogen *Colletotrichum siamense*. This study will be useful for the management of seed, soil and airborne pathogens of Indian sandalwood.

Keywords: Indian sandalwood; endophytes; pathogen; management