

A COMPREHENSIVE REVIEW ON A MARVEL FRUIT OF ANNONA MURICATA

Hetakshi V. Patil ¹ and Mansi A. Dhankani ^{2,*} and Amitkumar R. Dhankani ³

¹ P.S.G.V.P.Mandal's College of Pharmacy, Dist-Nandurbar, Shahada 425409, Maharashtra, India; patilhetakshi18@gmail.com

² Assistant Professor, Department of Pharmaceutics, P.S.G.V.P.Mandal's College of Pharmacy, Dist-Nandurbar, Shahada 425409, Maharashtra, India; laxmipremchandani3@gmail.com

³ Assistant Professor, Department of Quality Assurance, P.S.G.V.P.Mandal's College of Pharmacy, Dist-Nandurbar, Shahada 425409, Maharashtra, India; dhankaniamit@gmail.com

Abstract: The purpose of this review is to delve into the essence of *Annona muricata* Linn. *A. muricata*, also known as soursop, guanabana, and graviola, is a member of the Annonaceae family and has a long history of traditional uses. This is an evergreen plant that grows in tropical and subtropical regions of the world, primarily in Africa, South America, and Southeast Asia. The *A. muricata* plant's miraculous nature is a boon to mankind, and it has been widely used in folk medicine. *A. muricata* preparations on the market include candies, syrups, beverages, ice creams, and shakes. Several studies have concluded that the plant contains over 212 chemical constituents such as acetogenins, alkaloids, and phenols. The plant has antibacterial, antiviral, antifungal, antitumor, anthelmintic, analgesic, antiarthritic, hypotensive, anti-inflammatory, immune enhancing effects, and anti-diabetic activity. Although some toxicities have been reported, the extract of *A. muricata* has been found to be effective and safe. This review attempts to bring together the majority of the available information on *A. muricata* phytochemistry, traditional uses, biological activities, and toxicity.

Keywords: *Annona muricata*; Annonaceae; Annonaceous acetogenins; Cytotoxicity; Neurotoxicity

Citation: Patil, H.V.; Dhankani, M.A. A COMPREHENSIVE REVIEW ON A MARVEL FRUIT OF *ANNONA MURICATA*. 2023, 3, x. <https://doi.org/10.3390/xxxxx> Published: 21 April 2023

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Natural products, particularly those derived from plants, have been used to help mankind maintain its health since the dawn of medicine. Over the last century, plant phytochemicals have been an important pipeline for pharmaceutical discovery. The value of active ingredients. The use of plants in agriculture and medicine has piqued scientists' interest. ^[1]

According to the World Health Organization (WHO), more than 80% of the world's population relies on traditional medicines to meet their primary health care needs⁶. The primary characteristics were medicinal plant chemical substances that exerted a physiologic action on the human body. The most important plant bioactive compounds were thought to be alkaloids, flavonoids, tannins, and phenolic compounds. The plant chemical Phytochemical Ethno-pharmacologicaffective approach to discovering new anti-infective agents from higher plants. ^[2]

Intensive chemical studies of this species' leaves and seeds have resulted in the isolation of a large number of acetogenins. Some of the isolated compounds exhibit interesting biological or pharmacological activities, such as antitumoral, cytotoxic, antiparasitic, and pesticidal properties. Because of their anti-parasitic and pesticidal properties, the roots of these species are used in traditional medicine.^[3]

The extract of *A. muricata*'s fruits, seeds, bark, roots, and pericarp contain over 212 phytochemical varieties that are used to treat a variety of ailments. ⁵ According to his-

torical records, *A. muricata* was used as an insecticide and parasiticide. Fever, pain, respiratory illness, sedative, malaria, gastrointestinal problems, arthritis, and liver and kidney problems have all been treated the fruit juice and leaf and bark infusions. *A. muricata* has also been shown to have anti-diabetic and anti-tumor activity in recent studies.^[4]




Figure 1. Properties of Soursop.

The hepatoprotective effects of AM leaf extracts were effective in the normalization of function after liver damage, as determined by biochemical and histological tests^[5].

Because the fruit has economic value, it is widely cultivated and consumed as edible food. The plant's major pharmacological activities include cytotoxicity, antileishmanial activity, wound healing, and antimicrobial activity. It also has anticarcinogenic and genotoxic properties. The presence of tannins, steroids, and cardiac glycosides, which are the major phytochemical compounds, was revealed by phytochemical analysis of the plant^[6].

2. Drug Profile

- Synonyms: Annona Macrocarpa Wercle
- Kingdom: plantae
- Clade: magnoliids
- Order: magnoliales
- Family: annonaceae
- Genus: annona



Figure 2. *Annona Muricata* a) whole plant b) leaves c) flower d) fruit e) seeds.

The petals are yellowish and thick. Outer petals are broadly ovate, 2.8 centimeters (1.1 in) to 3.3 centimeters (1.3 in) by 2.1 centimeters (0.83 in) to 2.5 centimeters (0.98 in), tapering to a point with a heart-shaped base. They are evenly thick and covered with long, slender, soft hairs on the outside and finely matted soft hairs on the inside. The inner petals are oval and overlap. They are sharply angled and tapered at the base, measuring approximately 2.5 centimeters (0.98 in) to 2.8 centimeters (1.1 in) by 2 centimeters (0.79 in). The margins are thin, with fine matted soft hairs on both sides. The receptacle is hairy and conical. The stamens are 4.5 mm (0.18 in) long^[7]. Its pollen is shed in the form of permanent tetrads.^[8]

The fruits are prickly and dark green. They are ovoid in shape and can grow to be up to 30 centimeters (12 in) long, with a moderately firm texture. Their flesh is juicy, acidic, whitish in color, and aromatic^[9]

A. muricata is a member of the Annonaceae family, which includes approximately 130 genera and 2300 species.

A. muricata is common in the West Indies and northern South America. It has recently spread throughout the West Indies, as well as from southern Mexico to Peru and Argentina.^[10]

3. Cultivation

The plant is cultivated for its 20-30 cm (7.9-11.8 in) long, prickly, green fruit, which can weigh up to 6.8 kg (15 lb), making it the second largest *Annona* after the jungles. Away from its native range, some limited production occurs as far north as southern Florida within USDA Zone 10, but this is mostly for garden plantings for local consumption. It is also grown in parts of Southeast Asia and is plentiful oninauritius. Mexico is the primary supplier of the fruit, followed by Peru, Brazil, Ecuador, Guatemala, and Haiti^[27]. *Annona Suricata*'s complete genome was sequenced in 2021 to aid soursop breeders and stimulate further development of genomic resources for this globally important plant family^[28].

4. Ethnomedicinal Uses

All parts of the *A. muricata* tree, like those of other *Annona* species such as *A. squamosal* and *A. reticulata*, are widely used as traditional medicines for a variety of human ailments and diseases, particularly cancer and parasitic infections. The fruit is used as a natural remedy for arthritis, aches and pains, neuralgia, arthritis diarrhea, dysentery, fever, malaria, parasites, rheumatism, skin rashes, and other symptoms to raise a mother's milk after childbirth. The leaves are used to Cystitis, diabetes, headaches, and insomnia can all be treated. Furthermore, internal administration of the decoction of the leaf is thought to have anti-rheumatic and neuralgic properties, whereas the cooked leaves are topically applied used in the treatment of abscesses and rheumatism^[12].

5. Phytochemistry

The extract of *A. muricata* contains over 212 different phytoconstituents.^[13] The chemicals most abundant and contributing to bioactivity are acetogenins, alkaloids, phenols, and flavonoids. Several studies have revealed the presence of flavonol triglycerides, alkaloids, phenolics, megastigmanes, cyclopeptides, and essential oils. Minerals such as calcium, sodium, iron, potassium, copper, and magnesium are obtained by the body through consumption.^[14]

The presence of various major minerals such as K, Ca, Na, Cu, Fe, and Mg suggests that eating *A. muricata* fruit regularly can help provide essential nutrients and elements to the human body^[15]




Figure 3. Phytochemical present in *A. muricata*.

5.1. Acetogenesis





Figure 4. Acetogenin compounds in *a. muricata* (a) linear structure (b) epoxy acetogenin (c) mono the (d) monotetrahydrofuran acetogenin (e) bis the nonadjacent acetogenin (f) and bis the adjacent.

Acetogenins are a distinct class of compounds that are thought to be the primary bioactive compound of the Annonaceae family and are thus also known as annonaceous acetogenins. More than 120 acetogenins have been identified in plant parts such as the leaves, stem, bark, seed, pulp, and fruit peel. The structure of the acetogenins determines *A. muricata*'s bioactivity. Annonacin is an important acetogenin found in abundance in the fruit and leaf, as well as in lesser amounts in the seeds, roots, and peel. According to some studies, acetogenins are more cytotoxic than alkaloids. [16]

5.2. Alkaloids



1

2

3

4

5

6

7

8

9

10

11

12

13

14

(a) (b) 1
Figure 5. The most abundant alkaloids in *A. muricata* (a) coreximine; (b) reticuline. 2

Muricata contains two important alkaloids: reticuline and core mine. The alkaloids 3
 are more abundant in the leaves and less so in the roots, stems, and fruits in that order. 4
 Isoquinoline, apomorphine, and protoberberine are the most important alkaloids found. 5
 The alkaloids extracted from *A. muricata* have an affinity for 5-HT_{1A} receptors, which 6
 participate in dopamine synthesis and thus have antidepressant and cytotoxic effects in 7
 the body.^[17] 8

5.3. Flavonoid 9




Figure 6. Flavonoid in *A. muricata* (a) kaempferol (b) quercetin (c) rutin. 10 11

The most abundant flavonoid in *A. muricata* is a. kaempferol b. quercetin and c. ruti- 12
 n. According to reports, *A. muricata* contains up to 37 phenolic compounds. The two most 13
 important are quercetin and gallic acid. Tocopherols and tocotrienols are abundant in the 14
 fruit's pulp. According to research, the majority of phenols are soluble in aqueous extract. 15
 Antioxidant properties have been reported for phenolic compounds.^[18] 16

5.4. Essential oil 17

GC and GC-MS analyses of *A. muricata* leaf oil from Cameroon revealed the pres- 18
 ence of mostly sesquiterpenes, with α -caryophyllene being the most abundant compound. 19
 Another study on *A. muricata* from Vietnam discovered significant α -pinene volatile oil 20
 constituents. Geracrene D (20.6%), α -mentha-2,4(8)- diene (9.8%), α -pinene (9.4%), and el- 21
 ements (9.1%) derived from leaf oil. Other important compounds include α -cadinene, 22
 epicasinol, and α -cardinal compounds allegedly discovered in leaf oil extracts. The essen- 23
 tial oil extracted from fruit pulp was discovered to have aliphatic acid esters containing 24
 the major compounds 2-hexenoic acid methyl ester and 2-hexenoic acid ester of ethyl 25
 High concentrations of mono- and sesquiterpenes, including α -caryophyllene, on the other 26
 hand, The fruit pulp also yielded 1,8-cineole and linalool.^[19] 27

5.5. Other compounds 28

Several studies have found the presence of other compounds in the leaves, seeds, 29
 and fruit pulp, such as vitamins, carotenoids, amides, and cyclopeptides. The seeds con- 30
 tain anti-inflammatory and anti-tumor amides called N-p-coumaroyl tyramine and cy- 31
 clopeptides. Around 37 volatile compounds have been identified in the fruit pulp, and 32
 sesquiterpene derivatives have been discovered in the leaf. Soursop pulp also contains 33
 enzymes like pectinase, catalase, and peroxidase. During ripening, the enzyme amylase is 34
 detected, and its activity increases approximately 18-fold as the ethylene level in the fruit 35
 increases.^[19] 36

6. Pharmacological Activities

6.1. Anticancer

A. Suricata's anticancer activity is linked to its cytotoxic activity against cancer cells. The effects of *A. muricata* on cancer cells are shown in Table 3. Extracts from various parts of *A. muricata* act as anticancer agents through a variety of mechanisms. Extracts of the fruit, stems, seeds, and twigs of *A. muricata* were reported to inhibit matrix metalloproteinases (MMPs) such as MMP-2 and MMP-9, which play important roles in cancer progression [59]. Extracts from the leaves, twigs, and roots inhibited the proliferation of the human leukemia cell line HL-60 by disrupting MMPs, reactive oxygen species (ROS), and the G0/G1 cell cycle arrest, resulting in cancer cell growth inhibition [20].

Ic₅₀ of several parts of muricate against cancer cell line

The A549 lung cancer cell line was treated with an ethyl acetate extract of *A. muricata* leaves, which induced apoptosis by increasing Bax expression and decreasing Bcl-2 expression. Apoptosis induced by an ethyl acetate extract of *A. muricata* leaf has also been linked to cell cycle arrest at the G0/G1 phase. In the colorectal cancer cell line COLO-205[21] and the breast cancer cell line MDA-MB-231. Another study found that *A. muricata* leaf ethanol extract and ethyl acetate fractions were active against MCF7 cells via an apoptosis mechanism mediated by decreased Bcl-2 expression and increased caspase-3 and caspase-9 expression [22].

6.2. Antiulcer

Flavonoids, tannins, and phenolic acids are abundant in *A. muricata* and have therapeutic effects due to their antioxidant, anti-inflammatory, and gastroprotective properties. According to a survey, the leaves and bark of *A. muricata* are commonly used to make tea to treat gastrointestinal problems like gastritis and poor digestion [23].

6.3. Antidiabetic

Americana also exhibits antidiabetic activity. It contains flavonoids that inhibit α -glucosidase activity through hydroxylation bonding and substitution at the bring. This inhibition suppresses carbohydrate hydrolysis and glucose absorption and inhibits carbohydrate metabolism into glucose.[24]

In vitro, *A. muricata* fruit extracts were found to have antioxidant and anti-diabetic effects by inhibiting key enzymes involved in type 2 diabetes mellitus, such as -amylase and -glucosidase. According to one study, *A. muricata* pericarp has the highest inhibitory enzyme and antioxidant properties. When compared to the standard drug, the fruit pulp and leaf extract inhibited -amylase and -glucosidase and reduced the rate of glucose assimilation into the blood after feeding [25]

Antidiabetic effects of *A. muricata* aqueous extract are mediated by antioxidant mechanisms. In streptozotocin-induced diabetic mice, *A. muricata* leaf extract reduced lipid peroxidation processes, which are a sign of oxidative stress, and indirectly affected insulin production and endogenous antioxidants [26].

6.4. Antibacterial

When compared to the standard antibiotic streptomycin, *A. muricata* extracts demonstrated antibacterial activity against Gram-positive and Gram-negative bacteria. However, the solvent used for extraction can have an impact on the bioactive efficacy of the extracts. The combination of *A. muricata* ethanolic extract and antibiotic treatment reduced the potential of antibiotic multidrug-resistant *Escherichia coli* and *Staphylococcus aureus* strains. Another study found that bioactive compounds in *A. muricata*, such as alkaloids (annonaine, asimilobine, liriodenirine, nornuciferine, and so on), attack the bacterial membrane (plasma and outer membrane), resulting in broad-spectrum antibacterial activity.[27]

6.5. Antiviral

Antiviral activity of *A. muricata* extract has been reported, for example, by interfering with the replication process of HIV-I. In another study, ethanolic extracts of *A. muricata* stem and bark demonstrated in vitro antiviral activity against the herpes simplex virus. Another study found that after 1 hour of contact time, an acidified ethanolic extract of *A. muricata* reduced viral replication^[28]. This activity could be attributed to the presence of phenolic compounds like rutin. Acetogenins with good inhibitory activity against SARS- CoV-2 spike proteins include annomuricin a, annomuricin b, annomuricin c, muricatocin c, Suricata in, cis- annonacin, annonacin-10-one, cis-goniothalamycin, Ariana can, and javoricin (in silico)^[29]. Cis-annonacin had the lowest binding energy and the greatest ability to form hydrogen bonds, indicating that it was the most potent of the acetogenins tested in the study. This finding suggests that annonaceous acetogenins have the potential to be anti-SARS-CoV-2 agents and should be studied in vitro and in vivo ^[30]

Table 1. Medicinal uses of *A. Muricata*.

Plant part	Medicinal use	Preparation/application
Leaf	Insomnia, febrifuge, catarrh, snake bite, analgesic, arthritis pain, rheumatism, neuralgia, weight loss, malaria, digestive, typhoid fever, antimicrobial	Oral
Fruit	hypertension, diarrhea, dysentery, febrifuge, inflammation, lactagogue,a stringent	Juice/oral
Bark	Helminthiasis, diarrhea, stomach ulcer, Sedative, vermifuge, antidot	Decoction
Root	Dysentery, vermifuge, antidote, insomnia Catarrh, febrifuge, spasms, parasites	Decoction /oral
Seed	Sedative, cardi tonic, convulsion, renal and skin disorder, vaginal infection	Plaster/topical Infusion/oral

7. Nutritional Facts

One cup (225g) of soursop pulp contains 148 calories, 2.3g protein, 37.8g carbohydrates, and 0.7g fat. *Annona muricata* is high in vitamin C, fibre, and potassium. ^[31]

Calories:14;
 Fat:0.7g;
 Carbohydrades:37.8g;
 Fiber:7.4g;
 Sugars:30.5g;
 Protein:2.3g;
 Vitamin C :46.4mg;
 Potassium:626mg.

7.1. Crabs:

Annona muricata has nearly 38 grammes of carbohydrates per cup. Soursop's carbs come from naturally occurring sugars, and it has more than 7 grammes of fibre per serving (about a quarter of your recommended daily intake). Sorrel has a low glycemic index.^[32]

7.2. Fats:

This fruit contains very little fat, with less than one gramme per serving.

7.3. Protein:

Annona muricata like most fruits, contains little protein. A single serving of protein contains only 2.3 grammes. To meet your daily protein requirements, you'll need to include other protein sources in your diet, such as salmon, lean meats, and legumes .

7.4. Vitamins And Minerals:

Soursop, like most fruits, has a low protein content. Protein is only 2.3 grammes in a single serving. Other protein sources, such as salmon, lean meats, and legumes, will help you meet your daily protein requirements ^[33].

8. Conclusion

A. muricata is an evergreen plant that is found almost everywhere. *A. muricata* extract contains over 212 different phytoconstituents. Annonaceous acetogenins are important chemical constituents found almost everywhere in plants that contribute to human health problems. This plant's alkaloids, flavonoids, and phenols exhibit a wide range of activity. People all over the world use a decoction of the plant's fruit, leaves, roots, and bark for a variety of purposes. The plant's cytotoxic, antiparasitic, anti-inflammatory, and antioxidant properties are the most promising. Previous studies focused primarily on the biological activity of the plant extract. More research is needed on the biochemical and physiological functions of active compounds, as well as the detailed mechanisms underlying these activities. Several reports suggest that the important constituent annonacin has a neurotoxicological effect. A clear distinction between chemical constituents, dose, and toxicity is critical. Future research and clinical trials must focus on the compounds that contribute to toxicity, as well as their dose and effect on the body. This article is intended to provide researchers with information and motivation to continue studying this plant.

Funding: This review received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We would like to convey our obligation to Management and Principal, P.S.G.V.P. Mandal's College of Pharmacy, Shahada, Dist. Nandurbar.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Moghadamtousi SZ, Goh BH, Chan CK, Shabab T, Kadir HA. Biological activities and phytochemicals of *Swietenia macrophylla* king. *Molecules* 2013; 18:10465-10483. www.nhs.uk was first indexed by Google more than 10 years ago, <https://www.nhs.uk/live-well/addiction-support/addiction-what-is-it/>
2. Duraipandiyan V, Ayyanar M, Ignacimuthu S, Antimicrobial Activity of Some Ethnomedical Plants Used by Paliyar Tribe from Tamil Nadu, India. *BMC complementary and alternative medicine*. 20A
3. Sawant TP, Dongre RS. Bio-chemical compositional analysis of *Annona muricata*: a miracle fruit's review. *International Journal of Universal Pharmacy and Bio Sciences*. 2014;2(3):82-104.Christophe Gleye, Alain Laurens, Reynald

- Hocquemiller, Olivier Laprevote, Laurent Serani and Andre Cave, Cohibins A and B, Acetogenins from roots of *Annona muricata*. *Phytochemistry*, 1997; 44(8): 1541 -1545BMJ.
4. Gavamukulya Y, Wamunyokoli F, El-Shemy HA. *Annona muricata*: Is the natural therapy for most disease conditions including cancer growing in our backyard? A scientific review of its research history and prospects. *Asian Pac JTrop Dis*. 2017;10(9):835-48.
 5. Arthur FKN, Larbie C, Woode E, Terlabi EO. Evaluation of the hepatoprotective effect of aqueous extract of *Annona muricata* (Linn.) leaf against carbon tetrachloride acetaminophen-induced liver damage. *J Nat Pharm*. 2012;3(1)
 6. Gajalakshmi S, Vijayalakshmi S, Rajeshwari Devi V. Phytochemical and Pharmacological Properties of *Annona muricata*: A Review, *Int J Pharm Sci*. 2012; 4(2):3-6.
 7. http://www.efloras.org/florataxon.aspx?flora_id=3&taxon_id=200008507 Floras.org. Retrieved 2008-04
 8. Walker JW (1971) Pollen Morphology, Phyto geography, and Phylogeny of the Annonaceae. *Contributions from the Gray Herbarium of Harvard University*, 202: 1-130.
 9. http://www.hear.org/pier/species/annona_muricata.html Institute of Pacific Islands Forestry: Pacific Island Ecosystems at Risk (PIER). 2008-01-05. Archived from the original on 12 May 2008. Retrieved 2008-04-18
 10. Morton JF. *Soursop (Annona muricata)*. Fruits of warm climates. Purdue University. 1987:75-80
 11. Strijk, Joeri S.; Hinsinger, Damien D.; Roeder, Mareike M.; Chatrou, Lars W.; Couvreur, Thomas L. P.; Erkens, Roy H. J.; Sauquet, Hervé; Pirie, Michael D.; Thomas, Daniel C.; Cao, Kunfang (2021). "Chromosome-level reference genome of the soursop (*Annona muricata*): A new resource for Magnoliid research and tropical pomology". *Molecular Ecology Resources*. 21 (5): 1608–1619. doi:10.1111/1755-0998.13353. ISSN 1755-0998. PMC 8251617. PMID 33569882.
 12. Mishra, S.; Ahmad, S.; Kumar, N.; Sharma, B.K. *Annona muricata* (the cancer killer): A review. *Glob. J. Pharm. Res*. 2013, 2, 1613–1618.
 13. Adewole, S.O.; Caxton-Martins, E.A. Morphological changes and hypoglycemic effects of *Annona muricata* Linn. (*Annonaceae*) leaf aqueous extract on pancreatic B-cells of streptozotocin-treated diabetic rats. *Afr. J. Biomed. Res*. 2006, 9, 173–187.
 14. Prasad SK, Varsha V, Devegowda D. Anti-cancer properties of *Annona muricata* (L.): A Review. *Medicinal Plants*. 2019;11(2):117-29.
 15. J. Nawrocka and U. Małolepsza, "Diversity in plant systemic resistance induced by *Trichoderma*,"
 16. *Biological Control*, vol. 67, no. 2, pp. 149–156, 2013.
 17. C. M. J. Pieterse, D. Van der Does, C. Zamioudis, A. Leon-Reyes, and S. C. M. Van Wees, "Hormonal modulation of plant immunity," *Annual Review of Cell and Developmental Biology*, vol. 28, no. 1, pp. 489– 521, 2012.
 18. W. Zárate-Martínez, S. González-Morales, F. Ramírez-Godina, A. Robledo-Olivo, and A. Juárez-Maldonado, "Efecto de los ácidos fenólicos en plantas de tomate (*Lycopersicon esculentum* Mill.) inoculadas con *Clavibacter michiganensis*," *Revista mexicana de ciencias agrícolas*, vol. 9, pp. 4367–4379, 2018.
 19. Coria-Téllez AV, Montalvo-González E, Yahia EM, Obledo-Vázquez EN. *Annona muricata*: A comprehensive review of its traditional medicinal uses, phytochemicals, pharmacological activities, mechanisms of action, and toxicity. *Arab J Chem*. 2018;11(5):662-91.
 20. Patel MS, Patel JK. A review of miracle fruit of *Annona muricata*. *J Pharmacogn Phytochem*. 2016;5(1):137-48.
 21. Jirovetz, L.; Buchbauer, G.; Ngassoum, M.B. Essential oil compounds of the *Annona muricata* fresh fruit pulp from Cameroon. *J. Agric. Food Chem*. 1998, 46, 3719–37
 22. Drishya G., Nambiar J., Shaji S.K., Vanuopadath M., Achuthan A., Kumar A., Alias A., Sherif A., Joseph C., Divya P., et al. RECK and TIMP-2 Mediate Inhibition of MMP-2 and MMP-9 by *Annona muricata*. *J. Biosci*. 2020; 45:1. Doi: 10.1007/s12038-020-00056-z.
 23. Pieme A.A., Kumar G.G., Dongmo S.S., Moukette M.M., Boyoum F.F., Ngogang Y.Y., Saxena K.K. Antiproliferative Activity and Induction of Apoptosis by *Annona muricata* (*Annonaceae*) Extract on Human Cancer Cells. *BMC Complement. Altern. Med*. 2014; 14:1–10. DOI: 10.1186/1472-6882-14-516.
 24. Moghadamtousi S.Z., Kadir H.A., Paydar M., Rouhollahi E., Karimian H. *Annona muricata* Leaves Induced Apoptosis in A549 Cells through Mitochondrial-Mediated Pathway and Involvement of NF- κ B. *BMC Complement. Altern. Med*. 2014; 14:1–13. DOI: 10.1186/1472-6882-14-299.
 25. Abdullah M., Syam A.F., Meilany S., Laksono B., Prabu O.G., Bektı H.S., Indrawati L., Makmun D. The Value of Caspase-3 after the Application of *Annona muricata* Leaf Extract in COLO-205 Colorectal Cancer Cell Line. *Gastroenterol. Res. Pract*. 2017; 2017:165. DOI: 10.1155/2017/4357165.
 26. Kim J.Y., Dao T.T.P., Song K., Park S.B., Jang H., Park M.K., Gan S.U., Kim Y.S. *Annona muricata* Leaf Extract Triggered Intrinsic Apoptotic Pathway to Attenuate Cancerous Features of Triple Negative Breast Cancer MDA-MB-231 Cells. *Evid. -Based Complement. Altern. Med*. 2018; 2018:916. doi: 10.1155/2018/7972916.
 27. Hansra D.M., Silva O., Mehta A., Ahn E. Patient with Metastatic Breast Cancer Achieves Stable Disease for 5 Years on Graviola and Xeloda after Progressing on Multiple Lines of Therapy. *Adv. Breast Cancer Res*. 2014; 3:47526. DOI: 10.4236/abcr.2014.33012.
 28. Hardoko Y.H., Halim Y., Wijoyo S.V. In Vitro Antidiabetic Activity of "Green Tea" Soursop Leaves Brew through α -Glucosidase Inhibition. *Int. J. PharmTech Res*. 2015; 8:30–37.

29. Agu K.C., Eluehike N., Ofeimun R.O., Abile D., Ideho G., Ogedengbe M.O., Onose P.O., Elekofehinti O.O. Possible Anti-Diabetic Potentials of *Annona muricata* (Soursop): Inhibition of α -Amylase and α -Glucosidase Activities. *Clin. Phytosci.* 2019; 5:1–13. Doi: 10.1186/s40816-019-0116
30. Adewole S.O., Ojewole J.A.O. Protective Effects of *Annona muricata* Linn. (Annonaceae) Leaf Aqueous Extract on Serum Lipid Profiles and Oxidative Stress in Hepatocytes of Streptozotocin-Treated Diabetic Rats. *Afr. J. Tradit. Complement. Altern. Med.* 2009;6 DOI: 10.4314/at cam. v6i1.57071
31. Vieira G.H.F., Mourão J.A., Ângelo Â.M., Costa R.A., Vieira R.H.S.D.F. Antibacterial Effect (in Vitro) of *Moringa Oleifera* and *Annona muricata* against Gram Positive and Gram-Negative Bacteria. *Rev. Inst. Med. Trop. Sao Paulo.* 2010; 52:129–132. doi: 10.1590/S0036-46652010000300003.
32. Prasad S.K., Pradeep S., Shimavallu C., Kollur S.P., Syed A., Marraiki N., Egbuna C., Gaman M.A., Kosakowska O., Cho W.C., et al. Evaluation of *Annona muricata* Acetogenins as Potential Anti-SARS-CoV-2 Agents Through Computational Approaches. *Front. Chem.* 2021; 8:1–7. doi: 10.3389/fchem.2020.624716.
33. <https://www.verywellfit.com/soursop-nutrition-facts-4687162> - toc-soursop-nutrition-facts