

Benefits and Drawbacks on the Incorporation of Grape Seeds into Bakery Products: Is It Worth It? †

J. Echave ¹, A. Silva ^{1,2}, A.G. Pereira ¹, P. Garcia-Oliveira ^{1,3}, M. Fraga-Corral ^{1,3}, Paz Otero ^{1,*}, L. Cassani ^{1,3}, Cao H ¹, J. Simal-Gandara ¹, M.A. Prieto ^{1,3} and J. Xiao ^{1,4,*}

¹ Nutrition and Bromatology Group, Department of Analytical Chemistry and Food Science, Faculty of Science, Universidade de Vigo, E32004 Ourense, Spain.; javier.echave@uvigo.es (J.E.); mass@isep.ipp.pt (A.S.); antia.gonzalez.pereira@uvigo.es (A.G.P.); paula.garcia.oliveira@uvigo.es (P.G.-O.); mfraga@uvigo.es (M.F.-C.); lucia.victoria.cassani@uvigo.es (L.C.); hui.cao@uvigo.es (C.H.); jsimal@uvigo.es (J.S.-G.); mprieto@uvigo.es (M.A.P.)

² REQUIMTE/LAQV, Instituto Superior de Engenharia do Porto, Instituto Politécnico do Porto, Rua Dr António Bernardino de Almeida 431, 4200-072 Porto, Portugal

³ Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolonia, 5300-253 Bragança, Portugal

⁴ International Joint Research Laboratory of Intelligent Agriculture and Agri-products Processing, Jiangsu University, Zhenjiang 212013, China

* Correspondence: paz.otero@uvigo.es (P.O.); jianbo.xiao@uvigo.es (J.X.)

† Presented at the 2nd International Electronic Conference on Processes: Process Engineering—Current State and Future Trends (ECP 2023), 17–31 May 2023; Available online: <https://ecp2023.sciforum.net/>.

Abstract: Grape is the third most produced fruit in the world, owing to its taste and its use as raw material for winemaking. Due to this fact, large volumes of waste biomass are generated as a result of grape juice and wine production, mainly grape seeds (GS) and peels. In recent decades, scientific research has highlighted the high content of polyphenols in GS, especially condensed tannins, and resveratrol. These compounds have been correlated to various potential benefits to human health, such as antioxidant, hypoglycemic, hypolipidemic or anti-inflammatory, among others. GS polyphenols may be incorporated into functional foods, especially as flour in wheat-based bakery products, appears to be an attractive option. Two strategies may be followed. By one hand, the incorporation of GS flour increases fiber, mineral and protein content of bakery products, as well as their hardness and phenolic content. However, it seems that consumers may accept up to 10% of GS flour since higher doses strongly diminish the organoleptic properties of the product. The other alternative involves the incorporation of polyphenol-rich GS extracts into the bakery formulations which would carry their beneficial bioactivities to final product. Nonetheless, this method is more laborious since it requires a prior chemical extraction of GS, and the control and addition of a safe, food-grade extract into the flour. Both strategies have been reported to increase phenolic content and antioxidant capacity of bakery products. The direct incorporation of GS flour is affordable for industries while the incorporation of polyphenol-rich extracts leads to the development of functional products with additional beneficial properties. This work discusses the benefits and potential hurdles of functional bakery products with incorporated GS flour and extracts based on up-to-date evidence.

Keywords: grape seeds; bakery products; polyphenols; functional foods; circular economy

Citation: Echave, J.; Silva, A.; Pereira, A.G.; Garcia-Oliveira, P.; Fraga-Corral, M.; Otero, P.; Cassani, L.; Cao, H.; Simal-Gandara, J.; et al. Benefits and Drawbacks on the Incorporation of Grape Seeds into Bakery Products: Is it Worth it? *Eng. Proc.* **2023**, *37*, x. <https://doi.org/10.3390/xxxxx> Published: 17 May 2023



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Winemaking generates waste in the form of grape pomace, seeds, and skins that remain after the wine has been pressed. Based on some estimates, approximately 20% of the grape total weight is disposed as waste in the form of seeds, stems, and skins, which can represent over 8 million tons worldwide of grape pomace waste [1]. Of this, grape seeds

(GS) constitute the most environmentally persistent. Removal of this waste is costly, and if it is not treated effectively, it may pose an environmental issue, since these seeds contain high levels of condensed tannins (CT) and other phenolic compounds (PC) that if concentrated, may have negative impacts on the soil [1]. PC, however, have attracted a great deal of interest both by the industry and academia, since they have been reported to exert antioxidant, hypoglycemic, anti-obesity, anti-inflammatory, or cardioprotective effects, among many others [2]. Grape seeds have a high content of PC such as phenolic acids, monomeric PC (e.g., flavanols such as catechin or epicatechin), polymeric CT and in the case of red grape varieties, several anthocyanins [3]. In addition, GS is one of the richest natural sources of resveratrol, a stilbene accounted with highly promising anti-obesity effects [4]. On the other hand, there is an increasing demand of healthier foods prepared with natural ingredients, and this has given rise to functional foods, which have potential health benefits besides their nutritional ones [5]. Therefore, one possible alternative to the management of GS as organic waste, has been its application towards fortification of several foodstuff, of which bakery products have emerged as the most feasible to produce, attractive to consumers and easy to commercialize. Moreover, bakery products made with refined wheat flour generally lack a great quantity of nutrients and are often described as highly glycemic foods, since their only value is as highly energetic foods [6].

Non-conventional flours have been studied to increase the nutritional value of traditional ones, such as wheat, and compensate for the loss of nutrients that occurs after the refining process. For example, a mixture of flours from other options that are considered in baking, to which GS are a promising candidate [7]. Improvements in technological and nutritional characteristics have already been observed, without compromising the final acceptance of bakery products, resulting in potential health benefits [8]. As such, GS as part of flour appears as a promising alternative to develop fortified foods. Nonetheless, PC-rich extracts may be obtained from GS and applied to flour formulations, as a functional alternative to enrich these products (Figure 1).




Figure 1. Diagram of grape seed revalorization as a functional ingredient for bakery products.

2. Incorporation into Bakery Products

2.1. Incorporation of Grape Seed Flour

GS is composed of approximately 40% fiber and 7% complex phenols, including CT, in addition to about 10% protein and other constituents, such as fatty acids, sterols, or minerals [9]. This means that, although the PC and fiber content of GS is the main reason for its health benefits, the organoleptic properties of the final bakery products will be heavily altered. In particular, it has been consistently observed that combining GS flour

with other flours and particularly wheat, generally gives place to much denser doughs, at least on combinations up to 10% of the flour weight [9]. This generally leads to poorer gasification of the dough during fermentation and oven, in a positively correlation with increasing GS flours concentrations, making a denser dough [10]. The GS flour concentration may be increased up to 20%, but rheological and organoleptic properties are negatively affected to the point the product may be unacceptable [9]. In addition, although the addition of GS may negatively alter the textural properties of the dough, it has been observed that elasticity and stability is positively improved [5]. This probably due to its increased water absorption capacity, which impacts the moisture of the final product [11]. Moreover, there is a significant alteration of taste, as with increased GS concentrations PC (and especially CT) provide the bread with a strongly bitter/sour taste, a common feature of PC and which can be more nuanced in complex ones [3].

However, considering the GS composition, this provides the end-product with additional fiber and mineral content. This also results in increased antioxidant properties, as the incorporated GS flour retains PC embedded in the seed tissue. It has been reported that white grape pomace flour addition (10%) can increase the fiber content of fortified breads by 88% and increased levels of phenolic acids from 0.1 mg/g to 0.3 mg/g [12]. However, as the GS flour is kept at relatively low proportions, this antioxidant effect is much lower than that of GS alone [13]. For example, it was reported that following an *in vitro* ferric reducing power (FRAP) assay, bread made with 6% GS flour, achieved an antioxidant effect more than 10 times higher than the control bread, with 590 mg Trolox equivalents (TE)/100 g vs. 45 mg TE/100 g, but was much lower than the unmixed GS (2775 mg TE/100 g) [13]. Similar results were obtained in other recent studies, with 6% GS addition reaching 346 mg TE/100 g, while the control bread scored 70 mg TE/100 g [14]. This evidences that GS flour incorporation may prove beneficial to some extent, considering its very low economic cost and antioxidant properties. However, such effects are also limited, since the resulting bakery product will not be accepted if more than 10% of GS is added due to resulting poor texture and flavor [15].

2.2. Incorporation of Grape Seed Extracts

Another strategy, involves the processing of GS to obtain chemical extracts with high concentrations of polyphenols, which may be added to the flour, thus working as a natural additive. This fortification may provide the end product with natural antioxidants, thus furthering their range of action and effectiveness [16]. The application of GS extracts on breads has been accounted with greatly ameliorating acrylamide production during baking (66 µg/kg vs. 296 µg/kg in the control), as well with improved antioxidant capacity [17]. In contrast to the incorporation of GS flour, the addition of GS extracts is accounted to not significantly alter the texture of the produced breads [17]. Thus, not only the incorporation of GS extracts adds beneficial phytochemicals to the end-product, but also prevents the negative alterations flour incorporation may involve. Indeed, it appears as if PC are better retained in the form of extracts than in the form of flour. For example, following measure by Total Phenolic Content (TPC) analyses, GS flour breads displayed up to 45 mg gallic acid equivalents (GAE)/100 g [18], with a 5% substitution, while 1.5% addition of GS extracts increased this value more than 10 times, up to 580 mg GAE/100 g [17].

GS extracts have been applied to other foods such as mackerel, showing improved storage stability and delay of degradation [19]. In this sense, GS extracts have showed improved delay of spoilage by microorganisms [16]. Moreover, the anti-obesity and antioxidant effect of GS extracts is well-established [20]. Indeed, intake of GS extracts has been reported to inhibit postprandial glycemia in small-scale clinical trials. It was showed that even when subjects had high-calorie meals, a daily intake of 100 mg of GS extract could ameliorate glucose levels (5.63 mmol/L vs. 6.99 mmol/L) [21]. Therefore, it is suggested that, given that PC from GS are retained in the bakery product even after baking and at high concentrations, it can exert these anti-glycemic and anti-obesity effects more effectively as added extracts [17]. This was demonstrated on an *in vivo* study, on which breads

fortified with grape skin extracts (1.4%) significantly reduced fat and protein digestion of treated rats. Being fed a high fat diet, supplementation with the fortified bread powder resulted in lower low-density lipoprotein (LDL) cholesterol levels, decreased leptin secretion, and glycemia, among other markers. This suggested that grape waste-derived extracts were effective in ameliorating diet-induced obesity symptoms [22]. Notably, in the same study flour supplementation achieved slightly better results, but this was coupled with a more negative impact in organoleptic properties that the extract addition did not show [22]. There is although, a general lacking on more accurate and concise studies directly assessing the effect of GS extract addition on critical quality parameters such as storage time, organoleptic properties alterations and direct potential health benefits from the intake of the formulated bakery products. On the other hand, many studies assess whole pomace or skins, making difficult to discern the effect of seeds alone [12].

3. Conclusions

The valorization of winemaking wastes such as GS can be easily and beneficially re-integrated into the food industry. This waste represents natural sources of antioxidants whose reutilization enables the reduction of synthetic analogues related to negative side effects. The incorporation of these antioxidants as preservatives maintains the stability of the final product, while potential health benefits may be added. Moreover, this reutilization of GS can be an effective strategy to produce added-value products while reducing volumes of this biomass destined to landfilling, thus promoting a circular economy model. This fact can contribute to the higher consumer's perceived value that would balance the cost of the development of new formulations and optimization of the food-making processes. Considering both strategies of incorporation of GS and derived extracts into the final product, it can be suggested that GS flour appears as the most feasible and economic practice, in concentrations up to 10% of the total flour weight. However, this limitation is accompanied of a proportional limitation of the antioxidant activity of the fortified bakery products but if such limit is exceeded, acceptability of the product may be compromised. In this sense, the inclusion of GS extracts as natural additives results in higher antioxidant properties of the product, with reduced alteration of organoleptic properties. Altogether, considering available reports, it would seem that the current best strategy to develop bakery products valorizing GS would be to apply PC-rich extracts. Although, this could lead to additional costs in the end-product that could act as deterrents to its consumer acceptability. Further research could unveil alternative optimized methods to compensate current shortcomings to valorize GS for functional bakery products.

Author Contributions: Conceptualization, J.E., J.X. and P.G.P.; methodology, J.E. and P.G.-O.; software, C.H.; validation, M.F.-C. and P.O.; formal analysis, P.G.-O. and L.C.; investigation, J.E., A.S. and A.G.P.; resources, P.O.; data curation, P.G.-O., L.C. and P.O.; writing—original draft preparation, J.E., M.F.-C., A.G.P. and A.S.; writing—review and editing, J.E. and M.F.-C. visualization, P.O. and A.S.; supervision, M.A.P., M.F.-C. and J.X.; project administration, J.X., C.H. and J.S.-G.; funding acquisition, J.S.-G., M.A.P. and J.X. All authors have read and agreed to the published version of the manuscript.

Funding: The research is supported by MICINN supporting the Ramón y Cajal grant for M.A. Prieto (RYC-2017-22891) and Jianbo Xiao (RYC-2020-030365-I), and the Juan de la Cierva Incorporación for Hui Cao (IJC2020-046055-I). By Xunta de Galicia for supporting the program EXCELENCIA-ED431F 2020/12, the program EXCELENCIA-ED431F 2022/01, the post-doctoral grant of M. Fraga-Corral (ED481B-2019/096), and L. Cassani (ED481B-2021/152), and the pre-doctoral grants of P. Garcia-Oliveira (ED481A-2019/295), A.G. Pereira (ED481A-2019/0228).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Acknowledgments: Authors are grateful to Ibero-American Program on Science and Technology (CYTED—AQUA-CIBUS, P317RT0003), to the Bio Based Industries Joint Undertaking (JU) under grant agreement No 888003 UP4HEALTH Project (H2020-BBI-JTI-2019) that supports the work of P. Otero. The JU receives support from the European Union’s Horizon 2020 research and innovation program and the Bio Based Industries Consortium.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Tița, O.; Lengyel, E.; Stegăruș, D.I.; Săvescu, P.; Ciubara, A.B.; Constantinescu, M.A.; Tița, M.A.; Rață, D.; Ciubara, A. Identification and Quantification of Valuable Compounds in Red Grape Seeds. *Appl. Sci.* **2021**, *11*, 5124. <https://doi.org/10.3390/app11115124>.
2. Jimenez-Lopez, C.; Fraga-Corral, M.; Carpena, M.; García-Oliveira, P.; Echave, J.; Pereira, A.G.; Lourenço-Lopes, C.; Prieto, M.A.; Simal-Gandara, J. Agriculture waste valorisation as a source of antioxidant phenolic compounds within a circular and sustainable bioeconomy. *Food Funct.* **2020**, *11*, 4853–4877. <https://doi.org/10.1039/D0FO00937G>.
3. Boff, J.M.; Strasburg, V.J.; Ferrari, G.T.; de Oliveira Schmidt, H.; Manfroi, V.; de Oliveira, V.R. Chemical, Technological, and Sensory Quality of Pasta and Bakery Products Made with the Addition of Grape Pomace Flour. *Foods* **2022**, *11*, 3812. <https://doi.org/10.3390/foods11233812>.
4. Tabrizi, R.; Tamtaji, O.R.; Lankarani, K.B.; Akbari, M.; Dadgostar, E.; Dabbaghmanesh, M.H.; Kolahdooz, F.; Shamshirian, A.; Momen-Heravi, M.; Asemi, Z. The effects of resveratrol intake on weight loss: A systematic review and meta-analysis of randomized controlled trials. *Crit. Rev. Food Sci. Nutr.* **2020**, *60*, 375–390. <https://doi.org/10.1080/10408398.2018.1529654>.
5. Meral, R.; Doğan, İ.S. Grape seed as a functional food ingredient in bread-making. *Int. J. Food Sci. Nutr.* **2013**, *64*, 372–379. <https://doi.org/10.3109/09637486.2012.738650>.
6. Coe, S.; Ryan, L. White bread enriched with polyphenol extracts shows no effect on glycemic response or satiety, yet may increase postprandial insulin economy in healthy participants. *Nutr. Res.* **2016**, *36*, 193–200. <https://doi.org/10.1016/j.nutres.2015.10.007>.
7. Maman, R.; Yu, J. Chemical Composition and Particle Size of Grape Seed Flour and Their Effects on the Characteristics of Cookies. *J. Food Res.* **2019**, *8*, 111. <https://doi.org/10.5539/jfr.v8n4p111>.
8. Domínguez Díaz, L.; Fernández-Ruiz, V.; Cámara, M. An international regulatory review of food health-related claims in functional food products labeling. *J. Funct. Foods* **2020**, *68*, 103896. <https://doi.org/10.1016/j.jff.2020.103896>.
9. Koca, I.; Tekguler, B.; Yilmaz, V.A.; Hasbay, I.; Koca, A.F. The use of grape, pomegranate and rosehip seed flours in Turkish noodle (erişte) production. *J. Food Process. Preserv.* **2018**, *42*, e13343. <https://doi.org/10.1111/jfpp.13343>.
10. Tolve, R.; Simonato, B.; Rainero, G.; Bianchi, F.; Rizzi, C.; Cervini, M.; Giuberti, G. Wheat Bread Fortification by Grape Pomace Powder: Nutritional, Technological, Antioxidant, and Sensory Properties. *Foods* **2021**, *10*, 75. <https://doi.org/10.3390/foods10010075>.
11. Samohvalova, O.; Grevtseva, N.; Brykova, T.; Grigorenko, A. The effect of grape seed powder on the quality of butter biscuits. *Eastern-European J. Enterp. Technol.* **2016**, *3*, 61–66. <https://doi.org/10.15587/1729-4061.2016.69838>.
12. Mildner-Szkudlarz, S.; Bajerska, J.; Zawirska-Wojtasiak, R.; Górecka, D. White grape pomace as a source of dietary fibre and polyphenols and its effect on physical and nutraceutical characteristics of wheat biscuits. *J. Sci. Food Agric.* **2013**, *93*, 389–395. <https://doi.org/10.1002/jsfa.5774>.
13. Šporin, M.; Avbelj, M.; Kovač, B.; Možina, S.S. Quality characteristics of wheat flour dough and bread containing grape pomace flour. *Food Sci. Technol. Int.* **2018**, *24*, 251–263. <https://doi.org/10.1177/1082013217745398>.
14. Kapcsándi, V.; Hanczné Lakatos, E.; Sik, B.; Linka, L.Á.; Székelyhidi, R. Antioxidant and polyphenol content of different *Vitis vinifera* seed cultivars and two facilities of production of a functional bakery product. *Chem. Pap.* **2021**, *75*, 5711–5717. <https://doi.org/10.1007/s11696-021-01754-0>.
15. Marcos, J.; Carriço, R.; Sousa, M.J.; Palma, M.L.; Pereira, P.; Nunes, M.C.; Nicolai, M. Effect of Grape Pomace Flour in Savory Crackers: Technological, Nutritional and Sensory Properties. *Foods* **2023**, *12*, 1392. <https://doi.org/10.3390/foods12071392>.
16. Krasteva, D.; Ivanov, Y.; Chengolova, Z.; Godjevargova, T. Antimicrobial Potential, Antioxidant Activity, and Phenolic Content of Grape Seed Extracts from Four Grape Varieties. *Microorganisms* **2023**, *11*, 395. <https://doi.org/10.3390/microorganisms11020395>.
17. Chen, G.; Wang, J.; Li, Y. Extracts of sorghum bran, grape seed, and green tea: Chromatographic comparison of phenolic profiles and mitigation effect on acrylamide in antioxidant-fortified bread. *Food Chem. Adv.* **2022**, *1*, 100082. <https://doi.org/10.1016/j.focha.2022.100082>.
18. Elkatry, H.O.; Ahmed, A.R.; El-Beltagi, H.S.; Mohamed, H.I.; Eshak, N.S. Biological Activities of Grape Seed By-Products and Their Potential Use as Natural Sources of Food Additives in the Production of Balady Bread. *Foods* **2022**, *11*, 1948. <https://doi.org/10.3390/foods11131948>.
19. Sofi, F.R.; Raju, C.V.; Lakshmisha, I.P.; Singh, R.R. Antioxidant and antimicrobial properties of grape and papaya seed extracts and their application on the preservation of Indian mackerel (*Rastrelliger kanagurta*) during ice storage. *J. Food Sci. Technol.* **2016**, *53*, 104–117. <https://doi.org/10.1007/s13197-015-1983-0>.

20. Fernández-Iglesias, A.; Pajuelo, D.; Quesada, H.; Díaz, S.; Bladé, C.; Arola, L.; Salvadó, M.J.; Mulero, M. Grape seed proanthocyanidin extract improves the hepatic glutathione metabolism in obese Zucker rats. *Mol. Nutr. Food Res.* **2014**, *58*, 727–737. <https://doi.org/10.1002/mnfr.201300455>.
21. Sapwarobol, S.; Adisakwattana, S.; Changpeng, S.; Ratanawachirin, W.; Tanruttanawong, K.; Boonyarit, W. Postprandial blood glucose response to grape seed extract in healthy participants: A pilot study. *Pharmacogn. Mag.* **2012**, *8*, 192–196. <https://doi.org/10.4103/0973-1296.99283>.
22. Mildner-Szkudlarz, S.; Bajerska, J. Protective effect of grape by-product-fortified breads against cholesterol/cholic acid diet-induced hypercholesterolaemia in rats. *J. Sci. Food Agric.* **2013**, *93*, 3271–3278. <https://doi.org/10.1002/jsfa.6171>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.