

Recovering Antioxidant Compounds from the Juçara Residue by a Green Approach [†]

Matheus Lima ^{1,2}, Ravena Silva ^{1,3}, Alex Novo ¹, Claudete Kunigami ¹, Eliane Jung ¹ and Leilson Ribeiro ^{1,*}

¹ Laboratory of Organic and Inorganic Chemical Analysis, National Institute of Technology, Rio de Janeiro 20081-312, Brazil; matheus.lima@int.gov.br (M.L.); ravena.rodrigues@int.gov.br (R.S.); alex.novo@int.gov.br (A.N.); claudete.kunigami@int.gov.br (C.K.); eliane.jung@int.gov.br (E.J.)

² Escola de Química, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brasil

³ Faculdade de Farmácia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brasil

* Correspondence: leilson.oliveira@int.gov.br

[†] Presented at the 4th International Electronic Conference on Foods, 15–30 October 2023; Available online: <https://foods2023.sciforum.net/>.

Abstract: Juçara (*Euterpe edulis*) is a Brazilian Atlantic Forest palm that produces phenolic-rich fruits, used for pulp preparation. The pulp refining produces antioxidant and lipid-rich residue. This study applied glycerol as a green solvent for microwave-assisted extraction of antioxidant compounds from residue. Variables included glycerol concentration (0–100%), microwave time (2–18 min), and temperature (53–187 °C). Total phenolic compounds (TPC) and DPPH[•] antioxidant capacity were influenced by glycerol concentration and temperature ($p < 0.05$). TPC ranged from 1052 to 5329 mg GAE/100g and antioxidant capacity from 29 to 191 μmol Trolox/g. The highest values were at 187 °C, using 50% glycerol for 10 min. All models were significant, with R^2 above 0.95. Glycerol, as a green solvent in the microwave-assisted extraction, improves antioxidant compound recovery from juçara residue, presenting itself as a sustainable approach for adding value to the fruit production chain.

Keywords: *Euterpe edulis*; glycerol; microwave; extraction; Atlantic Forest

Citation: Lima, M.; Silva, R.; Novo, A.; Kunigami, C.; Jung, E.; Ribeiro, L. Recovering Antioxidant Compounds from the Juçara Residue by a Green Approach. *Biol. Life Sci. Forum* **2023**, *26*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Name

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Juçara (*Euterpe edulis*) is a native palm tree to the Atlantic Forest, one of the main biomes of Brazil. To obtain its main product, the palm heart, the trees had to be felled, which caused juçara to be placed on the list of Brazilian species threatened with extinction. Alternatively, harvesting the fruits of the juçara palm tree does not involve cutting it down, thus its utilization emerges as an alternative for specie preservation. Furthermore, the use of the fruit should be seen as an opportunity for the sustainable development by means of new technologies and research to promote the preservation of the biodiversity and contribute to the development of activities that fortify local communities [1].

The main product obtained from the fruits is the frozen pulp, a cheap and easy alternative to extend their shelf life. After refinement of the pulp, a residue rich in bioactive compounds and macronutrients is obtained. Studies have already been conducted with this residue, aiming to increase the value of the juçara productive chain. Ribeiro et al. [2] obtained an antioxidant extract from juçara residue using ethanol as a solvent. However, variations in extraction operational conditions and the use of other solvents were not evaluated by these authors. To reduce the environmental impact of the residues and the extraction process, green solvents and cleaner extraction techniques are becoming more important, such as the use of glycerol associated with microwave heating [3].

Glycerol is a byproduct of biodiesel production. With the increasing use of biodiesel as a replacement for fossil fuel-based fuels, glycerol turns into an abundant byproduct in the market. Concerning its characteristics, glycerol exhibits a series of interesting attributes that justify its application as a green solvent, such as being non-flammable, miscible in water, having low volatility, and low cost. However, its use as a solvent in the extraction of bioactive compounds from biomass has been relatively unexplored, in part due to its viscosity. To increase the efficiency of extraction, binary solvent systems can be employed by combining glycerol and water, as well as using high temperatures. However, the main advantage of glycerol that supports its application as a solvent is the non-toxicity, since it is classified by the FDA as GRAS, meaning it is a substance recognized as safe. This feature allows for direct use of extracts in formulations, without requiring purification or fractionation steps [3–6].

The use of microwave as an extraction technique offers advantages such as shorter processing time, automation, and reduced solvent consumption. Furthermore, its application in extraction is efficient due to increasing pressure inside the plant's matrix cells, leading to the rupture of the cell wall, and releasing its contents easier target compounds [7,8].

Therefore, this study aims to evaluate the use of glycerol as green solvent in the microwave-assisted solid-liquid extraction of the antioxidant compounds from juçara residue.

2. Materials and Methods

2.1. Samples

The juçara residue used in this study was obtained from juçara pulp centrifugation performed on the pilot plant of Embrapa Agroindústria de Alimentos (Guaratiba, Rio de Janeiro, Brazil). The residue was defatted and dried before antioxidant recovery. This material exhibited 59.46% of particles with size range between 0.297 mm and 0.590 mm.

2.2. Microwave-Assisted Solid-Liquid Extraction

Glycerol, water, and their combinations were used as a solvent, varying temperature from 53 to 187 °C, and process time, from 2 to 18 min, as shown in Table 1. The combination of independent variables (solvent, temperature, and time) was established through experimental design, using a rotational central composite, comprising 17 experiments. The extracts were subjected to microwave treatment (Ethos 1, Milestone, Brøndby, Denmark) and the solid-liquid proportion was fixed at 1:30 (w/v) for all experiments. After each process, the extracts were vacuum-filtered and stored frozen for subsequent experiments.

Table 1. Experimental design for antioxidant compounds recovery from juçara residue and results obtained for TPC and DPPH*.

Experiment	Glycerol (%)	Temperature (°C)	Time (minute)	TPC (mg GAE/100 g)	DPPH* (µmol Trolox/g)
1	20	80	5	1083	31
2	20	80	15	1052	30
3	20	160	5	3208	90
4	20	160	15	3228	98
5	80	80	5	1151	30
6	80	80	15	1587	40
7	80	160	5	4311	140
8	80	160	15	4222	134
9	0	120	10	1448	41
10	100	120	10	2478	63
11	50	53	10	1096	29

12	50	187	10	5329	191
13	50	120	2	2026	68
14	50	120	18	2447	90
15 (CP)	50	120	10	2711	72
16 (CP)	50	120	10	2388	64
17 (CP)	50	120	10	2286	61

CP—Central Point. GAE—Gallic Acid Equivalent.

2.3. Determination of Total Phenolic Compounds (TPC)

The TPC content was evaluated using the Folin-Ciocalteu reagent, following the method described by Singleton and Rossi [9]. In the method, 250 μL of the extract was mixed with 1.25 mL of 10% (w/v) Folin-Ciocalteu reagent. After 2 min, 2 mL of 7.5% (w/v) sodium carbonate solution was added. The reaction proceeded for 15 min at 50 °C and the tubes were cooled in an ice bath subsequently. The absorbance was measured with a spectrophotometer (SP 220, Biospectro) at 760 nm using ultrapure water as blank. For quantification, a calibration curve was prepared using gallic acid (Sigma-Aldrich, Brazil) as the standard, which ranged from 10 to 100 mg/L. The results were expressed as mg of gallic acid equivalent per 100 g of sample (mg AGE/100 g).

2.4. DPPH• Assay

The antioxidant capacity using the DPPH• radical method was determined as described by Hidalgo [10]. In this method, 100 μL of the extract was mixed with 2.9 mL of DPPH• solution at $6.0 \times 10^{-5}\text{M}$, which was diluted with methanol to obtain a solution with absorbance between 0.600 and 0.800. The reaction took place for 30 min after the addition of the DPPH• radical. The absorbance of the solution was measured with a spectrophotometer at 517 nm, with methanol as the blank. A calibration curve was constructed using Trolox standards and the results were expressed as μM of Trolox per gram (μM Trolox/g).

2.5. Statistical Analysis

The experimental data obtained was evaluated on *Statistica* software version 13 (Dell Inc.), through analysis of variance (ANOVA) and Pareto chart. The test to determine the model's lack of fit and determination of R^2 were employed to verify the model significance, considering a confidence interval of 95%.

3. Results and Discussion

Table 1 presents the results obtained from the conditions proposed by the experimental design. It is possible to observe that the independent variables had an influence on the TPC content of the extracts, ranging from 1052 to 5329 mg GAE/100 g (experiments 2 and 12, respectively). The same can be applied to the antioxidant capacity determined by the DPPH• method, with the best result being 191 μmol Trolox/g, also observed in experiment 12. These data indicated a significant influence caused by the increase in glycerol concentration on the extraction solution, as well as temperature, once experiment 12 employed 50% glycerol as the solvent and a temperature of 187 °C, both higher than the levels used in condition 2. It is important to emphasize that the TPC content and antioxidant capacity by DPPH• observed in this study are superior to those found by Ribeiro et al. [3], which reported a content of 1116 mg GAE/100 g and 39 μmol Trolox/g, respectively, for juçara residue extract obtained with 30% of ethanol at 70 °C for 1 h. This corroborates the efficiency of the extraction system using glycerol as solvent in the microwave-assisted extraction.

The Pareto chart presented in Figures 1 and 2 indicates that the TPC and antioxidant capacity of the extracts were influenced mostly by temperature and glycerol concentration in the extraction solution, with positive linear effects for both independent variables. For the temperature variable, the quadratic effect was also significant, meaning that high

temperatures can reduce the antioxidant potential of the extracts, possibly due to degradation or changes in the structure of the extracted compounds. Also, it is possible to observe that the variable time did not influence the evaluated responses.

The influences of the independent variables observed in this study can be explained through the thermodynamic aspects involved in extraction processes. According to Oliveira [11], increasing the temperature facilitates the extraction of phenolic compounds by increasing the solubility coefficients and the diffusion of compounds to be extracted, while also reducing the viscosity of the solvents. Furthermore, it is known that the polarity of the solvent is crucial for the efficiency of bioactive compound extraction processes. For example, low-polar solvents extract smaller amounts of phenolic compounds, resulting in extracts with lower antioxidant potential [11]. Thus, the glycerol/water mixture provides adequate polarity to the system and reduces the viscosity of pure glycerol, explaining the high values of bioactive compounds and antioxidant capacity in the present study compared to the data reported by Ribeiro et al. [2].

In terms of the statistical analysis, the models were significant in predicting the behavior of the responses concerning the independent variables, as the calculated F-values were higher than the listed F values ($F_{9,7} = 3.68$) at $p = 0.05$. For TPC content and antioxidant capacity using the DPPH• method, the calculated F values were 117 and 38, respectively. Moreover, it is worth mentioning that the lack of fit of the models was not significant, as it presented p -values higher than 0.05 and the calculated F values were lower than the listed F values for all responses. The R^2 values of the adjusted models were 0.98 and 0.95 for the TPC and DPPH•, respectively, indicating that the models explained, at least, 95% of the variability in the data obtained from the experimental design.




Figure 1. Influence of independent variables on the TPC content of the extracts.




Figure 2. Influence of independent variables on the antioxidant capacity by the DPPH* method.

4. Conclusions

Considering the results obtained, it can be concluded that the residue from juçara pulp refinement can be used as a raw material to obtain bioactive compounds. Additionally, the use of glycerol as a solvent in microwave-assisted extraction favored the recovery of these compounds, emerging as a promising approach to add value to the agro-industrial chain of the juçara fruit by a sustainable way.

Acknowledgments: The authors would like to thank Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) for the financial support and to National Institute of Technology (INT).

References

1. Juçara: extração responsável ajuda a manter palmeiras em pé e conserva a Mata Atlântica. #COLABORA. Available online: <https://projetcocolabora.com.br/ods12/jucara-extracao-responsavel-ajuda-a-manter-palmeiras-em-pe-e-conserva-a-mata-atlantica/> (accessed on 1 September 2023).
2. Ribeiro, L.O.; Freitas, S.P. Microencapsulation of the Extract from *Euterpe edulis* Co-product: An Alternative to Add Value to Fruit Agro-Chain. *Waste Biomass Valorization* **2020**, *12*, 1803–1814. <https://doi.org/10.1007/s12649-020-01122-1>
3. Exportações de glicerina têm seu segundo melhor resultado na história. Available online: <https://www.biodieselbr.com/biodiesel/glicerina/exportacoes-de-glicerina-tem-seu-segundo-melhor-resultado-na-historia-130521> (accessed on 1 September 2023).
4. de Los Angeles Perez Fernandez Palha, M.; da Silva, S.P.R. Potencial aproveitamento da glicerina gerada na cadeia produtiva do biodiesel. In Proceedings of the 6° Congresso da Rede Brasileira de Tecnologia de Biodiesel and 9° Congresso Brasileiro de Plantas Oleaginosas, Óleos, Gorduras e Biodiesel, Natal, Brazil, 22–25 November 2016.
5. Huang, H.; Belwal, T. Valorization of lotus byproduct (*Receptaculum Nelumbinis*) under Green extraction condition. *Food Bioprod. Process.* **2019**, *115*, 110–117. <https://doi.org/10.1016/j.seppur.2014.03.010>
6. Tan, H.W.; Abdul Aziz, A.R. Glycerol production and its applications as a raw material: A review. *Renew. Sustain. Energy* **2013**, *27*, 118–117. <https://doi.org/10.1016/j.fbp.2019.03.006>
7. Buratto, R.T.; Cocero, M.J. Characterization of industrial açai pulp residues and valorization by microwave-assisted extraction. *Chem. Eng. Process.-Process Intensif.* **2020**, *160*, 108269. <https://doi.org/10.1016/j.cep.2020.108269>
8. Bener, M.; Ozyürek, M. Optimization of microwave-assisted extraction of curcumin from *Curcuma longa* L. (turmeric) and evaluation of antioxidant activity in multi-test systems. *Rec. of Nat. Prod.* **2016**, *10*, 542.
9. Singleton, V.L.; Rossi, J.A. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enol. Vitic.* **1965**, *16*, 144.

10. Hidalgo, M.; Sánchez-Moreno, C. Flavonoid–flavonoid interaction and its effect on their antioxidant activity. *Food Chem.* **2010**, *121*, 691–696.
11. González-Montelongo, R.; Lobo, M.G. Antioxidant activity in banana peel extracts: Testing extraction conditions and related bioactive compounds. *Food Chem.* **2010**, *119*, 1030–1039.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.