



Proceeding Paper

Production Assay and Partial Characterization of a Protease Produced by *Idiomarina loihiensis*, a Moderately Halophilic Bacterium Strain [†]

Hania Benmebarek * and Karima Kharroub

Extremophile Metabolites Research Team, Biotechnology and Food Quality Research Laboratory (BIOQUAL), Institute of Nutrition, Food and Agro-Technologies (INATAA), Université Frères Mentouri-Constantine 1 (UFMC1), Route de Ain El Bey, Constantine 25000, Algeria; kharroub.karima@umc.edu.dz

* Correspondence: benmebarek.hania@umc.edu.dz

[†] Presented at the 2nd International Electronic Conference on Microbiology, 1–15 December 2023; Available online: <https://ecm2023.sciforum.net>.

Abstract: Halophiles are microorganisms that inhabit saline and hypersaline environments, requiring salinity to survive in such extreme conditions. These microorganisms are mainly researched for their biotechnological potential. The study aims to investigate the phenology of the studied strain, *Idiomarina loihiensis*, and to demonstrate its extracellular proteolytic activity as well as the production of a protease by batch fermentation in halophilic microorganisms. Macroscopic studies revealed small colonies (≤ 5 mm) with a convex spherical structure, regular outline, smooth surface and color ranging from beige to opaque cream. Protease production was investigated in high salinity conditions by a moderately halophilic bacterium using basal media with varying nitrogen sources. The study found that the highest proteolytic activity occurred in media with tryptone and casein peptone as nitrogen sources at pH 10, temperature 70 °C, and 22.5% salt concentration. The results also demonstrated that the protease was a thermostable enzyme.

Keywords: halophile; protease; *Idiomarina loihiensis*; enzyme; proteolytic activity

1. Introduction

Extremophile microorganisms dwelling in harsh environmental habitats have evolved unique properties; which can be of biotechnological and commercial significance. These non-conventional microorganisms living in such extremely hostile environmental habitats possess special adaptation strategies that make them interesting [1]. And have developed the capacity to produce extremophile enzymes such as proteases with biotechnological potential [2].

Besides being intrinsically stable and active at high salt concentrations, halophilic enzymes offer important opportunities in biotechnological applications, such as food processing, environmental bioremediation and biosynthetic process [3].

It is important to highlight that the use of enzymes from halophiles in industrial applications are not limited to their stability at high salt concentrations, since these extremozymes usually are also tolerant to high temperatures and they are stable in presence of organic solvents [2,3].

Halophilic microbes isolated from different saline soils have been found to produce proteases with potential industrial biotechnological value [4]. They are most significant as their properties can be easily modified through genetic manipulations to suit their various applications [5].

The industrial demand of proteases with special properties continues to stimulating the search for new enzymes. The genus *Bacillus* is the main producer of industrial alkaline

Citation: Benmebarek, H.; Kharroub, K. Production Assay and Partial Characterization of a Protease Produced by *Idiomarina loihiensis*, a Moderately Halophilic Bacterium Strain. *Biol. Life Sci. Forum* **2023**, *31*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Name

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

proteases, however other genus such as *Pseudomonas* and *Streptomyces* isolated from various environments have been reported as producers of alkaline and thermostable proteases with industrial importance. Likewise, *Idiomarina* could represent an important source of novel proteases with commercial applications [2].

The aim of this study was the characterisation of the phenology and the demonstration of the extracellular proteolytic activity of the studied strain *Idiomarina loihiensis* and the production of a protease by batch fermentation in halophilic microorganisms.

2. Material and Methods

2.1. Samples and Strain Isolation and Identification

A moderately halophilic bacterium was isolated from an evaporation table in Sebkheth Ez-Zemoul (Ain M'lila, eastern Algeria ♀ 35.874076, 6.544489) and its proteolytic potential was confirmed on agar with 1% casein/gelatin added (MGM solid medium-0.3% peptone and 0.1% yeast extract with 12.5% salt) and by an enzymatic test on liquid medium in the presence of azocasein [6]. Molecular identification was carried out using the 16S rRNA gene [7].

2.2. Morphology and Physiology

The strain was examined for morphology using cells from exponentially growing cultures. The aspect of the colonies was examined. Cell morphology and Gram staining were observed. Proteolytic activity was tested on MGM (modified growth medium) solid media (0.3% peptone and 0.1% yeast extract) supplemented with 1% (*w/v*) casein [8] or 1% (*w/v*) gelatin [9]; positive results were detected after 5 days at 37 °C by the presence of a precipitate around the colonies for casein and/or a translucent halo (after addition of Frazier's reagent) for gelatin. Growth rates were estimated on solid MGM medium at different salt concentrations (0–27%), pH (5–10) incubated at 37 °C and temperatures (12–55 °C) under aerobic conditions for 24–72 h.

2.3. Inoculum Preparation and Halophilic Proteases Production

The inoculum is prepared from a 24-h culture on Petri dishes, inoculated into tubes containing 5 mL MGM medium with 10% salt and incubated at 37 °C for 48 h. Enzyme extract was prepared by inoculating 10% *v/v* of bacterial suspension into 250 mL flasks containing 45 mL production medium (MGM 0.05% peptone and 0.01% yeast extract) supplemented with different nitrogen sources: glycine, peptone, gelatin, casein, casein-peptone, tryptone, cream, gelatin- tryptone. The pH, salt concentration and temperature are adjusted according to the optimal growth conditions for the culture [10];

The media are incubated in a shaking bath at 37 °C for 06 days. The cultures are then centrifuged (4000× *g* for 10 min) and the supernatant is cooled to 0 °C. To compare proteolytic activity on the different nitrogen sources, a two-factor analysis of variance was performed using Minitab 18 Statistical Software.

2.4. Enzymatique Assay

Enzymatic assay was performed using two methods to confirm proteolytic activity at 0% and 10% of salt. A positive control with proteinase K and a blank assay (addition of TCA prior to reaction) were performed each time.

2.4.1. Method 1

Proteolytic activity was determined by the method of Tsuchida et al. (1986). The reaction mixture contained 1 mL of 2% casein (prepared in phosphate buffer at pH 7) and 1 mL of the enzyme extract. The mixture was incubated in a water bath for 30 min at 40 °C (standard conditions). The enzymatic reaction was stopped by the addition of 2 mL of 0.4 M TCA and centrifuged at 4000× *g* for 10 min. Then 1 mL of the supernatant was transferred to another tube to which 5 mL of 0.4 M Na₂CO₃ and 1 mL of 10% *v/v* Folin's calcium

reagent were added. The reaction was incubated for 20 min at 40 °C in the dark. The absorbance was measured at a wavelength of 750 nm. One unit of proteolytic activity is defined as the amount of enzyme capable of releasing 1 µg of tyrosine per ml/min [11].

2.4.2. Method 2

Halophilic protease activity was determined according to the method of Brock et al. 1982 modified by Chuprom et al. 2016, using azocasein (sulfanilamide azocasein) as substrate. 1 mL of a reaction mixture consisting of 1/1 crude enzyme (cell-free supernatant) and 0.8% (*w/v*) azocasein (in 0.1 M Tris-HCl buffer (pH 8.0) at 10% (*w/v*) salinity) was incubated at 40 °C for 2 h in a Marie bath. The reaction was stopped by adding 1 mL of 10% (*w/v*) TCA solution and the mixture was allowed to stand for 30 min at room temperature. After centrifugation at 4000× *g* for 10 min, 600 µL of the supernatant was transferred to a microcentrifuge tube and 700 µL of 1.0 N NaOH was added. Absorbance was measured at 440 nm using a UV-vis spectrophotometer (one unit (U) of halophilic protease activity was defined as the amount of enzymatic activity that produced a 0.01 change in absorbance at 440 nm in 2 h at 40 C under standard assay conditions). Halophilic protease activity was calculated using the following equation:

$$\text{Halophilic protease activity (U/}\mu\text{L)} = \frac{(A-B) \cdot Vt}{Ve \cdot 0.01}$$

where *A* and *B* are the optical densities of the crude enzyme and the control, respectively, *Vt* is the total reaction volume and *Ve* is the volume of crude enzyme [10].

2.5. Partial Physico-Chemical Characterisation of the Enzyme Extract

In order to determine the parameters required for optimum proteolytic activity, the enzymatic reaction is carried out in different buffer solutions from pH 4 to 12, at different temperatures (30, 40, 50, 60, 70, 80 and 90 °C) for 30 min, in the presence of 0–30% salts and, finally, to study thermostability, the reaction mixture is heated to 70 °C for 4 periods (30, 60, 90 and 120 min) and the residual activity is compared with the optimum activity. Proteolytic activity was determined in duplicate by the Tsuchida method [12].

3. Results and Discussion

A moderately halophilic bacterium strain producing a halophilic extracellular protease was isolated from a hypersaline environment. The bacteria hydrolyzed casein and gelatin, demonstrating their proteolytic potential. The halophilic strain was identified as *Idiomarina loihiensis*.

3.1. Strain Isolation and Identification

The strain studied showed small colonies (≤5 mm) of domed spherical shape, regular outline, smooth surface, beige to opaque cream in colour. Gram staining using the technique of (Dussault, 1955) [13] showed that the strain was Gram-negative small rods with cells arranged in chains, pairs or dispersed, as is the case with most isolates of moderately halophilic, halotolerant and halophilic bacteria isolated from different saline habitats [14–16]. The halophilic bacterium was identified as *Idiomarina loihiensis* by molecular characterization.

3.2. Nitrogen Source

Different nitrogen sources affecting the production of halophilic protease by the isolate give differences in activity depending on the dosing method and the salinity of the medium, as shown in Figure 1. In the absence of salt, the most cost-effective nitrogen sources are peptone for the azocasein method and casein peptone for the Tsuchida method, yielding higher activity as positive control. However, in the presence of 10% (*w/v*) salt, the best nitrogen source for both methods is tryptone, which results in greater

proteolysis compared to proteinase K (control). Analysis of variance confirms that tryptone is the best carbon source.




Figure 1. Proteolytic activity of the enzymatic curd from *Idiomarina loihiensis* against various nitrogen sources. (a) at 0% salt and (b) at 10% salt concentration.

3.3. Partial Physico-Chemical Characterization

A characteristic curve of the pH effect on enzymatic activity is shown in Figure 2a. Maximum activity is observed at pH 10, before and after this value, it decreases until reaching a minimum at pH 7, leading to the conclusion that our extract contains alkaline proteases. Similar results were obtained with proteases produced by the moderate halophilic bacterium *Halobacillus karajensis*, whose optimum pH is 9 [17], other proteolytic enzymes of *Bacillus luteus* H11 have a pH optimum at 10.5 [18]. Other similar results are obtained with proteases produced by *Thermoactinomyces* sp. HS682, whose optimum pH is 11 [19].




Figure 2. Physiological effects on pH (a) and salinity (b) on the proteolytic activity of enzyme curd.

Figure 2b shows a characteristic curve of the effect of salts on enzymatic activity. Indeed, the maximum activity is observed at 22.5% salt with a maximum activity of 2.55 U/mL/min. Similar results were obtained for the halophilic bacteria *Halobacillus* sp. SR5-3, which showed optimum activity at 20% salt. The activity of the enzyme was increased approximately 2.5-fold by the addition of 20 to 35% salt, and the enzyme was strongly stabilised by the salt.

For the temperature effect, it can be seen that the proteolytic activity increases at a temperature of 70 °C, but decreases sharply above this temperature (Figure 3a). We can therefore conclude that the optimum temperature at which the enzyme extract shows maximum proteolytic activity is 70 °C. Results similar to ours have been obtained for the strain *Chromohalobacter* sp. TVSP101, which shows a protease with maximum activity at 75 °C [20]. The strain *Thermoactinomyces* sp. HS682 studied by Tsuchida et al. (1992) has an active protease at 70 °C.




Figure 3. Temperature effect (a) on proteolytic activity and thermal stability (b) of the enzyme curd.

Thermostability of the enzyme extract at the optimum temperature showed maximum activity at 30 min as shown in Figure 3b, a 33% loss of activity at 60 min, which remained stable until 90 min and increased to 86% after 120 min. The results obtained reveal that the protease from *Idiomarina loihiensis* strain is a thermostable enzyme. There are thermostable bacterial enzymes such as thermolysin, a protease produced by *Bacillus thermoproteolyticus*, which retains half of its activity after one hour at 80 °C [21].

Table 1. Variance analysis (Turkey test). Tukey's method and 95% confidence level for grouping information.

Midium	N	Average	Group
Tryptone	8	0.825750	A *
Cesein peptone	8	0.797250	A
Gelatine + Tryptone	8	0.759250	A
Skim. milk	8	0.755625	A
Peptone	8	0.589125	A B *
Gelatine	8	0.379750	B
Caseine	8	0.353375	B
Glycine	8	0.335375	B

* Groups. Averages with no common letters are significantly different.

4. Conclusions

The moderately halophilic strain *Idiomarina loihiensis* secretes a proteolytic enzyme extract with maximum activity at salinity 22.5%, pH 10 and temperature 70 °C, maintaining more than 80% of the proteolytic activity after 120 min at this temperature.

This shows that we are dealing with a halophilic, alkaliphilic and thermostable protease capable of maintaining a good hydrolytic activity under extreme conditions, which makes it very interesting from a biotechnological point of view.

Author Contributions:

Funding:

Institutional Review Board Statement:

Informed Consent Statement:

Data Availability Statement:

Acknowledgments: We kindly thank Guessoum Randa and Boumezouzd Fatima Zahra for their contribution.

Conflicts of Interest:

References

1. Jothi Basu, M.; Indra, V.; Hemapriya, J.; Vijayanand, S. Bioprocess Optimization of Halophilic Protease Production by a Halophilic Bacterial Strain JS4. *Int. J. Curr. Res. Acad. Rev.* **2015**, *3*, 309–315. Available online: www.ijcrar.com (accessed on 26 June 2019).
2. Flores-Fernández, C.N.; Chávez-Hidalgo, E.; Santos, M.; Zavaleta, A.I.; Arahal, D.R. Molecular characterization of protease producing *Idiomarina* species isolated from Peruvian saline environments. *Microbiol. Biotechnol. Lett.* **2019**, *47*, 4001–4411. <https://doi.org/10.4014/mbl.1809.09001>.
3. De Lourdes Moreno, M.; Pérez, D.; García, M.; Mellado, E. Halophilic bacteria as a source of novel hydrolytic enzymes. *Life* **2013**, *3*, 38–51. <https://doi.org/10.3390/life3010038>.
4. Corral, J.C.C.; De Los Santos Villalobos, S.; Barragán, L.A.P.; Figueroa, J.J.B.; Vásquez-Murrieta, M.S.; Alvarado, M.I.E.; Chávez, L.A.C. Isolation of moderately halophilic bacteria in saline environments of sonora state searching for proteolytic hydrolases. *Open Agric.* **2018**, *3*, 207–213. <https://doi.org/10.1515/opag-2018-0021>.
5. Asha, B.; Palaniswamy, M. Optimization of alkaline protease production by *Bacillus cereus* FT 1 isolated from soil. *J. Appl. Pharm. Sci.* **2018**, *8*, 119–127. <https://doi.org/10.7324/JAPS.2018.8219>.
6. Gomri, M.A.; Rico-Díaz, A.; Escuder-Rodríguez, J.-J.; Khaldi, T.E.M.; González-Siso, M.-I.; Kharroub, K. Production and Characterization of an Extracellular Acid Protease from Thermophilic *Brevibacillus* sp. OA30 Isolated from an Algerian Hot Spring. *Microorganisms* **2018**, *6*, 31. <https://doi.org/10.3390/microorganisms6020031>.
7. Benmebarek, H.; Escuder-Rodríguez, J.J.; González-Siso, M.I.; Kharroub, K. Test for the production and assay of the proteolytic activities of halophilic bacteria and archaea isolated from algerian hypersaline environments. *Proceedings* **2020**, *66*, 12. <https://doi.org/10.3390/proceedings2020066012>.
8. Cojoc, R.; Merciu, S.; Popescu, G.; Dumitru, L.; Enache, M.; Kamekura, M.; Enache, M. Extracellular hydrolytic enzymes of halophilic bacteria isolated from a subterranean rock salt crystal. *Rom. Biotechnol. Lett.* **2009**, *14*, 4658–4664. Available online: https://www.researchgate.net/publication/228413520_Extracellular_hydrolytic_enzymes_of_halophilic_bacteria_isolated_from_a_subterranean_rock_salt_crystal (accessed on 12 October 2023).
9. Gutiérrez, C.; González, C. Method for simultaneous detection of proteinase and esterase activities in extremely halophilic bacteria. *Appl. Microbiol.* **1972**, *24*, 516–517. <https://doi.org/10.1128/am.24.3.516-517.1972>.
10. Chuprom, J.; Bovornreungroj, P.; Ahmad, M.; Kantachote, D.; Dueramae, S. Approach toward enhancement of halophilic protease production by *Halobacterium* sp. strain LBU50301 using statistical design response surface methodology. *Biotechnol. Rep. (Amst.)* **2016**, *10*, 17–28. <https://doi.org/10.1016/j.btre.2016.02.004>.
11. Tsuchida, O.; Yamagata, Y.; Ishizuka, T.; Arai, T.; Yamada, J.-I.; Takeuchi, M.; Eiji, I. An Alkaline Proteinase of an alkalophilic *Bacillus* sp. *Curr. Microbiol.* **1986**, *14*, 7–12. <https://doi.org/10.1007/BF01568094>.
12. Boughachiche, F.; Rachedi, K.; Duran, R.; Lauga, B.; Karama, S.; Bouyoucef, L.; Boulezaz, S.; Boukrouma, M.; Boutaleb, H.; Boulahrouf, A.. Optimization of alkaline protease production by *Streptomyces* sp. strain isolated from saltpan environment. *Afr. J. Biotechnol.* **2016**, *15*, 1401–1412. <https://doi.org/10.5897/AJB2016.15259>.
13. Dussault, H.P. An improved technique for staining red halophilic bacteria. *J. Bacteriol.* **1955**, *70*, 484–485. <https://doi.org/10.1128/jb.70.4.484-485.1955>.
14. Ventosa, A.; De La Haba, R.R. Chromohalobacter. In *Bergey's Manual of Systematics of Archaea and Bacteria*, 1st ed.; Whitman, W.B., Ed.; Wiley: Hoboken, NJ, USA, 2020; pp. 1–16. <https://doi.org/10.1002/9781118960608.gbm01189.pub2>.
15. Arahal, D.R.; Ludwig, W.; Schleifer, K.H.; Ventosa, A. Phylogeny of the family *Halomonadaceae* based on 23S and 16S rDNA sequence analyses. *IJSEM* **2002**, *52*, 241–249. <https://doi.org/10.1099/00207713-52-1-241>.
16. Yoon, S.; Govind, C.K.; Qiu, H.; Kim, S.J.; Dong, J.; Hinnebusch, A.G. Recruitment of the ArgR/Mcm1p repressor is stimulated by the activator Gcn4p: A self-checking activation mechanism. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 11713–11718. <https://doi.org/10.1073/pnas.0404652101>.
17. Karbalaie-Heidari, H.R.; Amoozegar, M.A.; Hajighasemi, M.; Ziaee, A.A.; Ventosa, A. Production, optimization and purification of a novel extracellular protease from the moderately halophilic bacterium *Halobacillus karajensis*. *J. Ind. Microbiol. Biotechnol.* **2009**, *36*, 21–27. <https://doi.org/10.1007/s10295-008-0466-y>.
18. Kalwasińska, A.; Jankiewicz, U.; Felföldi, T.; Burkowska-But, A.; Swiontek Brzezinska, M. Alkaline and halophilic protease production by *Bacillus luteus* h11 and its potential industrial applications. *Food Technol. Biotechnol.* **2018**, *56*, 553–561. <https://doi.org/10.17113/ftb.56.04.18.5553>.
19. Tsuchiya, K.; Nakamura, Y.; Sakashita, H.; Kimura, T. Purification and characterization of a thermostable alkaline protease from alkalophilic *Thermoactinomyces* sp. H8682. *Biosci. Biotechnol. Biochem.* **1992**, *56*, 246–250. <https://doi.org/10.1271/bbb.56.246>.
20. Vidyasagar, M.; Prakash, S.; Mahajan, V.; Shouche, Y.S.; Sreeramulu, K. Purification and characterization of an extreme halothermophilic protease from a halophilic bacterium *Chromohalobacter* sp. TVSP101. *Braz. J. Microbiol.* **2009**, *40*, 12–19. <https://doi.org/10.1590/S1517-83822009000100002>.
21. Majumder, D.R.; Kanekar, P.P.; Gaikwad, S.M. Purification and characterization of a thermolysin like protease from *Thermoactinomyces thalophilus* MCMB-380. *Protein Pept. Lett.* **2013**, *20*, 918–925. <https://doi.org/10.2174/0929866511320080009>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.