

PGPR characteristics of rhizospheric bacteria to understand the mechanisms of faba bean growth

Rim MAOUGAL^{1,2}, Maya KECHID^{1,2}, Chaima LADJABI³ and Abdelhamid DJEKOUN¹

¹ Laboratoire Génétique Biochimie et Biotechnologie Végétale, SNV Université frères Mentouri Constantine 1

² Institut de la Nutrition, de l'Alimentation et des Technologies Agro-alimentaires (INATAA), Département de Biotechnologie alimentaire. Université Frères Mentouri, Constantine 1, Algérie.

³ Départements d'écologie et biologie végétale, SNV Université frères Mentouri Constantine 1 Constantine, Algérie.

rym.maougal@umc.edu.dz; maya.kechid@umc.edu.dz; ladjabichaima94@hotmail.com; djak2591@gmail.com*

* Correspondence: e-mail@e-mail.com; Tel.: (optional; include country code; if there are multiple corresponding authors, add author initials) +xx-xxxx-xxx-xxxx (F.L.)

† Presented at the 1st International Electronic Conference on Microbiology, 2–30 November 2020; Available online: <https://ecm2020.sciforum.net/>

Published: 2 November 2020

Abstract: Rhizobacteria play an important role in maintaining soil balance, among these bacteria there are those which have shown their ability to promote the growth of plants known as Plant Growth Promoting Rhizobacteria (PGPR). In our work, we are interested in characterizing 110 bacterial strains isolated in the field in the region of ben badis (Constantine Algeria) from 5 varieties of faba bean. Phenotypic and biochemical characterization showed that most of the isolates are cream colored, slightly raised, flat and opaque, gram-, catalase + and oxidase-, Bacillus form. PCA analysis allowed us to select 40 isolates with a high degree of variability to continue our work. The results obtained have directed us towards different taxonomic groups (rhizobium, Pseudomonas, bacillus etc.). The evaluation of the PGPR potential of bacteria (phytostimulation, biofertilization and biocontrol), showed that 100 % of bacteria are able to produce auxin at different concentrations with the highest concentration (177.77 µg / ml) for the isolate 6, that more than 50% of isolates are capable of producing nitrogen, ammonia and phytate mineralization. These PGPR traits are a direct effect on plant growth of five varieties of the faba bean and can be used to select the best performing bacteria for inoculation tests.

Keywords: Growth promotion; faba bean; rhizosphere; PGPR traits

1. Introduction:

The soil is an important reservoir for microorganisms, it contains a complex and varied microflora that plays essential roles for the telluric ecosystem and the multi-cellular organisms that live there. Among the higher soil organisms, plants benefit from the direct and / or indirect effects of microorganisms, in particular those of the rhizospheric flora. (Munees and Mulugeta, 2013).

PGPR bacteria (Plant Growth Promoting Rhizobacteria) are part of this type of microorganism because they do not establish a mutualistic symbiosis relationship, but promote the growth of plants with which they are associated (Kloepper and Beauchamp, 1992). The selection of an effective PGPR strain is linked to the characterization of its properties promoting plant growth such as: the solubilization of phosphate, the production of siderophores and phytohormones, the ability to fix nitrogen, and the biological control of diseases plants (Cattelan et al., 1999).

The objective of this work is to pre-identify bacteria previously isolated from the rhizosphere of faba bean, our plant model in this study, based on their morphological characters: macroscopic, microscopic, biochemical and physiological.

2. Methods

Strains of bacteria were isolated from rhizosphere of faba bean, roots are washed and soil solution obtained was used for isolation of non-symbiotic bacteria. Physiological and biochemical characteristics of the bacterial isolates were examined according to the methods described by the Bergey's manual (Holt et al. 1994). Different assays and tests allowed to test the PGPR functions.

3. Results and discussion

The phenotypic characterization of the 110 rhizobacteria showed that most isolates are Gram - (94.54%), catalase + (87.5%), oxidase - (65 %), slightly raised (27.27%), flat (61.81%), and opaque (50.9%), round in shape (60%), and in different colors, Bacillus form (60 %).

The detection of pyoverdinin pigment by observation of a yellow-green color show that 55.45 % of the bacteria exhibited a yellow-green fluorescence against 43.63 % without fluorescence This fluorescence could be indicative of the affiliation of the bacteria to the species *Pseudomonas fluorescens*.

The factorial analysis of these tests allowed us to realize a point cloud gathering the isolates with a certain degree of similarity. We used the PCA to select the 40 isolates with a high degree of variability to continue our work.

The study of the Plant Growth Promoting Rhizobacteria (PGPR) effect of rhizobacteria showed that 100% of the isolates tested are able of producing auxin at 177.77 µg/ml concentration (Strain 6).

The solubilization efficiency of phosphates from our isolates measured on NBRIP medium (Kumar V et al., 2013) showed that 50% of isolates were able to grow on a medium containing phytate as the sole source of phosphorus so they are able to degrade phytate.

The growth of bacteria on "Burk's N-free" medium depends on their ability to fix nitrogen our results showed that 72.5% of the isolates tested were able to fix nitrogen.

The appearance of the medium after incubation at 30 ° C and addition of Nessler's reagent showed that 67.5% of the media appear yellowish or orange in color, which corresponds ammonium-producing bacteria

4. Conclusion

Through this present work and the experiments carried out, we have affirmed the presence of a relationship between the bacteria of the rhizosphere of the plant *vicia faba* in the region of Constantine (Algeria) thanks to their important agro-economic, environmental and ecological interests.

Better knowledge of the mechanisms of plant-microorganism interaction makes it possible to promote populations already present in the rhizosphere and which are beneficial for plants. So its isolates can therefore be further studied and developed, in order to be used in agriculture and biotechnology.

References

1. Cattelan A. J., Hartel P. G., Fuhrmann J. J. (1999). Screening for plant growth-promoting rhizobacteria to promote early soybean growth. *Soil Sci. Soc. Am. J.* 63 1670–1680.
2. Holt et al., J.G. Holt, N.R. Krieg, P.H.A. Sneath, J.T. Stanley, S.T. Williams. (1994) *Bergey's Manual of Determinative Bacteriology* (9th ed.), Williams & Wilkins, Co., Baltimore
3. Kloepper, J. w., Beauchamp, C. J. (1992): A review of issues related to measuring colonization of plant roots by bacteria. *Can. J. Microbiol.* 38, 1219 -1232
4. Kumar ,A., A. Kumar, S. Devi, S. Patil, C. Payal, S. Negi (2012), Isolation, screening and characterization of bacteria from rhizospheric soils for different plant growth promotion (PGP) activities: an in vitro study *Recent Res. Sci. Technol.*, 4 (1) pp. 01-05
5. Munees A. ,Mulugeta K. (2013) .Mechanisms and applications of plant growth promoting rhizobacteria: Current perspective, *Journal of King Saud University – Science*, January Volume 26, Issue 1, Page 1–20

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).