

Proceedings

Synthesis and Structure of Novel Potentially Bioactive Amphiphilic -O-(N)-Glycosides [†]

Natividad Bejarano, Leticia Lafuente, Juliana Esteche, Cintia Santiago, Agustín Rojas and Agustín Ponzinibbio ^{*}

Centro de Estudio de Compuestos Orgánicos (CEDECOR), Departamento de Química, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, 47 y 115, 1900 La Plata, Argentina

* Correspondence: ponzinibbio@quimica.unlp.edu.ar

† Presented at the 24th International Electronic Conference on Synthetic Organic Chemistry, 15 November–15 December 2020; Available online: <https://ecsoc-24.sciforum.net/>.

Received: date; Accepted: date; Published: date

Abstract: Herein we present the synthesis and structural analysis of novel -O-(N)-glycosides. The biphasic reaction of NHS and acetobromo- α -D-glucose or D-galactose gives the β anomer glycoside in a straightforward manner. Further hydrazinolysis and condensation with decanal afforded the desired products. Their complete structures, including the anomeric and E/Z double bond configurations, were determined by spectroscopic analysis.

Keywords: amphiphilic carbohydrates; -O-(N)-glycosides; NMR

1. Introduction

Carbohydrate-based surfactants are molecules with a sugar hydrophilic moiety and a hydrophobic tail, usually a hydrocarbon chain such as a fatty acid or alcohol. These glycosides have been synthesized in the last few decades and their structures were analyzed in terms of supramolecular arrangements as described in the literature [1]. Furthermore, they were proposed as potential therapeutic active molecules. This bioactive profile lays on their capability to interact with the lipid bilayer of biological membranes due to the lipophilicity exhibited by the glycoside's aglycone. [2]. In this work, we show the design and synthesis of carbohydrate derivatives with amphiphilic properties including a bioactive moiety, a Schiff base. Oxime ether function is an important structural motif in many famous drug scaffolds. Many anti-viral agents have the -C=N-O- function in their structures, for example, the kinase inhibitor “enviroxime” and the recently reported “EIDD-2801” glycoside which has shown broad anti-influenza virus activity [3].

2. Materials and Methods

2.1. General

Commercially available reagents were directly used without further purification. All sugar derivatives precursors were prepared by known methods, carefully purified, and fully characterized by ¹H and ¹³C NMR. Reactions were monitored by TLC on silica gel (Merck 60 F254 plates), with detection by UV light (254 nm) or by charring with sulfuric acid. Flash chromatography was performed using silica gel (230–400 mesh). ¹H and ¹³C NMR spectra were recorded on a Bruker AVIII (600 and 150 MHz, respectively) in CDCl₃. HSQC and COSY spectra were used to establish peak assignments in ¹H and ¹³C NMR.

2.2. N-Hydroxysuccinimide Glycosylation

To a stirred mixture of acetobromo- α -D-glucose (1 mmol) and Na_2CO_3 solution in CH_2Cl_2 (10 mL, 1M) was added N-hydroxysuccinimide (3 mmol) followed by TBAHS (1 mmol) phase transfer catalyst. The mixture was stirred at 25 °C until the reaction was completed as indicated by TLC. Then CH_2Cl_2 (10 mL) was added and washed twice with 10 mL of water and brine. The organic layer was dried and concentrated in vacuo. The corresponding product was purified by flash column chromatography to obtain the pure glycoside derivative.

2.3. Hydrazinolysis Procedure

To a stirred solution of the aforementioned compound (0.5 mmol) in 10 mL of 1:1 CH_2Cl_2 -MeOH mixture was added hydrazine hydrate (1.5 mmol) at room temperature. The reaction was monitored by TLC, and after 30 min quenched with 5 mL of saturated NaHCO_3 solution. The mixture was extracted with 3 × 10 mL of CH_2Cl_2 . The organic layers were washed with brine, dried, and concentrated in vacuo. Products **1** (D-glucose derivative) and **2** (D-galactose derivative) were carefully purified by column chromatography to obtain the corresponding -O-NH₂ β-glycoside.

2.4. Synthesis of Oxime Derivatives

The aminoxyglycoside (1 mmol) and 1.2 eq of decanal were dissolved in dried CH_2Cl_2 (2 mL) then one drop of concentrated HCl was added. The mixture was stirred for 24 h at ambient temperature until the reaction was completed as indicated by TLC, then quenched with water (3 mL) and the aqueous phase was extracted with CH_2Cl_2 (3 × 10 mL), neutralized with NaHCO_3 solution and washed twice with water. The combined organic layers were dried and concentrated under reduced pressure. The residue was dried under high vacuum. The purification was performed by flash column chromatography (hexane/EtOAc 8:2) to afford the desired compounds **3** and **4**.

3. Results and Discussion

The biphasic reaction between N-hydroxysuccinimide and acetobromo- α -D-glucose or D-galactose gives the β-O-(N)-glycoside in a straightforward manner. Further hydrazinolysis and condensation with decanal, under acid catalysis, give the desired products. (Figure 1)




Figure 1. Multistep synthesis of amphiphilic glycosides Schiff bases from acetobromo- α -D-glucose and D-galactose.

In all cases, we obtained the products in good yields and high anomeric stereoselectivity. The values of $^3\text{J}_{\text{H}1-\text{H}2}$ coupling constants were measured and all of them are around 8–9 Hz. This, undoubtedly indicates the beta configuration at the anomeric centre. Their structures were determined and fully characterized by ^1H and ^{13}C NMR spectroscopy and were as expected. The oxime ether glycosidic linkage is a Schiff base moiety. The *E* isomer is the thermodynamically more

stable and consequently the major isomer obtained. ^1H NMR spectra of non-purified products reveal the presence of a minor by-product. An isomeric compound was detected in a 3:2 ratio in both, D-glucose and D-galactose derivatives. Particularly, the oximyl proton could be clearly differentiated, we present δ values obtained in Table 1.

Table 1. Oximyl proton displacements for compounds **3** and **4**, the values were extracted from ^1H NMR spectra of isomeric mixtures.

Product	$\delta 1$ (ppm)	$\delta 2$ (ppm)
3	6.81	7.51
4	6.80	7.53

The complete ^1H and ^{13}C NMR experiments demonstrate that this minor product was the Z isomer. Assignment of the E/Z configuration may be done from the ^1H NMR spectra of oxime protons. It is well reported that oximyl protons of (*E*)-oxime ethers are deshielded relatively to those of (*Z*)-oxime ethers [4].

After flash column chromatography with 15–25 μm silica gel, it was possible to separate both isomers. *E* and Z products were obtained pure and the NMR analysis confirms ours mentioned hypothesis. Comparing the data extracted from ^1H NMR spectra, with those reported previously by us and others we may consider that the *E* configuration isomer was the major product obtained [5]. (Figure 2).




Figure 2. Partial ^1H NMR spectra for both *E* and *Z* isomers of **3a** and **3b**.

With the method optimized and, after Zemplén deacetylation, we synthesized four new -O-(N)-glycosides bearing a hydrophobic tail at the anomeric position as seen in Figure 3 [6].




Figure 3. New amphiphilic -O-(N)- β -glycosides. Overall yields in brackets.

The glycosides synthesized were particularly stable. As previously described in literature [7], the stability towards hydrolysis of oximes is higher than simple hydrazones because of the

electronegativity differences in the heteroatom ($\chi_O = 4.5$ vs. $\chi_N = 3.0$). No acetal or oxime hydrolysis products were detected after several days in aqueous solution. Moreover, the protonation of the anomeric oxygen, to catalyze the acetal hydrolysis in an exocyclic cleavage, is also not favored. If an alternative endocyclic cleavage mechanism may be performed, the rate-limiting step involves delocalization of the electrons on oxygen over the glycosidic bond. This delocalization is completely disfavored [8].

4. Conclusions

A simple and efficient route for the synthesis of novel amphiphilic carbohydrate-derived oxime ethers has been described. The products were obtained in good overall yields and with high anomeric and *E/Z* stereocontrol. Both *E* and *Z* isomers were obtained pure after a careful chromatography work and a complete structural analysis were done compiling interesting data that may be used for setting future correlations. We recently initiated investigations to evaluate surfactant properties and cytotoxic effects of the novel compounds prepared.

Funding: This research received no external funding.

References

1. a) Allen, D.K.; Tao, B.Y Carbohydrate-alkyl ester derivatives as biosurfactants. *J. Surfactants Deterg.* **1999**, *2*, 383–390. b) Söderman, O.; Johansson, J. Polyhydroxyl-based surfactants and their physico-chemical properties and applications. *Curr. Opin. Colloid Interface Sci.* **1999**, *4*, 391–401. c) Ewan, H.S.; Muli, C.S.; Touba, S.; Bellinghiere, A.T.; Veitschegger, A.M.; Smith, T.B.; Pistel, W.L.; Jewell, W.T.; Rowe, R.K.; Hagen, J.P.; et al. Synthesis of sugar oxime ether surfactants. *Tetrahedron Lett.* **2014**, *55*, 4962–4965.
2. Xavier, N.M.; Rauter, A.P. Environmentally friendly approaches to the synthesis of new antibiotics from sugars. *Pure Appl. Chem.* **2012**, *84*, 803–816 and references cited therein.
3. a) Heinz, B.A.; Vance, L.M. The antiviral compound enviroxime targets the 3A coding region of rhinovirus and poliovirus. *J. Virol.* **1995**, *69*, 4189–4197. b) Toots, M.; Yoon, J.; Cox, R.M.; Hart, M.; Sticher, M.Z.; Makhsous, N.; Plesker, R.; Barrena, A.H.; Reddy, P.G.; Mitchell, D.G.; et al. Characterization of orally efficacious influenza drug with high resistance barrier in ferrets and human airway epithelia. *Sci. Transl. Med.* **2019**, *11*, eaax5866.
4. a) Karabatsos, G.J.; His, N. Structural studies by nuclear magnetic resonance—XI: Conformations and configurations of oxime o-methyl ethers. *Tetrahedron* **1967**, *23*, 1079–1095. b) Rodriguez, E.C.; Winans, K.A.; King, D.S.; Bertozzi, C.R. A Strategy for the Chemoselective Synthesis of O-Linked Glycopeptides with Native Sugar-Peptide Linkages. *J. Am. Chem. Soc.* **1997**, *119*, 9905–9906. c) Winans, K.A.; Bertozzi, C.R. An inhibitor of the human UDP-GlcNAc 4-epimerase identified from a uridine-based library: A strategy to inhibit O-linked glycosylation. *Chem. Biol.* **2002**, *9*, 113–129.
5. a) Lafuente, L.; Santiago, C.C.; Rojas, A.H.; Piro, O.E.; Echeverría, G.A.; Ponzinibbio, A. Selective Synthesis and Molecular Structure of Novel Aminoxyglycosyl Derivatives Bearing Hydroxyphenyl Moieties. *ChemistrySelect* **2020**, *5*, 864–868. b) Lafuente, L.; Rojas, A.H.; Piro, O.E.; Echeverría, G.A.; Ponzinibbio, A.; Synthesis, NMR and X-ray studies on novel heteroaromatic aldoxime O-ether 2-and 2, 3-unsaturated glycosides. *Tetrahedron Lett.* **2020**, *61*, 152241.
6. Representative NMR data. Compound **4a**. ^1H : δ 7.53 (t, $J = 6.3$ Hz, 1H, H7); 5.43 (dd, $J = 3.4, 1.1$ Hz, 1H, H4); 5.32 (dd, $J = 10.4, 8.4$ Hz, 1H, H2); 5.15 (d, $J = 8.45$ Hz, 1H, H1); 5.10 (dd, $J = 10.35, 3.4$ Hz, 1H, H3); 4.20 (d, $J = 6.8$ Hz, 2H, H6); 4.03 (td, $J = 6.84, 1.20$ Hz, 1H, H5); 2.24–1.27 (m, 16H, H 8–15); 2.17 (s, 3H, CH₃); 2.08 (s, 3H, CH₃); 2.06 (s, 3H, CH₃); 2.01 (s, 3H, CH₃); 0.90 (t, 3H, H16). ^{13}C : 170.35 (C=O); 170.28 (C=O); 170.14 (C=O); 169.50 (C=O); 155.11 (C7); 102.35 (C1); 71.15 (C3); 70.71 (C5); 67.54 (C2); 66.86 (C4); 61.01 (C6); 31.90–22.66 (CH₂); 20.80–20.59 (CH₃); 14.11 (C16).
7. Pifferi, C.; Daskhan, G.C.; Fiore, M.; Shiao, T.C.; Roy, R.; Renaudet, O. Aminoxylated carbohydrates: Synthesis and applications. *Chem. Rev.* **2017**, *117*, 9839–9873.
8. Kalia, J.; Raines, R.T. Hydrolytic stability of hydrazones and oximes. *Angew. Chem. Int. Ed.* **2008**, *47*, 7523–7526.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).