

Article

# New black hole solutions in $\mathcal{N} = 2$ and $\mathcal{N} = 8$ gauged supergravity

Antonio Gallerati <sup>1,2,\*</sup> 

<sup>1</sup> Politecnico di Torino, Dipartimento di Scienza Applicata e Tecnologia, corso Duca degli Abruzzi 24, 10129 Torino, Italy

<sup>2</sup> Istituto Nazionale di Fisica Nucleare, Sezione di Torino, via Pietro Giuria 1, 10125 Torino, Italy

\* Correspondence: antonio.gallerati@polito.it (A.G.)

**Abstract:** We present a  $\mathcal{N} = 2$  supergravity model that interpolates between all the single dilaton truncations of the gauged  $SO(8)$ ,  $\mathcal{N} = 8$  supergravity. We provide new explicit non-extremal charged black hole solutions and their supersymmetric limits, exploiting the non-trivial transformation of the Fayet-Iliopoulos parameters under electromagnetic duality to connect the electric and the magnetic configurations. We also provide the asymptotic charges, thermodynamics and boundary conditions of these black hole configurations. We then construct a new supersymmetric truncation of the maximal supergravity, this new sector featuring non-extremal and supersymmetric black holes.

**Keywords:** supergravity; black hole solutions; black holes thermodynamics; BPS black holes.

## 1. Introduction

Anti-de Sitter (AdS) black hole configurations are an amazing and stimulating field of research due to the role they play in high energy theory as well as in the phenomenology of the AdS/CFT conjecture [1]. In particular, classical AdS black hole solutions can reveal specific features about the dual, strongly coupled gauge theory, providing then a possible description of many condensed matter phenomena.

The thermodynamic properties of AdS black holes were first analyzed in [2] and subsequently extended to various other AdS black hole configurations [3–6]. These researches described how black holes feature specific phase structures, giving rise to critical phenomena analogous to other common thermodynamic systems. Of particular interest are black hole solutions preserving a certain amount of supersymmetry, since they allow to map a weak (string) coupling description of the system thermodynamics to the strong-coupling regime, where a formulation in terms of a black hole configuration is valid [7]. Moreover, these solutions can be exploited to study the BPS attractor flows in AdS spacetime [8–21].

In the following we will discuss the new exact charged hairy black hole solutions in gauged  $\mathcal{N} = 2$ ,  $D = 4$  supergravity of [22]<sup>1</sup>, interpolating between four single dilaton truncations of the maximal  $SO(8)$ ,  $\mathcal{N} = 8$  supergravity. In particular, we will provide the explicit expressions for two new different families of non-extremal black hole solutions, analysing the duality relation between them. We will then investigate the thermodynamic properties of our new solutions, together with the analysis of the related boundary conditions. We will also study which conditions for the parameters give rise to BPS configuration ones. Finally, we will characterize certain models within the general class under consideration as consistent truncations of the maximal  $\mathcal{N} = 8$ ,  $D = 4$  gauged supergravity.



Citation: . *Proceedings* **2021**, *1*, 0.

Academic Editor:

Received:

Accepted:

Published:

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

<sup>1</sup> these new solutions generalize the uncharged configurations of [23,24]

## 2. Results

Let us consider an extended  $\mathcal{N} = 2$  supergravity theory in  $D = 4$ , coupled to  $n_v$  vector multiplets and no hypermultiplets, in the presence of Fayet–Iliopoulos (FI) terms. The model describes  $n_v + 1$  vector fields  $A_\mu^\Lambda$ , ( $\Lambda = 0, \dots, n_v$ ) and  $n_s = n_v$  complex scalar fields  $z^i$  ( $i = 1, \dots, n_s$ ). The bosonic gauged Lagrangian has the explicit form

$$\frac{1}{\sqrt{-g}} \mathcal{L}_{\text{BOS}} = -\frac{R}{2} + g_{i\bar{j}} \partial_\mu z^i \partial^\mu \bar{z}^{\bar{j}} + \frac{1}{4} \mathcal{I}_{\Lambda\Sigma}(z, \bar{z}) F_{\mu\nu}^\Lambda F^{\Sigma\mu\nu} + \frac{1}{8\sqrt{-g}} \mathcal{R}_{\Lambda\Sigma}(z, \bar{z}) \varepsilon^{\mu\nu\rho\sigma} F_{\mu\nu}^\Lambda F_{\rho\sigma}^\Sigma - V(z, \bar{z}) \quad (1)$$

where  $g = \det(g_{\mu\nu})$  and with the  $n_v + 1$  vector field strengths:

$$F_{\mu\nu}^\Lambda = \partial_\mu A_\nu^\Lambda - \partial_\nu A_\mu^\Lambda. \quad (2)$$

The  $n_s$  complex scalars  $z^i$  couple to the vector fields in a non-minimal way through the real symmetric matrices  $\mathcal{I}_{\Lambda\Sigma}(z, \bar{z})$ ,  $\mathcal{R}_{\Lambda\Sigma}(z, \bar{z})$  and span a special Kähler manifold  $\mathcal{M}_{\text{SK}}$ , while the scalar potential  $V(z, \bar{z})$  originates from electric-magnetic FI terms.

### 2.1. The model

Let us now focus on a  $\mathcal{N} = 2$  theory with no hypermultiplets and a single vector multiplet ( $n_v = 1$ ), with a complex scalar field  $z$ . The geometry of the special Kähler manifold is characterized by a prepotential of the form:

$$\mathcal{F}(\mathcal{X}^\Lambda) = -\frac{i}{4} (\mathcal{X}^0)^n (\mathcal{X}^1)^{2-n}, \quad (3)$$

$\mathcal{X}^\Lambda(z)$  being components of a holomorphic section of the symplectic bundle over the manifold and the coordinate  $z$  being identified with the ratio  $\mathcal{X}^1/\mathcal{X}^0$ , in a local patch in which  $\mathcal{X}^0 \neq 0$ . For special values of  $n$ , the model is a consistent truncation of the STU model<sup>2</sup>.

If we set  $\mathcal{X}^0 = 1$ , the holomorphic section  $\Omega^M$  of the model reads:

$$\Omega^M = \begin{pmatrix} 1 \\ z \\ -\frac{i}{4} n z^{2-n} \\ -\frac{i}{4} (2-n) z^{1-n} \end{pmatrix}, \quad (4)$$

and the Kähler potential  $\mathcal{K}$  has the expression

$$e^{-\mathcal{K}} = \frac{1}{4} z^{1-n} (n z - (n-2)\bar{z}) + \text{c.c.} \quad (5)$$

The theory is deformed with the introduction of abelian electric-magnetic FI terms defined by a constant symplectic vector  $\theta_M = (\theta_1, \theta_2, \theta_3, \theta_4)$ , encoding the gauge parameters of the model. The scalar potential  $V(z, \bar{z})$  can be then obtained from:

$$V = \left( g^{i\bar{j}} U_i^M \bar{U}_{\bar{j}}^N - 3 \mathcal{V}^M \bar{\mathcal{V}}^N \right) \theta_M \theta_N = -\frac{1}{2} \theta_M \mathcal{M}^{MN} \theta_N - 4 \mathcal{V}^M \bar{\mathcal{V}}^N \theta_M \theta_N, \quad (6)$$

where  $\mathcal{V}^M = e^{\mathcal{K}/2} \Omega^M$ ,  $U_i^M = \mathcal{D}_i \mathcal{V}^M$  and  $\mathcal{M}(\phi)$  is the symplectic, symmetric, negative definite matrix encoding the non-minimal couplings of the scalars to the vector fields of the theory.

<sup>2</sup> the STU model [25–27] is a  $\mathcal{N} = 2$  supergravity coupled to  $n_v = 3$  vector multiplets and characterized, in a suitable symplectic frame, by the prepotential  $\mathcal{F}_{\text{STU}}(\mathcal{X}^\Lambda) = -\frac{i}{4} \sqrt{\mathcal{X}^0 \mathcal{X}^1 \mathcal{X}^2 \mathcal{X}^3}$ , together with symmetric scalar manifold of the form  $\mathcal{M}_{\text{STU}} = (\text{SL}(2, \mathbb{R})/\text{SO}(2))^3$  spanned by the three complex scalars  $z^i = \mathcal{X}^i/\mathcal{X}^0$  ( $i = 1, 2, 3$ ); this model is in turn a consistent truncation of the maximal  $\mathcal{N} = 8$  theory in four-dimensions with  $\text{SO}(8)$  gauge group [28–30]

Writing  $z = e^{\lambda\phi} + i\chi$ , the truncation to the dilaton field  $\phi$  (i.e.  $\chi = 0$ ) is consistent provided:

$$(2-n)\theta_1\theta_3 - n\theta_2\theta_4 = 0, \quad (7)$$

and the metric restricted to the dilaton reads:

$$ds^2 = 2g_{z\bar{z}} dz d\bar{z} \Big|_{\substack{\chi=0 \\ d\chi=0}} = \frac{1}{2}\lambda^2 n(2-n) d\phi^2, \quad (8)$$

and is positive provided  $0 < n < 2$ . Choosing  $\lambda = \sqrt{\frac{2}{n(2-n)}}$  the kinetic term for  $\phi$  is then canonically normalized. As a function of the dilaton only, the scalar potential has the following explicit form:

$$\begin{aligned} V(\phi) = & -2e^{\lambda\phi(n-2)} \left( \frac{2n-1}{n} \theta_1^2 + 4\theta_1\theta_2 e^{\lambda\phi} + \frac{2n-3}{n-2} \theta_2^2 e^{2\lambda\phi} \right) - \\ & - \frac{1}{8} e^{-\lambda\phi(n-2)} \left( (2n-1)n\theta_3^2 - 4\theta_3\theta_4 n(n-2) e^{-\lambda\phi} + (n-2)(2n-3)\theta_4^2 e^{-2\lambda\phi} \right). \end{aligned} \quad (9)$$

### 2.1.1. Redefinitions

Let us now make the shift

$$\phi \rightarrow \phi - \frac{2\nu}{\lambda(\nu+1)} \log(\theta_2 \xi), \quad (10)$$

and redefine the FI terms as:

$$\theta_1 = \frac{\nu+1}{\nu-1} \theta_2^{-\frac{\nu-1}{\nu+1}} \xi^{-\frac{2\nu}{\nu+1}}, \quad \theta_3 = 2\alpha(\xi\theta_2)^{\frac{\nu-1}{\nu+1}} s, \quad \theta_4 = \frac{2\alpha}{\theta_2 \xi s}, \quad (11)$$

having defined the quantity  $\nu = (n-1)^{-1}$  and having also introduced the parameters  $\alpha, s$  and

$$\xi = \frac{2L\nu}{\nu-1} \frac{1}{\sqrt{1-\alpha^2 L^2}}, \quad (12)$$

expressed in terms of the AdS radius  $L$ . Let us recall that the truncation to the dilaton is consistent provided equation (7) is satisfied and, in light of the new parametrization (11), this condition requires

$$(s^2 - 1)(\nu^2 - 1)\alpha\sqrt{1 - L^2\alpha^2} = 0, \quad (13)$$

which is solved, excluding values  $n = 0$  and  $n = 2$ , either for pure electric FI terms ( $\alpha = 0$ ) or for  $s = \pm 1$ . Since we are interested in dyonic FI terms, we shall restrict ourselves to the latter case.

After the shift (10), the scalar field  $z$  is expressed as

$$z = (\theta_2 \xi)^{-\frac{2\nu}{\nu+1}} e^{\lambda\phi}, \quad (14)$$

and the same redefinition for the potential (with  $s = \pm 1$ ) yields

$$\begin{aligned} V(\phi) = & -\frac{\alpha^2}{\nu^2} \left( \frac{(\nu-1)(\nu-2)}{2} e^{-\phi\ell(\nu+1)} + 2(\nu^2-1) e^{-\phi\ell} + \frac{(\nu+1)(\nu+2)}{2} e^{\phi\ell(\nu-1)} \right) + \\ & + \frac{\alpha^2 - L^{-2}}{\nu^2} \left( \frac{(\nu-1)(\nu-2)}{2} e^{\phi\ell(\nu+1)} + 2(\nu^2-1) e^{\phi\ell} + \frac{(\nu+1)(\nu+2)}{2} e^{-\phi\ell(\nu-1)} \right), \end{aligned} \quad (15)$$

where  $\ell = \frac{\lambda}{\nu}$  and having disposed of  $\theta_2$  by the above redefinitions.

After the restriction to the dilaton truncation, the matrix  $\mathcal{R}_{\Lambda\Sigma}$  vanishes and the action has the form

$$\mathcal{S} = \frac{1}{8\pi G} \int d^4x \sqrt{-g} \left( -\frac{R}{2} + \frac{\partial_\mu \phi \partial^\mu \phi}{2} + \frac{1}{4} \mathcal{I}_{\Lambda\Sigma}(\phi) F_{\mu\nu}^\Lambda F^{\Sigma\mu\nu} - V(\phi) \right), \quad (16)$$

If we define the canonically normalized gauge fields

$$\bar{F}^1 = \frac{1}{2} \sqrt{\frac{1+\nu}{\nu}} (\theta_2 \xi)^{\frac{1-\nu}{1+\nu}} F^1, \quad \bar{F}^2 = \frac{1}{2} \sqrt{\frac{-1+\nu}{\nu}} (\theta_2 \xi) F^2, \quad (17)$$

the action can be expressed as

$$\mathcal{S} = \frac{1}{8\pi G} \int d^4x \sqrt{-g} \left( -\frac{R}{2} + \frac{\partial_\mu \phi \partial^\mu \phi}{2} - \frac{1}{4} e^{(-1+\nu)\ell\phi} (\bar{F}^1)^2 - \frac{1}{4} e^{-(1+\nu)\ell\phi} (\bar{F}^2)^2 - V(\phi) \right). \quad (18)$$

## 2.2. Hairy black hole solutions

Now we construct two distinct families of solutions, which we refer to as electric and magnetic, respectively.

### 2.2.1. Family 1 - Electric solutions

A first family of solutions is given by

$$\phi = -\ell^{-1} \ln(x), \quad \bar{F}_{tx}^1 = Q_1 x^{-1+\nu}, \quad \bar{F}_{tx}^2 = Q_2 x^{-1-\nu}, \quad Y(x) = \frac{x^{\nu-1} \nu^2 L^2}{\eta^2 (x^\nu - 1)^2}, \quad (19a)$$

$$f(x) = \frac{x^{2-\nu} \eta^2 (x^\nu - 1)^2}{\nu^2} \frac{k}{L^2} + \alpha^2 L^2 \left( -1 + \frac{x^2}{\nu^2} \left( (\nu + 2) x^{-\nu} - (\nu - 2) x^\nu + \nu^2 - 4 \right) \right) + 1 + \frac{x^{2-\nu} \eta^2 (x^\nu - 1)^3}{\nu^3 L^2} \left( \frac{Q_1^2}{(\nu + 1)} - \frac{Q_2^2}{(\nu - 1)} x^{-\nu} \right), \quad (19b)$$

$$ds^2 = Y(x) \left( f(x) dt^2 - \frac{\eta^2}{f(x)} dx^2 - L^2 d\Sigma_k \right), \quad (19c)$$

where  $d\Sigma_k^2 = d\theta^2 + \frac{\sin^2(\sqrt{k}\theta)}{k} d\varphi^2$  is the metric on the 2D-surfaces  $\Sigma_k = \{\mathbb{S}^2, \mathbb{H}^2, \mathbb{R}^2\}$  (sphere, hyperboloid and flat space) with constant scalar curvature  $R = 2k$ .

Boundary conditions, mass and thermodynamics for the electric solutions.

To make contact with AdS canonical coordinates, we consider the following fall-off for the metric function:

$$Y(x) = \frac{r^2}{L^2} + O(r^{-2}). \quad (20)$$

The change of coordinates that provides the above asymptotic behaviour is given by

$$x = 1 \pm \left( \frac{L^2}{\eta r} + L^6 \frac{1 - \nu^2}{24 (\eta r)^3} \right) + L^8 \frac{\nu^2 - 1}{24 (\eta r)^4}, \quad (21)$$

where the  $+$  ( $-$ ) sign holds for  $x > 1$  ( $x < 1$ ). The corresponding asymptotic behaviour of the scalar field is

$$\phi = L^2 \frac{\phi_0}{r} + L^4 \frac{\phi_1}{r^2} + O(r^{-3}) = \mp L^2 \frac{1}{\ell \eta r} + L^4 \frac{1}{2\ell \eta^2 r^2} + O(r^{-3}), \quad (22)$$

with normalized  $\phi_0$  and  $\phi_1$ , to match their conformal and engineering dimension. The coefficients of the leading and subleading terms in the scalar asymptotic expansion read

$$\phi_0 = \mp \frac{1}{\ell \eta}, \quad \phi_1 = \frac{\ell}{2} \phi_0^2, \quad (23)$$

and corresponds to AdS invariant boundary conditions, the boundary conformal symmetry being then preserved. The asymptotic expansion of the spacetime (19) reads

$$g_{tt} = \frac{r^2}{L^2} + k - \frac{\mu_E L^4}{r} + O(r^{-2}), \quad g_{rr} = -\frac{L^2}{r^2} - L^6 \frac{k L^{-2} + \frac{1}{2} \phi_0^2}{r^4} + O(r^{-5}), \quad (24)$$

where

$$\mu_E = \pm \left( \frac{\nu^2 - 4}{3\eta^3} \alpha^2 L^2 - \frac{k}{\eta L^2} + \frac{Q_2^2}{\eta(\nu-1)L^2} - \frac{Q_1^2}{\eta(\nu+1)L^2} \right). \quad (25)$$

The black hole mass can be read-off from the above expansion [31–33] and reads

$$M_E = L^4 \frac{\mu_E \sigma_k}{8\pi G}, \quad (26)$$

where  $\sigma_k = \int d\Sigma_k$ , while the temperature and the entropy are given by

$$T = \left. \frac{|f(x)'|}{4\pi \eta} \right|_{x=x_+}, \quad S = \frac{L^2 Y(x_+) \sigma_k}{4G}, \quad (27)$$

where  $f(x_+) = 0$ . The physical charges and electric potentials are

$$q_1 = \frac{L^2 Q_1 \sigma_k}{8\pi G \eta}, \quad \Phi_1^E = Q_1 \frac{x_+^\nu - 1}{\nu}; \quad q_2 = \frac{L^2 Q_2 \sigma_k}{8\pi G \eta}, \quad \Phi_2^E = Q_2 \frac{1 - x_+^{-\nu}}{\nu}, \quad (28)$$

and it is possible to verify that these quantities satisfy the first law of thermodynamics:

$$dM_E = T dS + \Phi_1^E dq_1 + \Phi_2^E dq_2. \quad (29)$$

### 2.2.2. Family 2 - Magnetic Solutions

A second family of solutions is given by

$$\phi = \ell^{-1} \ln(x), \quad \bar{F}_{\theta\varphi}^1 = P_1 \frac{\sin(\sqrt{k}\theta)}{\sqrt{k}}, \quad \bar{F}_{\theta\varphi}^2 = P_2 \frac{\sin(\sqrt{k}\theta)}{\sqrt{k}}, \quad Y(x) = \frac{x^{\nu-1} \nu^2 L^2}{\eta^2 (x^\nu - 1)^2}, \quad (30a)$$

$$f(x) = \frac{x^{2-\nu} \eta^2 (x^\nu - 1)^2}{\nu^2} \frac{k}{L^2} + \left( 1 - \alpha^2 L^2 \right) \left( -1 + \frac{x^2}{\nu^2} \left( (\nu+2)x^{-\nu} - (\nu-2)x^\nu + \nu^2 - 4 \right) \right) + 1 + \frac{x^{2-\nu} \eta^4 (x^\nu - 1)^3}{\nu^3 L^6} \left( \frac{P_1^2}{(\nu+1)} - \frac{P_2^2}{(\nu-1)} x^{-\nu} \right), \quad (30b)$$

$$ds^2 = Y(x) \left( f(x) dt^2 - \frac{\eta^2}{f(x)} dx^2 - L^2 d\Sigma_k \right). \quad (30c)$$

The electric and magnetic solutions are related to each other by means of electromagnetic duality

$$\phi \rightarrow -\phi, \quad \alpha^2 \rightarrow L^{-2} - \alpha^2, \quad (31)$$

and corresponding transformation of the electromagnetic fields. In each family, the asymptotic region is located at the pole of order 2 of the conformal factor  $Y(x)$ , namely  $x = 1$ . The

geometry and scalar field are singular at  $x = 0$  and  $x = \infty$  and, therefore, the configuration contains two disjoint geometries given by  $x \in (1, \infty)$  or  $x \in (0, 1)$ .

Boundary conditions, mass and thermodynamics for the magnetic solutions.

The metric of the magnetic family can be obtained from the electric solutions by means of the transformation

$$Q_i \rightarrow \frac{\eta}{L^2} P_i, \quad \alpha^2 \rightarrow L^{-2} - \alpha^2, \quad (32)$$

therefore all the discussions on mass and thermodynamics carry on from the electric to the magnetic case. The main difference is that the scalar boundary leading and subleading terms now satisfy

$$\phi_1 = -\frac{\ell}{2} \phi_0^2, \quad (33)$$

which correspond again to AdS invariant boundary conditions. We define now

$$\mu_M = \pm \left( \frac{\nu^2 - 4}{3\eta^3} (1 - \alpha^2 L^2) - \frac{k}{\eta L^2} + \frac{\eta P_2^2}{(\nu - 1)L^6} - \frac{\eta P_1^2}{(\nu + 1)L^6} \right), \quad (34)$$

where the  $+$  ( $-$ ) corresponds to  $x > 1$  ( $x < 1$ ). The black hole mass is

$$M_M = L^4 \frac{\mu_M \sigma_k}{8\pi G}, \quad (35)$$

while the magnetic charges and potentials are

$$p_1 = \frac{P_1 \sigma_k}{8\pi G}, \quad \Phi_1^M = \frac{P_1 \eta}{L^2} \frac{x_+^\nu - 1}{\nu}; \quad p_2 = \frac{P_2 \sigma_k}{8\pi G}, \quad \Phi_2^M = \frac{P_2 \eta}{L^2} \frac{1 - x_+^{-\nu}}{\nu}. \quad (36)$$

It is possible to verify that these quantities satisfy the first law of thermodynamics:

$$dM_M = T dS + \Phi_1^M dp_1 + \Phi_2^M dp_2. \quad (37)$$

### 3. Discussion

#### 3.1. Duality Relation between the Two Families of Solutions

The two families of solutions are related by a non-perturbative electric-magnetic duality symmetry. The latter is a global symmetry of the ungauged theory and is extended to the gauged one if the constant tensor  $\theta_M$  is made to transform under it as well. In general, a transformation of  $\theta_M$  would imply a change in the theory and the duality would be interpreted as an equivalence between different models. Such transformation can be absorbed in a redefinition of a single parameter  $\alpha$  in  $\theta_M$ : the two solutions related by duality satisfy the field equations of the same model with two (dual) values of the  $\alpha$  parameter.

Consider a generic  $\mathcal{N} = 2$  theory described by the Lagrangian (1), where the scalar manifold (which is of special Kähler type in the absence of hypermultiplets) is now described in terms of real scalars  $\phi(x) \equiv \phi^s(x)$  and where  $\mathcal{G}_{rs}(\phi)$  is the metric on the scalar manifold. The scalar potential  $V(\theta, \phi)$  is given by eq. (6), explicitly depending on both the scalar fields and the FI  $\theta_M$ . To describe the duality it is useful to define the magnetic field strengths as:

$$G_{\Lambda\mu\nu} = -\epsilon_{\mu\nu\rho\sigma} \frac{\delta \mathcal{L}_{\text{BOS}}}{\delta F_{\rho\sigma}^\Lambda} = \mathcal{R}_{\Lambda\Sigma} F_{\mu\nu}^\Sigma - \mathcal{I}_{\Lambda\Sigma} {}^* F_{\mu\nu}^\Sigma, \quad (38)$$

and to introduce the symplectic field strength vector:

$$\mathbb{F}^M = \begin{pmatrix} F_{\mu\nu}^\Lambda \\ G_{\Lambda\mu\nu} \end{pmatrix}, \quad (39)$$

together with the  $2(n_v + 1) \times 2(n_v + 1)$  matrices

$$\mathcal{M}_{MN}(\phi) \equiv \begin{pmatrix} \mathcal{I}_{\Lambda\Sigma} + (\mathcal{R}\mathcal{I}^{-1}\mathcal{R})_{\Lambda\Sigma} & -(\mathcal{R}\mathcal{I}^{-1})_{\Lambda}{}^{\Gamma} \\ -(\mathcal{I}^{-1}\mathcal{R})^{\Delta}{}_{\Sigma} & (\mathcal{I}^{-1})^{\Delta\Gamma} \end{pmatrix}, \quad \mathbb{C} \equiv \begin{pmatrix} \mathbf{0} & \mathbb{1} \\ -\mathbb{1} & \mathbf{0} \end{pmatrix}. \quad (40)$$

A feature of special Kähler manifolds is the existence of a flat symplectic bundle structure on it, within which the matrix  $\mathcal{M}_{MN}$  can be regarded as a metric on the symplectic fiber. As a consequence of this, with each isometry  $\mathbf{g}$  in the isometry group  $G$  of the scalar manifold, there corresponds a constant  $2(n_v + 1) \times 2(n_v + 1)$  symplectic matrix  $\mathcal{R}[\mathbf{g}] = \mathcal{R}[\mathbf{g}]^M{}_N$  such that, if  $\phi' = \phi'^s = \phi'(\phi)$  describe the (non-linear) action of  $\mathbf{g}$  on the scalar fields, we have that  $\mathcal{M}_{MN}(\phi)$  transforms as a fiber metric [34,35]:

$$\mathcal{M}(\phi') = (\mathcal{R}[\mathbf{g}]^{-1})^T \mathcal{M}(\phi) \mathcal{R}[\mathbf{g}]^{-1}, \quad (41)$$

where we have suppressed the symplectic indices. If, for any isometry  $\mathbf{g}$ , we transform the fields  $\mathbb{F}_{\mu\nu}^M$  and the constant vector  $\theta_M$  correspondingly,

$$\mathbb{F}_{\mu\nu}^M \rightarrow \mathbb{F}'_{\mu\nu}{}^M = \mathcal{R}[\mathbf{g}]^M{}_N \mathbb{F}_{\mu\nu}^N, \quad \theta_M \rightarrow \theta'_M = (\mathcal{R}[\mathbf{g}]^{-1})^N{}_M \theta_N, \quad (42)$$

we find that the equations of motion are formally left invariant. Aside from this duality equivalence, one can just redefine the field strengths and the FI constants without altering the physics of the model:

$$\mathbb{F}_{\mu\nu} \rightarrow \begin{pmatrix} \mathbf{A} & \mathbf{0} \\ \mathbf{0} & (\mathbf{A}^{-1})^T \end{pmatrix} \mathbb{F}_{\mu\nu}, \quad \theta \rightarrow \begin{pmatrix} ((\mathbf{A}^{-1})^T & \mathbf{0}) \\ \mathbf{0} & \mathbf{A} \end{pmatrix} \theta, \quad (43)$$

where  $\mathbf{A} = (A^{\Lambda}{}_{\Sigma})$  is a generic real, invertible matrix, not related to the isometries of the scalar manifold. This just amounts to choosing a different basis in the symplectic fiber.

The above mechanism also works for our 1-scalar truncated model (with trivial metric  $\mathcal{G}(\phi) = 1$ ), provided the isometry does not switch on the truncated axion  $\chi$  (imaginary part of  $z$ ). The absence of the axion also implies  $\mathcal{R}_{\Lambda\Sigma} = 0$ . The isometries of the scalar field are

$$\phi \rightarrow \phi' = \phi + \beta, \quad (44a)$$

$$\phi \rightarrow \phi' = -\phi, \quad (44b)$$

$\beta$  being a constant. The first of the above isometries was used in Subsect. 2.1.1 to reabsorb the  $\theta_2$  dependence of the FI terms, making them basically only depending on  $\alpha$ . This was then followed by a corresponding redefinition of the field strengths and of the FI parameters of the kind (43). Let us now consider the duality transformation associated with the isometry (44b). Using the property  $\mathcal{I}_{\Lambda\Sigma}(-\phi) = (\mathcal{I}^{-1})^{\Lambda\Sigma}(\phi)$ , it is easily verified that the associated duality transformation  $\mathcal{R}[\mathbf{g}]$  is simply given by  $\mathbb{C}$ . Indeed we have:

$$\mathcal{M}(\phi') = \mathcal{M}(-\phi) = (\mathbb{C}^{-1})^T \mathcal{M}(\phi) \mathbb{C}^{-1}. \quad (45)$$

By the same token we can compute the duality transformed field strengths:

$$\begin{pmatrix} \bar{F}'_{\mu\nu}{}^{\Lambda} \\ \bar{G}'_{\Lambda\mu\nu} \end{pmatrix} = \mathbb{C} \begin{pmatrix} \bar{F}_{\mu\nu}{}^{\Lambda} \\ \bar{G}_{\Lambda\mu\nu} \end{pmatrix} = \begin{pmatrix} \bar{G}_{\Lambda\mu\nu} \\ -\bar{F}_{\mu\nu}{}^{\Lambda} \end{pmatrix}. \quad (46)$$

After the action of the shift symmetry (44a) and the above mentioned redefinition, the resulting FI parameter vector, to be denoted by  $\bar{\theta}_M$ , has the form:

$$\bar{\theta}_M = \left( \sqrt{\frac{v+1}{v}} \sqrt{L^{-2} - \alpha^2}, \sqrt{\frac{v-1}{v}} \sqrt{L^{-2} - \alpha^2}, \alpha s \sqrt{\frac{v+1}{v}}, \frac{\alpha}{s} \sqrt{\frac{v-1}{v}} \right). \quad (47)$$

We denote by  $\bar{\theta}'$  the transformed of  $\bar{\theta}$  by  $\mathbb{C}$ , that is  $\bar{\theta}' = \mathbb{C}\bar{\theta}$ . One can then verify that

$$V(\bar{\theta}', \phi') = V(\bar{\theta}, \phi). \quad (48)$$

Since  $\bar{\theta}$  only depends on  $\alpha$ , we will simply denote the potential by  $V(\alpha, \phi)$ . This action, apart from signs, basically amounts in  $\bar{\theta}$  to changing  $\alpha^2 \rightarrow L^{-2} - \alpha^2$ . In particular we find:

$$V(\alpha, \phi) = V(\alpha', \phi'), \quad (49)$$

where  $\alpha'^2 = L^{-2} - \alpha^2$ . It is possible to verify that, if the electric solutions

$$\phi(x), \quad \bar{F}_{\mu\nu}^\Lambda(x), \quad g_{\mu\nu}(x), \quad (50)$$

solve the field equations with FI terms defined by the parameter  $\alpha$ , then the magnetic configuration

$$\phi'(x) = -\phi(x), \quad \bar{G}'_{\Lambda\mu\nu}(x) = -\bar{F}_{\mu\nu}^\Lambda(x), \quad g'_{\mu\nu}(x) = g_{\mu\nu}(x), \quad (51)$$

is a solution with parameter  $\alpha'$ . The identification  $\bar{G}'_{\Lambda\mu\nu}(x) = -\bar{F}_{\mu\nu}^\Lambda(x)$  means that the magnetic field strengths (apart from a sign) will have the same form as the original electric ones, so that we can relate its magnetic charge parameters  $P'_i$  with the original electric ones:  $P'_i = -Q_i$ .

### 3.2. Supersymmetric solutions

Now we want to study supersymmetric configurations for our model, imposing the vanishing of the SUSY variations [22,36]. First it is useful to make a change of coordinates that puts metric (19c) in the standard form

$$ds^2 = e^{2U(r)} dt^2 - e^{-2U(r)} \left( dr^2 + e^{2\Psi(r)} d\Sigma_k^2 \right). \quad (52)$$

This can be achieved through the change of coordinate

$$x(r) = \left( 1 + \frac{L^2 \nu}{\eta (r - c)} \right)^{\frac{1}{\nu}}, \quad (53)$$

$c$  being a constant.

#### 3.2.1. Family 1

The scalar field  $z$  in the new parametrization has the form

$$z = (\theta_2 \xi)^{-\frac{2\nu}{1+\nu}} \left( 1 + \frac{L^2 \nu}{\eta (r - c)} \right)^{-1}, \quad (54)$$

while the electric-magnetic charges explicitly read

$$\Gamma^M = \begin{pmatrix} m^\Lambda \\ e_\Lambda \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \frac{L^2}{2\eta} Q_1 \sqrt{\frac{1+\nu}{\nu}} (\theta_2 \xi)^{\frac{1-\nu}{1+\nu}} \\ \frac{L^2}{2\eta} Q_2 \sqrt{\frac{-1+\nu}{\nu}} \theta_2 \xi \end{pmatrix}. \quad (55)$$



The solution is supersymmetric if

$$\begin{aligned} Q_1 &= -Q_2 \sqrt{\frac{-1+\nu}{1+\nu}} + \frac{k\eta}{\alpha L^2} \sqrt{\frac{\nu}{1+\nu}}, \\ Q_2 &= \left( \frac{k\eta}{2\alpha L^2} + \frac{\alpha L^2(1+\nu)}{2\eta} \right) \sqrt{\frac{-1+\nu}{\nu}}. \end{aligned} \tag{56}$$

### 3.2.2. Family 2

The scalar field  $z$  reads

$$z = (\theta_2 \xi)^{-\frac{2\nu}{1+\nu}} \left( 1 + \frac{L^2 \nu}{\eta(r-c)} \right), \tag{57}$$

while the electric-magnetic charges have the form

$$\Gamma^M = \begin{pmatrix} m^\Lambda \\ e_\Lambda \end{pmatrix} = \begin{pmatrix} 2P_1 \sqrt{\frac{\nu}{1+\nu}} (\theta_2 \xi)^{\frac{-1+\nu}{1+\nu}} \\ 2P_2 \sqrt{\frac{\nu}{-1+\nu}} (\theta_2 \xi)^{-1} \\ 0 \\ 0 \end{pmatrix}. \tag{58}$$

The solution is supersymmetric if

$$\begin{aligned} P_1 &= -P_2 \sqrt{\frac{-1+\nu}{1+\nu}} + \frac{kL}{\sqrt{1-\alpha^2 L^2}} \sqrt{\frac{\nu}{1+\nu}}, \\ P_2 &= \left( \frac{kL}{2\sqrt{1-\alpha^2 L^2}} + \frac{L^3(1+\nu)}{2\eta^2} \sqrt{1-\alpha^2 L^2} \right) \sqrt{\frac{-1+\nu}{\nu}}. \end{aligned} \tag{59}$$

The supersymmetric magnetic condition can be obtained from the supersymmetric electric condition by means of the duality transformation

$$Q_i \rightarrow \frac{\eta}{L^2} P_i, \quad \alpha^2 \rightarrow L^{-2} - \alpha^2. \tag{60}$$

### 3.3. BPS black holes of finite area

#### 3.3.1. Family 1: BPS electric black holes

We found that the electric family has BPS black holes of finite area only when  $\alpha^2 = L^{-2}$ . In this case, the lapse function has a double zero, as expected:

$$f(x_+) = 0 \implies x_\pm^\nu = \frac{\nu^2 - 1 - k\eta^2 L^{-2}}{\nu - 1 - k\eta^2 L^{-2}} \pm \nu \frac{\sqrt{\nu^2 - 1 - 2k\eta^2 L^{-2}}}{\nu - 1 - k\eta^2 L^{-2}}. \tag{61}$$

It is possible to verify that  $x_\pm(\nu) = x_\pm(-\nu)$ , namely,  $x_\pm$  is an even function of  $\nu$ . Hence, we can restrict our analysis to the  $\nu > 1$  interval.

The asymptotic region of the spacetime is located at  $x = 1$  and, therefore, the region of the spacetime at  $x > 1$  is disconnected from the region at  $x < 1$ , representing then different spacetimes. The location of the horizon can be characterized as follows:

- $k =$  in the flat case the location of the horizon is very simple (see the above (61))
- $0:$  and it follows that  $x_+^\nu > 0$  and  $x_-^\nu < 0$ , so we conclude that only  $x_+$  exists;
- $k =$  in the hyperbolic case,  $x_+ > 1$  always exists while the solution  $0 < x_- < 1$
- $-1:$  exists provided  $\eta^2 L^{-2} > \nu + 1$ ;
- $k =$  for spherical black hole only  $x_+$  exists, provided  $\nu - 1 > \eta^2 L^{-2} > 0$ .
- $+1:$

### 3.3.2. Family 2: BPS Magnetic Black Holes

Again, we shall consider only the case  $\nu > 1$ . This family has BPS black holes of finite area only when  $\alpha^2 = 0$ , namely when the gauging is purely electric. In this case the metric of the magnetic solution exactly coincides with the metric of the electric solutions. Hence the analysis of the location of the horizons is exactly the same. The extremality of the magnetic solutions is the same as for the electric solutions when the electric charges and potentials are interchanged by their magnetic counterparts.

### 3.4. $\mathcal{N} = 8$ Truncations

#### 3.4.1. Uncharged Case

The infinitely many theories we have described in the previous sections contain all the possible one-dilaton consistent truncations of the  $\omega$ -deformed  $SO(8)$  gauged maximal supergravities. Let us briefly discuss this result.

In the dyonic models with gauge group  $SO(8)$ , the 28 generators are gauged by the 28 vector fields in a symplectic frame which is related to the one of [37,38] by an  $SO(2)$  transformation parameterized by an angle  $\omega$ . The physically independent values of  $\omega$  lie within the interval  $0 \leq \omega \leq \pi/8$ ,  $\omega = 0$  corresponding to the original theory by de Wit and Nicolai. The 70 scalar fields parameterize the coset  $E_{7(7)}/SU(8)$  and, upon truncation to gravity and scalar field sector<sup>3</sup>, we are led to consider the following action [39,40]:

$$I(g_{\mu\nu}, \vec{\phi}) = \int_{\mathcal{M}} d^4x \sqrt{-g} \left[ -\frac{R}{2} + \frac{1}{2} (\partial\vec{\phi})^2 - V(\vec{\phi}) \right], \quad (62)$$

where  $\vec{\phi} = (\phi_i)$ , with  $i = 1, \dots, 8$  and  $\sum_{i=1}^8 \phi_i = 0$ . The potential is given by

$$V(\vec{\phi}) = -\frac{g^2}{32} \left[ \cos^2(\omega) \left( \left( \sum_{i=1}^8 X_i \right)^2 - 2 \sum_{i=1}^8 X_i^2 \right) + \sin^2(\omega) \left( \left( \sum_{i=1}^8 X_i^{-1} \right)^2 - 2 \sum_{i=1}^8 X_i^{-2} \right) \right], \quad (63)$$

where

$$X_i = e^{2\phi_i}, \quad \prod_{i=1}^8 X_i = 1. \quad (64)$$

Let us consider now a single scalar field reduction preserving  $SO(p) \times SO(8-p)$ . This is effected through the following identification:

$$\phi_1 = \dots = \phi_p = \frac{1}{2\sqrt{2}} \sigma \phi, \quad \phi_{p+1} = \dots = \phi_8 = -\frac{1}{2\sqrt{2}} \frac{\phi}{\sigma}, \quad (65)$$

where we have defined:

$$\sigma = \sqrt{\frac{8-p}{p}} = \sqrt{\frac{\nu-1}{\nu+1}}, \quad p = \frac{4(\nu+1)}{\nu}. \quad (66)$$

With the above choice we have:

$$X_1 = \dots = X_p = X := e^{\frac{1}{\sqrt{2}} \sigma \phi}, \quad X_{p+1} = \dots = X_8 = Y := e^{-\frac{1}{\sqrt{2}} \frac{\phi}{\sigma}}, \quad (67)$$

and the previous action (62) reduces to the one we are studying in this paper. The action is invariant under  $\sigma \rightarrow 1/\sigma$ ,  $\phi \rightarrow -\phi$  and  $p \rightarrow 8-p$ . This action, consistent truncation

<sup>3</sup> we choose a parametrization covariant under  $SU(8)$ , the scalars splitting into the representations  $35_c$  and  $35_v$  of the gauge group  $SO(8)$ : the former can be truncated out, while the latter scalars span the submanifold  $SL(8, \mathbb{R})/SO(8)$ ; the local  $SO(8)$  transformations can be used to diagonalize the coset representative and thus to further truncate the theory to the seven scalars parameterizing the non compact Cartan subalgebra of  $SL(8, \mathbb{R})$

of the  $\omega$ -rotated  $\text{SO}(8)$ -gauged maximal supergravity, coincides, in the absence of vector fields, with the action (16) upon the following identification:

$$g = \frac{\sqrt{2}}{L}, \quad \cos(\omega) = L\alpha, \quad \sin(\omega) = \sqrt{1 - L^2\alpha^2}. \quad (68)$$

Changing  $\phi$  into  $-\phi$  in (67), the above identifications change correspondingly:

$$g = \frac{\sqrt{2}}{L}, \quad \sin(\omega) = L\alpha, \quad \cos(\omega) = \sqrt{1 - L^2\alpha^2}. \quad (69)$$

From the above relation between  $p$  and  $v$ , we conclude that the single scalar field models considered in this work, if all vector fields are set to zero, coincide with truncations of the  $\omega$ -deformed  $\text{SO}(8)$  gauged maximal supergravity to the singlet sector with respect to the following subgroups of  $\text{SO}(8)$  gauge group :

$$\begin{aligned} v = \frac{4}{3} &\rightarrow \text{SO}(7), \\ v = 2 &\rightarrow \text{SO}(6) \times \text{SO}(2), \\ v = 4 &\rightarrow \text{SO}(5) \times \text{SO}(3), \\ v = \infty &\rightarrow \text{SO}(4) \times \text{SO}(4). \end{aligned} \quad (70)$$

The values  $v = \infty$  or  $v = \pm 2$  correspond to models which can be embedded in the STU truncation of the  $\text{SO}(8)$  gauged  $\mathcal{N} = 8$  supergravity. Therefore the black hole solutions discussed in this work, in the absence of electric and magnetic charges, can all be embedded in the maximal supergravity. Since the  $\text{SO}(8)$  gauged maximal supergravity can be uplifted to  $D = 11$  supergravity only for  $\omega = 0$  [41], only for  $L\alpha = 0$  or equivalently for  $L\alpha = \pm 1$  our solutions can be embedded in the eleven dimensional theory through maximal supergravity, by means of the formulas presented in [39,40].

### 3.4.2. Charged Case

The  $v = 4$  case.

Let us focus first on the  $\omega = 0$  case. The solutions describe gravity coupled to one scalar field and two vector fields. When identified with fields in the maximally supersymmetric model, the scalar and the two vectors should not excite the other fields in the model, such as the scalars in the  $\mathbf{35}_c$  of  $\text{SO}(8)$  (see footnote 3). This condition in turn require in our solution  $F^\Lambda \wedge F^\Sigma = 0$ .<sup>4</sup> Of the remaining scalar fields in the  $\mathbf{35}_v$  of  $\text{SO}(8)$ , 28 are gauged away so that we are left with the seven independent scalar fields  $\vec{\phi} = \phi_i$ , with  $i = 1, \dots, 8$  and  $\sum_{i=1}^8 \phi_i = 0$ , parameterizing the Cartan subalgebra of  $\mathfrak{e}_{7(7)}$  (Lie algebra of  $E_{7(7)}$ ).

Next we want to further truncate the theory to the scalar field  $\phi$ , which is singlet with respect to the subgroup  $\text{SO}(5) \times \text{SO}(3)$  of  $\text{SO}(8)$ . When the solution is charged, the two vector fields involved in it should be identified with two of the 28 vectors on the maximal

<sup>4</sup> only in this subsection we shall denote by  $\Lambda, \Sigma$  the indices labelling the 28 vectors of the maximal theory, by  $\lambda, \sigma = 1, 2$  those of the two vectors surviving the truncation, by  $M, N$  the symplectic indices of the 56 electric and magnetic charges, by  $m, n = 1, \dots, 4$  the symplectic index labelling the two electric and two magnetic charges in the truncated model; the corresponding indices in the new,  $\omega$ -rotated symplectic frame are distinguished from those in the original frame by a hat.

model which do not source the six scalar fields among  $\vec{\phi}$  that we wish to set to zero. To this end, let us write  $\vec{\phi}$  as follows:

$$\begin{aligned}\phi_1 &= -\frac{1}{2} \sqrt{\frac{3}{10}} \phi + \varphi_1, & \phi_2 &= -\frac{1}{2} \sqrt{\frac{3}{10}} \phi + \varphi_2, & \phi_3 &= -\frac{1}{2} \sqrt{\frac{3}{10}} \phi + \varphi_3, \\ \phi_4 &= -\frac{1}{2} \sqrt{\frac{3}{10}} \phi + \varphi_4, & \phi_5 &= -\frac{1}{2} \sqrt{\frac{3}{10}} \phi - \sum_{k=1}^4 \varphi_k, & \phi_6 &= \frac{1}{2} \sqrt{\frac{5}{6}} \phi + \varphi_5, \\ \phi_7 &= \frac{1}{2} \sqrt{\frac{5}{6}} \phi + \varphi_6, & \phi_8 &= \frac{1}{2} \sqrt{\frac{5}{6}} \phi - \varphi_5 - \varphi_6.\end{aligned}\quad (71)$$

Writing the SO(8) generators as  $T_{IJ} = -T_{JI}$  ( $I, J = 1, \dots, 8$ ), the equations for the scalars  $\varphi_\ell$  ( $\ell = 1, \dots, 6$ ) are satisfied when  $\varphi_\ell \equiv 0$ , and the scalar  $\phi$  enters the kinetic terms of the vector fields as in (18) with  $\nu = 4$  if  $J_1$  and  $J_2$  are chosen as follows:

$$J_1 = \sqrt{\frac{2}{5}} \left( T_{12} + \frac{\varepsilon_1}{\sqrt{2}} T_{34} + \frac{\varepsilon_2}{\sqrt{2}} T_{35} + \frac{\varepsilon_3}{\sqrt{2}} T_{45} \right), \quad J_2 = \frac{1}{\sqrt{3}} (T_{67} + \varepsilon_4 J_{68} + \varepsilon_5 J_{78}), \quad (72)$$

where  $\varepsilon_\ell^2 = 1$ . This identifies the two vector fields  $A_\mu^1$  and  $A_\mu^2$  out of  $A_\mu^{IJ}$ :

$$\frac{1}{2} A_\mu^{IJ} T_{IJ} = A_\mu^1 J_1 + A_\mu^2 J_2, \quad (73)$$

so that the two field strengths  $\bar{F}_{\mu\nu}^1, \bar{F}_{\mu\nu}^2$  in (18) are identified with the  $F_{\mu\nu}^{IJ}$  of the maximal theory as follows:

$$\begin{aligned}F_{\mu\nu}^{12} &= \sqrt{\frac{2}{5}} \bar{F}_{\mu\nu}^1, & F_{\mu\nu}^{34} &= \frac{\varepsilon_1}{\sqrt{5}} \bar{F}_{\mu\nu}^1, & F_{\mu\nu}^{35} &= \frac{\varepsilon_2}{\sqrt{5}} \bar{F}_{\mu\nu}^1, & F_{\mu\nu}^{45} &= \frac{\varepsilon_3}{\sqrt{5}} \bar{F}_{\mu\nu}^1, \\ F_{\mu\nu}^{67} &= \frac{1}{\sqrt{3}} \bar{F}_{\mu\nu}^2, & F_{\mu\nu}^{68} &= \frac{\varepsilon_4}{\sqrt{3}} \bar{F}_{\mu\nu}^2, & F_{\mu\nu}^{78} &= \frac{\varepsilon_5}{\sqrt{3}} \bar{F}_{\mu\nu}^2.\end{aligned}\quad (74)$$

When  $\omega \neq 0$ , the same generators  $T_{IJ}$  are gauged by linear combinations of  $A_\mu^{IJ}$  and  $A_{IJ\mu}$  of the form

$$\hat{A}_\mu^{IJ} = \cos(\omega) A_\mu^{IJ} - \sin(\omega) A_{IJ\mu}, \quad (75)$$

which means that the gauging of SO(8) is performed in a different symplectic frame in which the electric vector fields are  $\hat{A}_\mu^{IJ}$ . Let us denote by  $\hat{A}_\mu^{\hat{M}} = (\hat{A}_\mu^{\hat{\Lambda}}, \hat{A}_{\hat{\Lambda}\mu})$ , where  $\hat{A}_\mu^{\hat{\Lambda}} = \hat{A}_\mu^{IJ}$ , the vector fields and their magnetic duals in the new symplectic frame, and by  $A_\mu^M = (A_\mu^{\hat{\Lambda}}, A_{\hat{\Lambda}\mu})$  the same vectors in the old frame. Let  $\mathbb{F}_{\mu\nu}^{\hat{M}} = \partial_\mu \hat{A}_\nu^{\hat{M}} - \partial_\nu \hat{A}_\mu^{\hat{M}}$  and  $\mathbb{F}_{\mu\nu}^M = \partial_\mu A_\nu^M - \partial_\nu A_\mu^M$  be the corresponding field strengths. We have the following relation:

$$\mathbb{F}_{\mu\nu}^{\hat{M}} = E_M^{\hat{M}} \mathbb{F}_{\mu\nu}^M, \quad E_M^{\hat{M}} = \begin{pmatrix} \cos(\omega) \mathbb{1}_{28 \times 28} & \sin(\omega) \mathbb{1}_{28 \times 28} \\ -\sin(\omega) \mathbb{1}_{28 \times 28} & \cos(\omega) \mathbb{1}_{28 \times 28} \end{pmatrix}. \quad (76)$$

In the new frame, the parameter  $\omega$  will also enter the kinetic matrices  $\mathcal{I}_{\hat{\Lambda}\hat{\Sigma}}(\phi, \omega)$ ,  $\mathcal{R}_{\hat{\Lambda}\hat{\Sigma}}(\phi, \omega)$ , that is, the components of the new symplectic matrix  $\mathcal{M}_{\hat{M}\hat{N}}(\phi, \omega)$ , which is expressed in terms of the  $\omega$ -independent  $\mathcal{M}_{MN}(\phi)$  in the original frame through the relation:

$$\mathcal{M}_{\hat{M}\hat{N}}(\phi, \omega) = E^{-1}(\omega)_{\hat{M}}^M E^{-1}(\omega)_{\hat{N}}^N \mathcal{M}_{MN}(\phi). \quad (77)$$

Upon truncating scalar and vector fields as described above, the two vectors will only enter the bosonic action through the corresponding field strengths. Written in the new symplectic frame, the kinetic terms of  $\hat{A}_\mu^{\hat{\Lambda}} = (\hat{A}_\mu^1, \hat{A}_\mu^2)$  will depend on the  $\omega$  parameter through the restrictions  $\mathcal{I}_{\hat{\Lambda}\hat{\Sigma}}(\phi, \omega)$ ,  $\mathcal{R}_{\hat{\Lambda}\hat{\Sigma}}(\phi, \omega)$  of  $\mathcal{I}_{\hat{\Lambda}\hat{\Sigma}}(\phi, \omega)$ ,  $\mathcal{R}_{\hat{\Lambda}\hat{\Sigma}}(\phi, \omega)$  to the two vectors.

This dependence can, however, be undone at the level of the bosonic field equations and Bianchi identities since the latter depend on  $\mathbb{F}_{\mu\nu}^{\hat{m}} = (F_{\mu\nu}^{\hat{\lambda}}, G_{\hat{\lambda}\mu\nu})$  only in symplectic-invariant contractions with the matrix  $\mathcal{M}_{\hat{m}\hat{n}}(\phi, \omega)$  and its derivatives. This means that the dependence on  $\omega$  of the terms involving the vector field strengths can be disposed of through a redefinition of the latter, which amounts to writing them in terms of the field strengths  $\mathbb{F}_{\mu\nu}^m$  in the original frame (consisting of  $\bar{F}_{\mu\nu}^1, \bar{F}_{\mu\nu}^2$  and their magnetic duals) through the matrix  $E$ . Upon this redefinition, the bosonic field equations of the truncated model coincide with those obtained from the action (18), with  $\nu = 4$ , provided we identify:

$$g = \frac{\sqrt{2}}{L}, \quad \sin(\omega) = L\alpha, \quad \cos(\omega) = \sqrt{1 - L^2\alpha^2}. \quad (78)$$

The embedding of the  $\nu = 4/3$  model in the maximal theory is more subtle and will be dealt with in the future. In the remaining cases  $\nu = \infty$  or  $\nu = \pm 2$ , our solutions can be extended to charged solutions within  $\mathcal{N} = 8$  SO(8) gauged supergravity, within the bosonic part of the STU truncation of it.

**Acknowledgments:** I would like to thank profs. M Trigiante, A Anabalón, D Astefanesei for discussions.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

1. Maldacena, J.M. The Large N limit of superconformal field theories and supergravity. *Int.J.Theor.Phys.* **1999**, *38*, 1113–1133, [[arXiv:hep-th/hep-th/9711200](#)]. doi:10.1023/A:1026654312961.
2. Hawking, S.; Page, D.N. Thermodynamics of Black Holes in anti-De Sitter Space. *Commun. Math. Phys.* **1983**, *87*, 577. doi:10.1007/BF01208266.
3. Chamblin, A.; Emparan, R.; Johnson, C.V.; Myers, R.C. Charged AdS black holes and catastrophic holography. *Phys. Rev. D* **1999**, *60*, 064018, [[hep-th/9902170](#)]. doi:10.1103/PhysRevD.60.064018.
4. Chamblin, A.; Emparan, R.; Johnson, C.V.; Myers, R.C. Holography, thermodynamics and fluctuations of charged AdS black holes. *Phys. Rev. D* **1999**, *60*, 104026, [[hep-th/9904197](#)]. doi:10.1103/PhysRevD.60.104026.
5. Cvetič, M.; Gubser, S.S. Phases of R charged black holes, spinning branes and strongly coupled gauge theories. *JHEP* **1999**, *04*, 024, [[hep-th/9902195](#)]. doi:10.1088/1126-6708/1999/04/024.
6. Caldarelli, M.M.; Cognola, G.; Klemm, D. Thermodynamics of Kerr-Newman-AdS black holes and conformal field theories. *Class. Quant. Grav.* **2000**, *17*, 399–420, [[hep-th/9908022](#)]. doi:10.1088/0264-9381/17/2/310.
7. Strominger, A.; Vafa, C. Microscopic origin of the Bekenstein-Hawking entropy. *Phys.Lett.* **1996**, *B379*, 99–104, [[arXiv:hep-th/hep-th/9601029](#)]. doi:10.1016/0370-2693(96)00345-0.
8. Cacciatori, S.L.; Klemm, D. Supersymmetric AdS(4) black holes and attractors. *JHEP* **2010**, *01*, 085, [[arXiv:hep-th/0911.4926](#)]. doi:10.1007/JHEP01(2010)085.
9. Hristov, K.; Looyestijn, H.; Vandoren, S. BPS black holes in N=2 D=4 gauged supergravities. *JHEP* **2010**, *08*, 103, [[arXiv:hep-th/1005.3650](#)]. doi:10.1007/JHEP08(2010)103.
10. Hristov, K.; Vandoren, S. Static supersymmetric black holes in AdS<sub>4</sub> with spherical symmetry. *JHEP* **2011**, *04*, 047, [[arXiv:hep-th/1012.4314](#)]. doi:10.1007/JHEP04(2011)047.
11. Hristov, K.; Toldo, C.; Vandoren, S. On BPS bounds in D=4 N=2 gauged supergravity. *JHEP* **2011**, *12*, 014, [[arXiv:hep-th/1110.2688](#)]. doi:10.1007/JHEP12(2011)014.
12. Toldo, C.; Vandoren, S. Static nonextremal AdS<sub>4</sub> black hole solutions. *JHEP* **2012**, *09*, 048, [[arXiv:hep-th/1207.3014](#)]. doi:10.1007/JHEP09(2012)048.
13. Chow, D.D.K.; Compère, G. Dyonic AdS black holes in maximal gauged supergravity. *Phys. Rev. D* **2014**, *89*, 065003, [[arXiv:hep-th/1311.1204](#)]. doi:10.1103/PhysRevD.89.065003.
14. Gnechchi, A.; Hristov, K.; Klemm, D.; Toldo, C.; Vaughan, O. Rotating black holes in 4d gauged supergravity. *JHEP* **2014**, *01*, 127, [[arXiv:hep-th/1311.1795](#)]. doi:10.1007/JHEP01(2014)127.
15. Gnechchi, A.; Halmagyi, N. Supersymmetric black holes in AdS<sub>4</sub> from very special geometry. *JHEP* **2014**, *04*, 173, [[arXiv:hep-th/1312.2766](#)]. doi:10.1007/JHEP04(2014)173.
16. Lü, H.; Pang, Y.; Pope, C. An  $\omega$  deformation of gauged STU supergravity. *JHEP* **2014**, *04*, 175, [[arXiv:hep-th/1402.1994](#)]. doi:10.1007/JHEP04(2014)175.
17. Faedo, F.; Klemm, D.; Nozawa, M. Hairy black holes in N = 2 gauged supergravity. *JHEP* **2015**, *11*, 045, [[arXiv:hep-th/1505.02986](#)]. doi:10.1007/JHEP11(2015)045.

18. Klemm, D.; Marrani, A.; Petri, N.; Santoli, C. BPS black holes in a non-homogeneous deformation of the stu model of  $N = 2$ ,  $D = 4$  gauged supergravity. *JHEP* **2015**, *09*, 205, [[arXiv:hep-th/1507.05553](#)]. doi:10.1007/JHEP09(2015)205.
19. Chimento, S.; Klemm, D.; Petri, N. Supersymmetric black holes and attractors in gauged supergravity with hypermultiplets. *JHEP* **2015**, *06*, 150, [[arXiv:hep-th/1503.09055](#)]. doi:10.1007/JHEP06(2015)150.
20. Hristov, K.; Katmadas, S.; Toldo, C. Rotating attractors and BPS black holes in  $AdS_4$ . *JHEP* **2019**, *01*, 199, [[arXiv:hep-th/1811.00292](#)]. doi:10.1007/JHEP01(2019)199.
21. Daniele, N.; Faedo, F.; Klemm, D.; Ramírez, P.F. Rotating black holes in the FI-gauged  $N = 2$ ,  $D = 4$   $\overline{CP}^n$  model. *JHEP* **2019**, *03*, 151, [[arXiv:hep-th/1902.03113](#)]. doi:10.1007/JHEP03(2019)151.
22. Anabalon, A.; Astefanesei, D.; Gallerati, A.; Trigiante, M. New non-extremal and BPS hairy black holes in gauged  $\mathcal{N} = 2$  and  $\mathcal{N} = 8$  supergravity **2020**. [[arXiv:hep-th/2012.09877](#)].
23. Anabalón, A.; Astefanesei, D.; Gallerati, A.; Trigiante, M. Hairy Black Holes and Duality in an Extended Supergravity Model. *JHEP* **2018**, *04*, 058, [[arXiv:hep-th/1712.06971](#)]. doi:10.1007/JHEP04(2018)058.
24. Anabalón, A.; Astefanesei, D.; Choque, D.; Gallerati, A.; Trigiante, M. Exact holographic RG flows in extended SUGRA **2020**. [[arXiv:hep-th/2012.01289](#)].
25. Duff, M.J.; Liu, J.T.; Rahmfeld, J. Four-dimensional string-string-string triality. *Nucl. Phys. B* **1996**, *459*, 125–159, [[hep-th/9508094](#)]. doi:10.1016/0550-3213(95)00555-2.
26. Behrndt, K.; Kallosh, R.; Rahmfeld, J.; Shmakova, M.; Wong, W.K. STU black holes and string triality. *Phys. Rev. D* **1996**, *54*, 6293–6301, [[hep-th/9608059](#)]. doi:10.1103/PhysRevD.54.6293.
27. Behrndt, K.; Lust, D.; Sabra, W.A. Stationary solutions of  $N=2$  supergravity. *Nucl. Phys. B* **1998**, *510*, 264–288, [[hep-th/9705169](#)]. doi:10.1016/S0550-3213(97)00633-0.
28. Duff, M.; Liu, J.T. Anti-de Sitter black holes in gauged  $N = 8$  supergravity. *Nucl. Phys. B* **1999**, *554*, 237–253, [[hep-th/9901149](#)]. doi:10.1016/S0550-3213(99)00299-0.
29. Andrianopoli, L.; D’Auria, R.; Gallerati, A.; Trigiante, M. Extremal Limits of Rotating Black Holes. *JHEP* **2013**, *1305*, 071, [[arXiv:hep-th/1303.1756](#)]. doi:10.1007/JHEP05(2013)071.
30. Andrianopoli, L.; Gallerati, A.; Trigiante, M. On Extremal Limits and Duality Orbits of Stationary Black Holes. *JHEP* **2014**, *01*, 053, [[arXiv:hep-th/1310.7886](#)]. doi:10.1007/JHEP01(2014)053.
31. Henneaux, M.; Martinez, C.; Troncoso, R.; Zanelli, J. Asymptotic behavior and Hamiltonian analysis of anti-de Sitter gravity coupled to scalar fields. *Annals Phys.* **2007**, *322*, 824–848, [[hep-th/0603185](#)]. doi:10.1016/j.aop.2006.05.002.
32. Anabalon, A.; Astefanesei, D.; Martinez, C. Mass of asymptotically anti-de Sitter hairy spacetimes. *Phys. Rev.* **2015**, *D91*, 041501, [[arXiv:hep-th/1407.3296](#)]. doi:10.1103/PhysRevD.91.041501.
33. Anabalon, A.; Astefanesei, D.; Choque, D.; Martinez, C. Trace Anomaly and Counterterms in Designer Gravity. *JHEP* **2016**, *03*, 117, [[arXiv:hep-th/1511.08759](#)]. doi:10.1007/JHEP03(2016)117.
34. Trigiante, M. Gauged Supergravities. *Phys. Rept.* **2017**, *680*, 1–175, [[arXiv:hep-th/1609.09745](#)]. doi:10.1016/j.physrep.2017.03.001.
35. Gallerati, A.; Trigiante, M. Introductory Lectures on Extended Supergravities and Gaugings. *Springer Proc. Phys.* **2016**, *176*, 41–109. doi:10.1007/978-3-319-31352-8\_2.
36. Gallerati, A. Constructing black hole solutions in supergravity theories. *Int. J. Mod. Phys.* **2020**, *A34*, 1930017, [[arXiv:hep-th/1905.04104](#)]. doi:10.1142/S0217751X19300175.
37. de Wit, B.; Nicolai, H.  $N=8$  Supergravity with Local  $SO(8) \times SU(8)$  Invariance. *Phys. Lett. B* **1982**, *108*, 285. doi:10.1016/0370-2693(82)91194-7.
38. de Wit, B.; Nicolai, H.  $N=8$  Supergravity. *Nucl. Phys. B* **1982**, *208*, 323. doi:10.1016/0550-3213(82)90120-1.
39. Cvetic, M.; Gubser, S.; Lu, H.; Pope, C. Symmetric potentials of gauged supergravities in diverse dimensions and Coulomb branch of gauge theories. *Phys. Rev. D* **2000**, *62*, 086003, [[hep-th/9909121](#)]. doi:10.1103/PhysRevD.62.086003.
40. Cvetic, M.; Lu, H.; Pope, C.; Sadrzadeh, A. Consistency of Kaluza-Klein sphere reductions of symmetric potentials. *Phys. Rev. D* **2000**, *62*, 046005, [[hep-th/0002056](#)]. doi:10.1103/PhysRevD.62.046005.
41. de Wit, B.; Nicolai, H. Deformations of gauged  $SO(8)$  supergravity and supergravity in eleven dimensions. *JHEP* **2013**, *05*, 077, [[arXiv:hep-th/1302.6219](#)]. doi:10.1007/JHEP05(2013)077.