

## Eucalyptus biochar as a sustainable nanomaterial for electrochemical sensors



Annalisa Scroccarello<sup>1\*</sup>, Flavio Della Pelle<sup>1</sup>, Qurat Ul Ain Bukhari<sup>1</sup>, Filippo Silveri<sup>1</sup>, Daniele Zappi<sup>2</sup>, Enrico Cozzoni<sup>3</sup>, Dario Compagnone<sup>1</sup>




<sup>1</sup>Faculty of Bioscience and Technology for Food, Agriculture and Environment University of Teramo, Via Renato Balzarini 1, 64100, Teramo, Italy

<sup>2</sup>Biosensor, Via Olmetti 44, 00060, Formello, Roma, Italy

<sup>3</sup>GRADO ZERO INNOVATION, Via nove 2/A, 50056, Montelupo F.no, Firenze, Italy

\*ascroccarello@unite.it

# State of the art



Electrochemical devices obtained from biochar: Advances in renewable and environmentally-friendly technologies applied to analytical chemistry

Leandro S. de Almeida, Eliézer Q. Oreste, Juliana V. Maciel, Mônica G. Heinemann, Daiane Dias\*




Sustainable materials for the design of forefront printed (bio)sensors applied in agrifood sector

Fabiana Arduini <sup>a,b</sup>, Laura Micheli <sup>a</sup>, Viviana Scognamiglio <sup>c</sup>, Vincenzo Mazzaracchio <sup>a</sup>, Danila Moscone <sup>a,\*</sup>



Review  
Biochar: Production, properties and emerging role as a support for enzyme immobilization


Deepshikha Pandey, Achlesh Daverey\*, Kusum Arunachalam\*\*



Application of biochar-based materials in environmental remediation: from multi-level structures to specific devices

Lun Lu<sup>1,2</sup>, Wentao Yu<sup>1,2</sup>, Yaofeng Wang<sup>1,2</sup>, Kun Zhang<sup>1,2</sup>, Xiaomin Zhu<sup>1,2</sup>, Yuecan Zhang<sup>1,2</sup>, Yajing Wu<sup>1,2</sup>, Habib Ullah<sup>1,2</sup>, Xin Xiao<sup>1,2</sup>, Baoliang Chen<sup>1,2</sup>

Received: 1 January 2020 / Accepted: 25 February 2020




★ Ramanayaka, S., Vithanage, M., Alessi, D. S., Liu, W. J., Jayasundera, A. C., & Ok, Y. S. (2020). *Environmental Science: Nano*, 7(11), 3279-3302.


★ Xiao, X., Chen, B., Chen, Z., Zhu, L., & Schnoor, J. L. (2018). *Environmental science & technology*, 52(9), 5027-5047.

# Biochar preparation, electrode set-up and fabrication




## Preparation of Biochar dispersion through liquid-phase sonochemical approach







## Biochar-based electrode assembling strategies



# Biochar as sensing element: amount optimization



**BH-Film**




CVs: 5 mM  $\text{Fe}(\text{CN})_6^{4-/-3-}$  in 0.1M KCl; scan rate 25 mV s<sup>-1</sup>.

EIS: 5 mM  $\text{Fe}(\text{CN})_6^{4-/-3-}$  in 0.1M KCl; frequency range of  $10^5$  -  $10^{-1}$  Hz using 5mV as sinusoidal wave amplitude, fixing the potential at open circuit.


# Biochar as sensing element: mono- and ortho-phenols evaluation

## Explorative cyclic voltammetries


Ortho-phenol







Mono-phenol



Bare electrode  
(black line)



BH-SPE  
(red line)




# Biochar as sensing element: mono- and ortho-phenols evaluation

## Analysis of Tyrosol and Hydroxytyrosol in extra virgin olive oil


**BH-SPE**




Dose-response curve




EVOO Sample analysis



Recovery & LOD







HTR : 97 - 108 %  
LOD : 0.5 μM




TR : 91 - 111 %  
LOD : 3.0 μM

**BH-Film**



HTR : 96 - 106 %  
LOD : 0.2 μM



TR : 95 - 110 %  
LOD : 0.6 μM

- ✓ A water-soluble Biochar nanofiber dispersion was obtained
- ✓ The water-soluble biochar was employed for sensors realization
- ✓ The Biochar nanofiber was used as SPE electrode modifier and to realize biochar-based conductive film
- ✓ The conductive film was integrated into an entirely lab-made flexible electrode
- ✓ The Biochar-based electrodes present great analytical performance for ortho-phenols compounds
- ✓ The Biochar-based electrode allow the selective detection of ortho-phenols in presence of mono-phenols compounds
- ✓ Sensors showed useful linear ranges and limits of detections
- ✓ The sensors were applied in EVOO samples analysis and quantitative recoveries were obtained

# Acknowledgments



Prof. Dario Compagnone

Dr. Flavio Della Pelle

Ph.D. Filippo Silveri

Ph.D. Qurat Ul Ain Bukhari



Dr. Maria Teresa Giardi

Dr. Daniele Zappi



Dr. Enrico Cozzoni

*... THANKS FOR  
YOUR ATTENTION!*

