

# Systematic review: Antioxidant and neuroprotective capacity of species of the genus *Asplenium* (Monilophyta: Aspleniaceae) †

Mathias Flores-González, Mario Simirgiotis and Alfredo Torres-Benítez \*

Instituto de Farmacia, Facultad de Ciencias, Universidad Austral de Chile, 5090000 Valdivia, Chile; mathy.power.96@gmail.com (M.F.-G.); mario.simirgiotis@gmail.com (M.S.)

\* Correspondence: aljotobe19@hotmail.com; Tel.: +59-9-9054-9002

† Presented at the 2<sup>nd</sup> International Electronic Conference on Brain Sciences, 15-30 July 2021; Available online: <https://iecbcs2021.sciforum.net>

**Abstract:** The genus *Asplenium* L. comprises about 700 species of terrestrial, epiphytic, and saxicolous habits distributed in temperate and tropical regions around the world, exhibiting a high chemical richness with variable biological activity. In this review, compounds with antioxidant activity that constitute a pharmacological potential in diseases of the central nervous system are detailed. *Asplenium nidus* L. species presents a high concentration of phenols and flavonoids evidenced by antioxidant activity assays such as DPPH, FRAP, total phenols, total flavonoids, and ORAC, and represent compounds with bioactive potential including neuroprotection. The species *Asplenium adiantum-nigrum* L. presents a high antioxidant potential of its rhizomes extracts exhibited in DPPH and ABTS assays, attributed to the high concentration of mangiferin; the xanthone mangiferin is a compound also present in other species of the genus such as *Asplenium ceterach* L. and *Asplenium montanum* Willd. in significant amounts. This xanthone has studies on its neuroprotective effect through different targets, some of them being the acetylcholinesterase enzymes, the 5-lipoxygenase enzyme, and the antioxidant activity itself. All these ways of action of mangiferin constitute an object of study for its effect on memory loss, which can be delayed in Alzheimer's disease.

**Keywords:** *Asplenium*; bioactive compounds; antioxidant capacity; neuroprotection

## 1. Introduction

Ancestral ferns have been widely used in many cultures as phytotherapeutic agents, covering the treatment of diverse and varied conditions, from infections of pathogens to combat kidney stones, asthma, stomach pains, and also as anti-inflammatory agents [1]. This in turn has prompted the emergence of multiple studies around the world to identify the components responsible for the biological activity of these plants.

*Asplenium* L. is a genus that has about 700 species of terrestrial, epiphytic, and saxicolous habitats distributed in temperate and tropical regions in the world [1]. All this diversity in species also translates into a great diversity of compounds such as flavonoids, which are a group of bioactive compounds, being among the most common that can be found in this genus. All this has aroused the interest of part of the scientific research, due to the phytotherapeutic potential of these compounds, such as the effect of antioxidant activity and neuroprotective effect [2].

Among the variety of compounds identified in the genus, *Asplenium* L. one can name the natural C-glucoside mangiferin (2-C-β-D-gluco-pyranosyl-1,3,6, 7-tetrahydroxyxanthone; C<sub>19</sub>H<sub>18</sub>O) a type of xanthone that has been detected in the species *Asplenium balearicum* Shivas, *Asplenium bradleyi* D.C. Eaton, *Asplenium pinnatifidum* Nutt., *Asplenium × stotleri* Wherry (pro. sp.), *Asplenium serra* Langsd. & Fisch, *Asplenium ceterach* L. and *Asplenium montanum* Willd. [1]. The importance of this compound lies in its potential for the treatment of neurodegenerative diseases in cellular and animal models, together with its

**Citation:** Lastname, F.; Lastname, F.; Lastname, F. Title. *Proceedings* **2021**, *68*, x. <https://doi.org/10.3390/xxxxx>

Published: date

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

high antioxidant activity and the ability to scavenge hydroxyl radicals. In addition, it reports an important effect on  $\beta$ -amyloid peptide activity and the ability to inhibit acetylcholinesterase (AChE) and lipoxygenase (LOX) enzymes [3].

## 2. Methods

A literature search was performed until March 2021 in the databases WoS, Scopus, ScienceDirect, and PubMed, using the keywords: Asplenium, antioxidant, neurodegenerative diseases, neuroprotection. References from relevant review articles were also identified, as well as others registered in the section of similar articles in PubMed.

## 3. Results and Discussion


Methanolic extracts of *Asplenium adiantum-nigrum* L. and *Asplenium ruta-muraria* L. evaluated by UHPLC/DAD/-HESI-MS/MS have shown different phenolic compounds, among which are phenolic acids and derivatives, flavonoids and derivatives, and xanthonenes. Table 1 shows the concentrations of the most relevant compounds present in three plant organs sampled for each species (rhizome, fronds, roots), expressed in milligrams per gram of fresh weight (FW) [1].

**Table 1.** Compounds present in the vegetative organs of *A. adiantum-nigrum* and *A. ruta-muraria*.

Compounds	Concentration (mg/g FW)		
	<i>A. adiantum-nigrum</i>	<i>A. ruta-muraria</i>	
Phenolic acids and derivatives	Gallic acid	0.69±0.01 (Rhizome)	-
	Protocatechuic acid	1.61±0.06 (Rhizome)	1.29±0.01 (Rhizome, Roots) 1.29±0.04 (Fronds)
	Gentisic acid	0.13±0.08 (Fronds)	0.11±0.01 (Rhizome, Roots), 0.11±0.06 (Fronds)
	p-Hydroxybenzoic acid	0.63±0.05 (Fronds)	0.44±0.11 (Fronds)
	Aesculin	0.39±0.05 (Fronds)	0.51±0.03 (Fronds)
	Chlorogenic acid	0.62±0.03 (Rhizome)	2.05±1.05 (Roots)
	Caffeic acid	0.59±0.06 (Rhizome)	0.21±0.02 (Fronds)
	p-Coumaric acid	2.82±0.74 (Rhizome)	1.00±0.08 (Fronds)
	Ferulic acid	0.23±0.01 (Rhizome)	-
	Rosmarinic acid	1.69±0.01 (Roots)	0.03±0.01 (Fronds)
	Flavonoids and derivatives	Epigallocatechin	3.71±1.33 (Rhizome)
Epicatechin		-	1.26±0.01 (Rhizome)
Gallocatechin gallate		-	10.29±0.67 (Fronds)
Epigallocatechin gallate		1.15±0.09 (Fronds)	1.74±0.18 (Fronds)
Xanthonenes	Rutin	0.27±0.01 (Roots)	0.24±0.01 (Roots)
	Mangiferin glucoside	507.69±19.58 (Rhizome)	0.51±0.01 (Rhizome)
	Mangiferin	598.29±20.82 (Rhizome)	0.70±0.12 (Rhizome)

It should be noted that xanthonenes represent the most abundant compounds, especially in *A. adiantum-nigrum*, with concentrations greater than 500 (mg/g FW) [1].

Mangiferin, a compound present in a large number of species of the genus *Asplenium* [1] (Figure 1), possesses qualities that make it a potential candidate for use in phytotherapy. Its performance in AChE and LOX enzyme inhibition tests, antioxidant activity by the DPPH method, and its action on the activity of  $\beta$ -amyloid peptides in neurons in vitro are presented below.



**Figure 1.** Chemical structure of mangiferin ( $C_{19}H_{18}O_{11}$ ).

The inhibition produced on the AChE enzyme, by the compound mangiferin, is quite significant obtaining a value for  $IC_{50}$  of  $55.42 \pm 1.52$   $\mu\text{g/ml}$  in comparison with galantamine, a compound used as a standard in AChE assays ( $IC_{50}$ :  $5.543 \pm 0.213$   $\mu\text{g/ml}$ ). Therefore, its action is considered potential to prevent the loss of the neurotransmitter acetylcholine in the progression of Alzheimer's disease [3].

On LOX enzyme, mangiferin has presented inhibition results for  $IC_{50}$  of  $42.28 \pm 0.63$   $\mu\text{g/ml}$ , which is significant, compared to the routine standard ( $IC_{50}$ :  $7.46 \pm 0.18$   $\mu\text{g/ml}$ ). It should be noted that LOX is a key enzyme in inflammatory processes as responsible for the production of leukotrienes and other inflammatory mediators [3].

As an antioxidant agent, mangiferin has shown an  $IC_{50}$  value in the DPPH method of  $18.50 \pm 0.18$   $\mu\text{g/ml}$  [3]. This activity corroborates the neuroprotective action of this compound, through the pathway of reducing  $H_2O_2$ -induced oxidative damage [4].

Mangiferin has demonstrated a concentration-dependent activity to have a reduction effect on  $\beta$ -amyloid peptide deposits, with significant results at concentrations of 10  $\mu\text{g/mL}$  and maximal at 20  $\mu\text{g/mL}$  in vitro tests [3]. The  $\beta$ -amyloid peptide is responsible for toxic effects to neurons in the course of Alzheimer's disease, in addition to its action in ROS production that increases neurotoxicity [5].

In general, the flavonoid profiles present in the species of the Aspleniaceae family are complex to characterize and differentiate. Regarding flavonols, kaempferol derivatives such as 3,7-glycosides mainly as well as 3,4'-diglycosides and 3,7,4'-triglycosides predominate; in *Asplenium sulcatum* Lam. the presence of O-methylated kaempferol glycosides has been reported. On the other hand, C-glycoside flavones are represented by luteolin and apigenin, and C-sugars may occasionally occur O-glycosides; O-glycoside flavones have been found in three *Asplenium* species, including scutellarein 6-O-glucoside in *Asplenium belangeri* Bory [6, 7].

#### 4. Conclusions

Species of the genus *Asplenium* present bioactive compounds of the polyphenolic type that stand out for their significant antioxidant activity and potential to reduce causes of neurotoxicity in Alzheimer's disease such as amyloid plaques and oxidative stress. In addition, their metabolomic profiles contribute to taxonomic separation with other groups of ferns with which they share morphological similarities.

**Author Contributions:** All authors contributed equally to this work. All authors discussed the results and implications and commented on the manuscript at all stages. A.T.-B. and M.F.-G. wrote the paper. All authors have read and agreed to the published version of the manuscript.

**Funding:** Please add: This research was funded by CONICYT PFCHA/National Doctorate Scholarship/2357-2021 (A-T.-B.).

**Acknowledgments:** To the laboratory of Natural Products of the Institute of Pharmacy of the Universidad Austral de Chile.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Živković, S.; Milutinović, M.; Maksimović, V.; Ćirić, A.; Ivanov, M.; Božunović, J.; et al. Antioxidant and antimicrobial activity of two *Asplenium* species. *South African Journal of Botany* **2020**, *132*, 180-187.
2. Jarial, R.; Thakur, S.; Sakinah, M.; Zularisam, A.W.; Sharad, A.; Kanwar, S.S.; Singh, L. Potent anticancer, antioxidant and antibacterial activities of isolated flavonoids from *Asplenium nidus*. *Journal of King Saud University-Science* **2018**, *30*, 185-192.
3. Sethiya, N.K.; Mishra, S.H. Investigation of mangiferin, as a promising natural polyphenol xanthone on multiple targets of Alzheimer's disease. *Journal of Biologically Active Products from Nature* **2014**, *4*, 111-119.
4. Kawpoomhae, K.; Sukma, M.; Ngawhirunpat, T.; Opanasopit, P.; Sripattanaporn, A. Antioxidant and neuroprotective effects of standardized extracts of *Mangifera indica* leaf. *Thai Journal of Pharmaceutical Sciences* **2010**, *34*, 32-43.
5. Cheignon, C.; Tomas, M.; Bonnefont-Rousselot, D.; Faller, P.; Hureau, C.; Collin, F. Oxidative stress and the amyloid beta peptide in Alzheimer's disease *Redox biology* **2018**, *14*, 450-464.
6. Cao, H.; Chai, T.T.; Wang, X.; Morais-Braga, M.; Yang, J.H.; Wong, F.C.; et al. Phytochemicals from fern species: potential for medicine applications. *Phytochemistry Reviews* **2017**, *16*, 379-440.
7. Umikalsom, Y.; Grayer-Barkmeijer, R.J.; Harborne, J.B. A comparison of the flavonoids in Athyriaceae and Aspleniaceae. *Biochemical Systematics and Ecology* **1994**, *22*, 587-594.