

Proceedings

Preferential use of Bamboos for Industrial Production of Incense sticks

George Jenner V^{1*}, Thiru Selvan² and Kumaresan P³¹Tripura University; vgjenner@gmail.com,²Tripura University; tselvan@tripurauniv.ac.in,³Tamilnadu Agricultural University; acfkumagri@gmail.com* Correspondence: vgjenner@gmail.com; Tel.: (+919436135005)

Abstract: The incense stick or agarbatti stick is used exclusively for religious purposes in almost every home in India, but now it is being used worldwide for its medicinal values. Bamboo, popularly known as green gold or poor man's timber, is a multipurpose, fast-growing woody species, which occupies an essential place in the people's diverse phases of life and culture. The primary concern of the different stakeholders of the bamboo sector is identifying the right choice of raw material to maximize incense sticks' production. On the other hand, the market functionaries take advantage of farmers' lack of knowledge of the demand-supply situation and make substantial margin money. Therefore, the paper evaluates the preference of the notable species, age, and the part of the culm of bamboo used for the industrial production of incense sticks. In Tripura, India, the investigation indicated that all study parameters were significantly different among the species *viz.*, *Bambusa polymorpha*, *B. vulgaris*, *B. cacharensis*, *B. tulda*, *B. balcooa*, *M. baccifera*, *Dendrocalamus asper*, and *D. longispachus*. Among the bamboo species, *B. tulda* is the most preferred one having a mean Likert scale of 4.89 followed by *Dendrocalamus longispachus* (4.06), *B. cacharensis* (3.54), *B. polymorpha* (3.50), and so on. The most preferred age of bamboo culm is three-year old culm (4.32), followed by four and two year old culms. Therefore, it can be concluded that middle portion of the third year old culms of *B. tulda* should be preferably harvested to become the raw material for agarbatti industry for better outturn and profit.

Citation: Jenner V, G.; Kumaresan P3 Preferential use of Bamboos for Industrial Production of Incense sticks. 2021, volume number, x. <https://doi.org/10.3390/xxxxx>

Received: date

Accepted: date

Published: date

Keywords: Bamboo raw material; Outturn; Incense Stick; Likert scale; Index score

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The incense stick is also known as "agarbatti" in India, "joss stick" in China, and "incense" stick in other countries, appears to be an integral part of the worship of all cultures and is used for therapeutic purposes, as well as mosquito repellents and fresheners [1]. Incense sticks are long, cylindrical structures, having a bamboo core coated with fragrant materials, burnt during religious ceremonies, fills the atmosphere with fresh air and aroma eliminate negative energy from our surroundings [2,3,4]. Bamboo is an important raw material in the agarbatti industry [5]. They are fast-growing, widely present, renewable, versatile, and a low-cost natural resource, due to which they are aptly known as 'green gold' and 'poor man's timber' [6]. More than 1600 species of bamboo are distributed in the tropical and sub-tropical regions of the world [7]. As per [8], India has 125 indigenous and 11 exotic bamboos belonging to 23 genera. Four species of bamboo *viz.* Barak (*Bambusa balcooa*), Bari (*Bambusa vulgaris*), Mirtinga (*Bambusa tulda*), and Muli (*Melocanna baccifera*) are used in bamboo-based incense sticks [9]. The durability and strength of bamboo are due to the physico-chemical characteristics of bamboo culms, which determines

its end uses [10]. Bamboo constitutes around 50% parenchyma, 40% fibers, and 10% vessels and sieve tubes [11] with microscopic structures of the bamboo fibers consisting of cellulose, hemicellulose, lignin, and pectin [12] thus showing remarkable variation both between and within species [13]. Bamboo culms have similar chemical constitutions to that of wood except for its high alkaline extract content, ash and silica contents [14] with lignin comparable to that of softwood and hardwood [15,16]. The scarcity of raw materials due to flowering [17], rapid depletion of bamboo resources [18], and increasing popularity of bamboo in the industrial sector as an alternative to wood [17] have resulted in a shortage of the species mainly used by agarbatti industry. Bamboo incense stick production plays a vital role in the rural subsistence economy of the household income basket [19] by providing livelihood and enhancing income levels of women [20]. Indian agarbatti has a high demand both in the local and international markets [21]. India is presently one of the largest producers and exporters of incense sticks with global domination by countries like the U.S.A., Brazil, and China [22]. India's incense stick market is likely to reach Rs. 7500-8000 crore with its exports to more than 150 countries [23]. Since there is no study on the choice of appropriate bamboo species, suitable age, and position of the culm the present research evaluates the industrial production of incense sticks.

2. Methodology

2.1. Study area

The study was undertaken in one of the North-Eastern states of India, Tripura, having an area of 10,491 km². It lies between 22°56' and 24°32' N latitude and 91°09' and 92°22' E longitude. For evaluating the preferred bamboo species, preferred age, and a desired portion of the culm, a questionnaire was developed using the *Likert scale* (1932) to test among 104 respondents, including entrepreneurs, managers, supervisors, workers, and academicians. The majority of respondents were women (59.6%); who were stick makers and artisans (45.2%), aged between 22 and 64 years, with educational qualification of secondary school (49%) working in private industries (83.7%). Respondents were invited to define their attitude to the statement by choosing many *n* grades. In this paper, *n*=5, which denotes five scales viz., strongly disagree, disagree, neither agree nor disagree (neutral), agree, and strongly agree.

$$X_i^{(\rho)} = \sum_{j=m_\rho}^{m_{\rho+1}-1} X_{ij} \quad \rho=1, \dots, v$$

$$X_i = \sum_{\rho=1}^v X_i^{(\rho)} = \sum_{j=1}^M X_{ij}$$

The above equation implies the response vector of the respondent *i* in the dimension ρ , respectively, the total response vectors in all *M* items. Likert scale analysis was performed using MS-Excel to identify the preferred bamboo species among the eight bamboo species (*viz.*, *Bambusa polymorpha*, *B. vulgaris*, *B. cacharensis*, *M. baccifera*, *B. tulda*, *Dendrocalamus asper*, *Dendrocalamus longispathus* and *Bambusa balcooa*), preferred age, and portion of the culm.

3. Results and Discussion

3.1. Preferred Bamboo Species

The grades for the Likert strongly disagree to strongly agree were assigned as 1 to 5, respectively. The analyzed data indicated that the most preferred bamboo species for industrial incense sticks was *Bambusa tulda*, with a Likert score of 4.88 out of five. The least preferred bamboo species was *Dendrocalamus asper* with a Likert score of 2.89 out of five (Table 1). Many bamboo species exist, but only six species *viz.* *B. balcooa*, *B. vulgaris*, *B. tulda*, *B. nutans*, *B. bambos*, and *Melocanna baccifera* are presently used in bamboo-based incense sticks [9, 24, 25]. [24] in its operational guidelines have mentioned *B tulda* as the best-suited species for the Indian Agarbatti industry. *B. tulda* shows higher bulk density,


basic density and lesser moisture content [26]. [27] mentions the fibre dimensions of the culm being- length 1.45-3.0 mm, diameter 15-20 µm, lumen diameter 5-5.6 µm, wall thickness 3.2-7.5 µm with approximate chemical composition of holocellulose 64%, pentosans 18%, lignin 25%, and ash 2-3% which may be the reason for its preference to other species. It is one of the most grown/planted species by the rural people because of its clean and straight culm, medium height, strength and easy availability [28].

Table 1. Scores and ranking of preferential attributes from questionnaire survey for choice of Bamboo species used in incense stick industry.

Parameters	<i>B. polymorpha</i>	<i>B. tulda</i>	<i>B. vulgaris</i>	<i>B. balcooa</i>	<i>M. baccifera</i>	<i>D. longispachus</i>	<i>B. cacharensis</i>	<i>D. aspera</i>
Ease in availability/supply	3.40	4.92	3.29	2.56	3.75	4.03	3.42	2.80
Price of culm	3.40	4.83	3.30	2.15	3.62	4.00	3.46	2.83
Nodal length	3.51	4.92	3.43	2.44	3.56	4.13	3.61	2.89
Wall thickness	3.53	4.91	3.48	3.05	3.16	4.05	3.62	2.97
Outturn/max production	3.50	4.92	3.44	2.81	3.27	4.05	3.53	2.94
Number of sticks per kg	3.53	4.91	3.37	2.84	3.38	4.03	3.52	2.90
Smoothness of the stick	3.58	4.79	3.35	2.66	3.64	4.11	3.59	2.91
Overall mean	3.49	4.89	3.38	2.64	3.48	4.05	3.53	2.89
Rank	4	1	6	8	5	2	3	7

The easy availability or supply, intermodal length of the culm, and the productivity or outturn of agarbatti sticks of *B. tulda* contributed to being the best raw material for the agarbatti industry. The percentage contribution (Fig-1) of all the attributes viz., the availability, cost, intermodal length and wall thickness of the culm, the stick outturn, number of sticks per kilogram, and the sticks' smoothness were almost identical, varying from 14% to 15%. The availability/supply and the intermodal length [29,30] contributed 15 percent each.

Figure 1. Percent contribution of different attributes for the choice of species for the industrial incense stick.



3.2. Preferred age of the Bamboo Culm


The respondent data revealed that the most preferred age of the culm for the production of industrial round sticks for Agarbatti is three-year-old culm with a Likert score of 4.30 out of five, and the least preferred age of the culm is one-year-old culm with a Likert score of 2.42 out of five (Table 2). [31,32,33] reported that selective cutting of mature bamboo culms of more than three years of age in plots with a three-four year rotation cycle appears to be sustainable and more productive. The strength of bamboo increases as it becomes older due to the hardening of the culm walls. [34] concluded that with age increment, mature tissues start to develop and continue to influence density, strength properties, growth of branches, and established root system in one to three-year-old Malaysian bamboos. He reiterated that as bamboo matures, the culm wall thickness becomes hard, resulting in maximum strength. Bamboo matures in about three years; it reaches its full strength [35,36,37].

Table 2. Scores and ranking of preferential attributes from questionnaire survey for choice of age of culm used in incense stick industry.

Parameters	One year	Two year	Three year	Four year	Five year and above
Ease in availability/supply	2.34	3.83	4.23	3.75	3.66
Price of culm	2.62	3.91	4.27	3.88	3.76
Internodal length	2.45	3.84	4.38	3.91	3.77
Wall thickness	2.36	3.81	4.34	3.88	3.75
Outturn/max production	2.33	3.75	4.31	3.84	3.69
Number of sticks per kg	2.42	3.82	4.34	3.85	3.70
Smoothness of the stick	2.50	3.84	4.30	3.94	3.80
Overall mean	2.43	3.83	4.31	3.86	3.73
Rank	5	3	1	2	4

The wall thickness and the strength properties have contributed to being the most preferred age for industrial incense sticks. The percentage contribution (Fig-2) of all the attributes viz., the availability, cost, intermodal length and wall thickness of the culm, the stick outturn, number of sticks per kilogram, and the sticks' smoothness were almost the same, varying from 14% to 15%. The wall thickness and the intermodal length [29,30] contributed 15 percent each.

Figure 2. Percent contribution of different attributes for the choice of the age of culm for the industrial incense stick.



3.3. Preferred Portion of the culm


The analyzed data revealed that the most preferred portion of the bamboo culm for the production of incense sticks is the middle portion of the culm, followed by the bottom part (Table 3). [29,38] reported increasing specific gravity or density with increasing

height levels of the culm. [29] studied the variation in density of selected Philippine bamboos *B. blumeana* and *Gigantochloa levis*. He observed a significant increase in density from base to top height levels in a culm, where the density value increased toward the top for both species. The internodal length and the physical properties of the middle portion of the culm contributed to the reasons for the preferred portion of the culm for industrial incense sticks. The percentage contribution (Fig-3) of all the attributes viz., the availability, cost, internodal length and wall thickness of the culm, the stick outturn, number of sticks per kilogram, and the sticks' smoothness were almost the same, varying from 14% to 15%. The number of sticks per kilogram and the internodal length [29,30] contributed 15 percent each.

Table 3. Scores and ranking of preferential attributes from questionnaire survey for choice of a portion of the culm used in incense stick industry.

Parameters	Bottom	Middle	Top
Ease in availability/supply	4.09	4.33	3.23
Price of culm	4.09	4.52	3.37
Internodal length	3.91	4.65	3.23
Wall thickness	4.19	4.63	2.74
Outturn/max production	4.20	4.64	2.78
Number of sticks per kg	4.05	4.64	2.89
Smoothness of the stick	4.12	4.46	3.34
Overall mean	4.09	4.55	3.08
Rank	2	1	3

Figure 3. Percent contribution of different attributes for the choice of portion of culm for the industrial incense stick.



4. Conclusion

From the present research, it can be concluded that the most preferred raw material for the industrial production of agarbatti sticks is the middle portion of the three-year-old culm of *B.tulda*. The variation in internodal length, wall thickness, and physical properties could be the factors for the above. Further studies are required to find out the best species in terms of outturn, preferred physical and mechanical properties in demand at the national level to make the agarbatti sector more profitable.

Acknowledgments: The authors sincerely thank all the respondents viz., the managers, supervisors, workers, entrepreneurs, and academicians who had participated as respondents in the survey. Our sincere gratitude also to the Department of Forestry and Biodiversity, Tripura University, for providing the opportunity to take up this work.

Conflicts of Interest: There is no conflict of interest

References

1. See, S.W; Rajasekhar, B; Umid Man Joshi. Physical characteristics of nanoparticles emitted from incense smoke Science and Technology of Advanced Materials. **2007**, 825–32
2. Lin, T; Krishnaswamy, C.; Chi, D. S. Incense smoke: clinical, structural and molecular effects on airway disease. Clinical and molecular allergy: **2008**, C.M.A., 6, 3. <https://doi.org/10.1186/1476-7961-6-3>.
3. Hyams, G.;Cushner S. Incense: Rituals, Mystery, Lore. Chronicle Books. 2004. ISBN 0-81183993-1
4. Jetter, J. J;Guo, Z; McBrian, J. A. Flynn, M. R. Characterization of emissions from burning incense. *Science of The Total Environment*,**2002**, 295(1-3), 51–67. doi:10.1016/s0048- 9697(02)00043-8
5. Dutta, P. Agarbatti: A bamboo-based Enterprise – Retrospect and Prospect in Tripura, In: K. S. Chakraborty (Ed.) *Entrepreneurship and Small BusinessDevelopment*, 2006, Mittal Publications, New Delhi, India. 239-240.
6. Tripathi, H. K; Somasundaram, P; Rajesh Kumar; Prakash Lakchaura., Assessment of Bamboo Resources of India, Bamboos in India. **2015**, pp. 15-26
7. Vorontsova, M. S; Clark, L. G; Dransfield, J; Govaerts, R; Baker; W. J; World checklist of bamboos and rattans. INBAR Technical Report No. 37. International Network of Bamboo & Rattan, Beijing, China. **2016**,pp. 466.
8. FSI, State of Forest Report- 2019, Available at: <http://www.fsi.nic.in/forest-report-2019>
9. Jenner, V. G.; Selvan, T. Prospects of Bamboo in Livelihood Enhancement of Rural Women through Incense Stick and its Role in Mitigating Rural Poverty- A model. International Forestry Working Group Newsletter- Working Group B3-Society of American Foresters. **2020**, 8(3): 17-22.
10. Guleria, V; Choudhary, P; Vashisth, A; Gupta, A; Selvan, T. Studies on the physical characteristics, properties, and productivity potential of exotic and indigenous bamboos in rainfed subtropical environment *Journal of Pharmacognosy and Phytochemistry*. **2020**,9(3): 1552-1550
11. Liese W. Research on Bamboo **1987** Wood Sci. Technol., 21, no. 3, 189-209.
12. Dessalegn Y; Singh B. Effect of Analysis on Morphology of Bamboo Fiber: A Review. International Journal of Engineering Sciences & Research Technology **2019** 8(3), 7-17.
13. Pande P.K. Variation in fibre dimensions of some commercially important bamboo species of India. **2009** Journal of the Indian Academy of Wood Science, 6(1-2): 7-11.
14. Liese W. Anatomy of bamboo. 161- 164. In: Proc. Workshop on Bamboo Research in Asia. Singapore. **1980** (Eds.) G. Lessard and A. Chouinard. IDRC, Ottawa, Canada.
15. Fengel, D; Wegener G. Wood: chemistry, ultrastructure, reactions. **1984** Berlin: Walter de Gruyter Publishers. pp. 613.
16. Dence C.W. The determination of lignin. In: (S.Y. Lin and C.W. Dence, eds.). Methods in lignin chemistry. Springer-verlag: **1992** Heidelberg. pp. 33-61.
17. Selvan, T; Bamboo Resources, Their Status, Conservation and Strategies for Improvement In: Forest, Climate Change and Biodiversity, **2018**, K.K. Sood & Mahajan, V (Eds.) Kalyani Publishers New Delhi, India. pp. 263-286.
18. Singh, N. B.; Sarswat, C. V.; Selvan, T; Genetic Improvement of Bamboos in India: Present Status and Future Strategies. In: *Advances in Hort. and Forestry*,**1999**, Singh, S. P. (Ed.) 6: 193-231.
19. Reza S. Bamboopreneur. **2021**,ISBN 978 1-63669-728-4 P.51
20. Rao, I. R.; Kumar, A.; Reza, S.; Motukuri, B. A pathway out of poverty. Bamboo Incense Sticks Production as a Livelihood Option for Rural Women in Tripura, India. International Network for Bamboo and Rattan (INBAR) and Beijing and Centre for Indian Bamboo Resource and Technology (CIBART), **2009**,New Delhi, pp. 7-39.
21. Hazarika, P; Nibedita, B; Dutta, S.C; Biswas, R.C; Dutta; Jayaraj. R.S.C. Status of Agarbatti Industry in India with Special Reference to Northeast. *Int. J. Adv. Res. Biol. Sci.***2018**,5(1): 173-186. DOI: <http://dx.doi.org/10.22192/ijarbs.2018.05.01.024>
22. Yadav, V.K.; Kumar, P.; Kalasariya, H.; Choudhary, N.; Singh, B.; Gnanamoorthy, G.; Gupta, N.; Khan, S.H.; Khayal, A. The Current Scenario of Indian Incense Sticks Market and Their Impact on the Indian Economy, *Ind. J. Pure App. Biosci.***2020**,8(3), 627-636. DOI: <http://dx.doi.org/10.18782/2582-2845.816>
23. Agarwal, P. **India's self-reliant incense industry has wide global appeal; S.M.B. story, 2021. Your story.com Accessed on 10th May, 2021.**
23. MSME, Operational Guidelines for the Pilot Projects of Agarbatti Industry under polymer and chemical-based industry (PCBI) Vertical of Gramodyog Vikas Yojana (G.V.Y.). **2020**, <http://dcmsme.gov.in/Agarbatti%20making%20Project.pdf>.
24. N.B.M. Consultation on opportunities and Challenges for Bamboo in India. (Concept Note), National Bamboo Mission, **2021**, Department of Agriculture, Cooperation and Farmers Welfare, Government of India.
25. Thirunirai Selvan R; Parthiban K.T; Umesh Khanna S. Physio-Chemical Properties of Bamboo Genetic Resources at Various Age Gradations **2017** Int. J. Curr. Microbiol. App. Sci. 6(9), 1671-1681
26. Duriyaprapan, S., Jansen, P.C.M. Bambusa tulda Roxb. In: Dransfield, S, Widjaja, E.A. (Eds.) Plant Resources of South-East Asia No. 7. **1995**. Bamboos. Backhuyes, Leiden, the Netherlands, pp. 69–72
27. Singha Bebija L; Hassan Y. Analysis of Carbohydrate, Moisture Content and Specific Gravity of *Bambusa tulda* with Special Reference to its Harvesting Season **2017** International Journal of Science and Research (IJSR) 6 (1), 279-283.

28. Espiloy, Z.B. Physico-mechanical properties and anatomical relationships of some Philippine bamboos. In: Rao, A.N. *et al.*, (E.D.s). Recent Research on Bamboo. Proceedings of an International Workshop. 16-14 October **1985**, Hangzhou. pp. 257-264.
29. Yan, C.K. Joss Stick Manufacturing: A study of traditional Industry in Hong Kong, Journal of the Royal Asiatic Society of Hong Kong Branch, **1989**, 29, 94-120.
30. Chaturvedi, N.A. Management of bamboo forests. In Bamboos: Current Research (eds I.V.R. Roa, R. Gnanaharan, B.C. Sastry), **1988**, pp. 80-82. Proceedings of the International Bamboo Work-shop held in Cochin, India from 14-18 November, Kerala Forest Research Institute, India and International Development Research Center, Canada.
31. Prasad, R. Scope for change in the management of natural bamboo stands. In Bamboos: Current Research (eds I.V.R. Roa, R. Gnanaharan, & B.C. Sastry), pp. 76-82. Proceedings of the International Bamboo Workshop held in Cochin, India from 14-18 November **1988**, Kerala Forest Research Institute, India and International Development Research Center, Canada.
32. Suwannapinunt, W. Horse-shoe harvesting trials in natural *Gigantochloa hasskarliana* stands. In Bamboos: Current Research (eds I.V.R. Roa, R. Gnanaharan, & B.C. Sastry), pp. 83-86. *Proceedings of the International Bamboo Workshop held in Cochin, India* from 14-18 November **1988**, Kerala Forest Research Institute, India and International Development Research Center, Canada.
33. Abd. Latif, M. Perusahaan Membuat Bidai dan Pencucuk Sate Kepong, Malaysia. **1987**.
34. Liese, W. Anatomy and properties of bamboo. In: *International Bamboo Workshop*, Hangzhou, **1985**, 6-14 October. Recent research on bamboos: Proceedings edited by A.N. Rao; G. Dhanrajan and C.B. Sastry. Canada, D.R.C. pp. 196-208.
35. Young, R. A.; Haun, J. R. Bamboo in the United States: description, culture, and Utilization. In *Agricultural Handbook*. **1961**, No. 193. Washington, D.C.: USDA Crops Research Division, Agricultural Research Service
36. Zhou, F. C.; Studies on physical and mechanical properties of bamboo woods. Journal of Nanjing Technology College of Forest Products, **1981**, 2, 1-32.
37. Liese, W. Characterization and Utilization of bamboo. Pro. IUFRO 18th World congress on Bamboo production and Utilization, 7-21 September 1986, Ljubljana, Yugoslavia, 11-16.