

Abstract

Morphological Value of Nephrotoxic Effects of Doxorubicin and PLGA-Doxorubicin †

Ivan Tsvetkov ¹, Natalia Zolotova ¹, Anna Kosyreva ¹, Dzhuliia Dzhililova ¹, Liliya Mikhailova ¹, Vera Kudelkina ¹, Artem Shelkov ¹, Olga Makarova ¹, Valeriy Chernikov ¹, Victoria Razzhivina ^{2,*}

¹ Federal State Budgetary Institution «Research Institute of Human Morphology», Moscow, Russia

² LLC Technology of medicines, Khimki, Russia

† Presented at the 1st International Electronic Conference on Molecular Sciences: Druggable Targets of Emerging Infectious Diseases, online, 01-14 September 2021.

Academic Editor: Clemente Capasso

Published: 31 August 2021

Abstract: Doxorubicin (DOX) is a chemotherapy drug that causes nephrotoxicity in rodent models and, to a lesser extent, in cancer patients. doxorubicin hydrochloride or doxorubicin - loaded poly(lactide-co-glycolide acid) (Dox-PLGA) nanoparticles at a therapeutic dose were injected intravenously male Wistar rats. PLGA is a biodegradable polymer used as a drug delivery vehicle. However, the therapeutic effect of Dox-PLGA is not clearly understood. The aim of the study. To estimate a comparative assessment of the nephrotoxic effects of different forms of doxorubicin. A study was carried out on male Wistar rats weighing about 200-250 g (n=24). Morphological assessment of kidneys was performed using light microscopy. Ultrastructural changes were studied using TEM Libra120 transmission electron microscope. The levels of ALT, AST, urea, creatinine in blood serum were determined. Comparison of digital data between experimental groups was performed using the Kruskal-Wallis test (ANOVA). Differences were considered statistically significant at $p < 0.05$. Degeneration changes in the proximal tubules with the destruction of the brush border were revealed by light and electron microscopy in the kidneys on the 8th and 21st days of the experiment after the administration of doxorubicin hydrochloride and its nanosomal form Dox-PLGA. It was found that on the 8th day, PLGA-doxorubicin causes less pronounced degenerative changes in the epithelium of the proximal tubules. The short-term effect after administration of Dox-PLGA is characterized by an increase in creatinine levels, and the long-term effect by an increase in the level of ALT activity, as well as the concentration of urea and creatinine.

Keywords: nephrotoxicity; doxorubicin; PLGA; nanoparticles; toxic effect; morphology; biochemistry