

Proceeding Paper

Evaluation of Technologies for the Co-Extraction of Phenolic Compounds and Proteinaceous Material from Olive-Derived Biomasses †

María del Mar Contreras ^{1,2,*}, Irene Gómez-Cruz ^{1,2,*}, Inmaculada Romero ^{1,2} and Eulogio Castro ^{1,2}

¹ Department of Chemical, Environmental and Materials Engineering, Universidad de Jaén, 23071 Jaén, Spain; iromero@ujaen.es (I.R.); ecastro@ujaen.es (E.C.)

² Centre for Advanced Studies in Earth Sciences, Energy and Environment (CEACTEMA), Universidad de Jaén, Campus Las Lagunillas, 23071 Jaén, Spain

* Correspondence: mcgamez@ujaen.es (M.d.M.C.); igcruz@ujaen.es (I.G.-C.)

† Presented at the 2nd International Electronic Conference on Foods, 15–30 October 2021; Available online: <https://foods2021.sciforum.net/>.

Abstract: The current interest in using olive biophenols to promote functional ingredients and anti-oxidant additives is increasing. These compounds can be obtained from olive fruit and olive-derived biomasses using different technologies. However, other components can be co-extracted. Therefore, the main objective of this study was to evaluate the effect on protein solubilization of several extraction technologies, which were applied to obtain olive biophenols from olive-derived biomasses. For this purpose, conventional (Soxhlet and water bath) and non-conventional technologies (ultrasound and microwave) have been evaluated. The total phenolic content was measured using the Folin & Ciocalteu method and the protein content using the Dumas combustion method. The phenolic profile and the hydroxytyrosol content were also determined. Overall, the highest total phenolic content was obtained using the Soxhlet method, while the microwave-assisted extraction at 100 °C led to the highest protein solubilization (closer to 60%) using water.

Keywords: green extraction; microwave-assisted extraction; olive-derived biomass; protein solubilization

Citation: Contreras, M.d.M.; Gómez-Cruz, I.; Romero, I.; Castro, E.

Evaluation of Technologies for the Co-Extraction of Phenolic Compounds and Proteinaceous Material from Olive-Derived Biomasses. *Biol. Life Sci. Forum* **2021**, *1*, x. **2021**, *1*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s):

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The current interest in using phenolic compounds to promote functional ingredients and antioxidant additives is increasing. This is also the case of olive biophenols obtained from olive fruits and by-products due to their high antioxidant activity and generally no adverse health effects have been shown. Besides the antioxidant activity in food systems, their biological properties make them to be potentially used as multipurpose additives [1].

To recover olive biophenols, conventional technologies such as maceration and Soxhlet extraction have been applied, but also new green trends include the use of ultrasound-assisted extraction (UAE), microwave-assisted extraction (MAE), supercritical fluid extraction, and pressurized liquid extraction [2–6]. The latter technologies can shorten the extraction time, reduce solvent consumption and present a lower energy cost; thus, being more respectful with the environment.

In order to develop a bio-based economy for an efficient conversion of these biomasses, proteinaceous material can be another bioproduct obtained from olive byproducts [2] with potential to be applied in different sectors [7,8]. Some of the latter extraction technologies have also been applied to recover intact and partially hydrolyzed proteins from agri-food bioresources, generally, using water, alkaline solutions and buffers, obtaining

different recoveries [7,9]. Therefore, the main objective of this study was to evaluate the effect of some extraction technologies to co-extract olive biophenols and protein.

2. Material and methods

2.1. Samples and Reagents

Samples were obtained from different industries placed in Jaén, Spain ('Spuny SA', 'SCA Unión Oleícola Cambil' and 'Peláez Renovables'). These biomasses were milled with an Ultra Centrifugal Mill ZM 200 (Retsch, Haan, Germany).

The following reagents were purchased from Sigma-Aldrich (St. Louis, MO, USA): Folin and Ciocalteu's phenol reagent, sodium carbonate, sodium hydroxide and gallic acid. Ethanol was procured from AppliChem (Barcelona, Spain). Hydroxytyrosol was obtained from Extrasynthese (Genay, France).

2.2. Extraction Technologies

Soxhlet extraction with a solid:solvent ratio of around 2.9:100 (*w/v*) was performed using sequentially water and ethanol for 24 h each step. The rest of methodologies were previously optimized in the laboratory to maximize the extraction of phenolic compounds from olive-derived biomasses. Aqueous water extraction using a water bath with agitation was performed at 85 °C for 90 min at 10% solid loading (*w/v*) [10]. UAE (probe-type) was performed using Branson Ultrasonics Corporation device (Danbury, CT, USA). The amplitude was 80%, the extraction time was 16 min and the solid loading was 12% (*w/v*). Finally, MAE was performed at 100 °C for 16 min and the solid loading was 12% (*w/v*) in an Anton Paar microwave (Monowave 400, Graz, Austria). For both, UAE and MAE, water was used as solvent.

After extraction, the samples were vacuum-filtered to separate the extract from the extracted solid fraction. The latter was dried and weighted to determine the solid recovery and the protein content.

2.3. Determination of the Total Phenolic Content and Phenolic Profile

The TPC was determined in the filtered extracts using the Folin & Ciocalteu method according to [10] using gallic acid as standard and a Bio-Rad iMark™ microplate reader (Hercules, CA, USA) was applied.

The phenolic profile was determined by reversed phase (RP)-high-performance liquid chromatography (HPLC) with a diode array detector (Shimadzu Prominence UFLC system) (Kyoto, Japan) according to Contreras et al. [3]. The hydroxytyrosol content was determined at 280 nm using the external standard method.

2.4. Determination of the Protein Content

The protein content was determined through the nitrogen content, which was measured using an elemental analyzer TruSpec Micro (Leco, St. Joseph, MI, USA), and applying a conversion factor of 6.25.

3. Results and Discussion

3.1. Protein Content of Olive Biomasses

Figure 1 shows the protein content of raw olive-derived biomasses. The protein content in olive mill leaves and in the exhausted olive pomace (EOP) was higher than that of olive pulp and intermediate compared to that of other agri-food residues [7,9]. Thus, the former residues are interesting bioresources to obtain proteins; especially, EOP.




Figure 1. Crude protein content in olive-derived biomasses, adapted from Contreras et al. [3].

3.2. Total Phenolic Content of the Extracts

Exhausted olive pomace (EOP) presented the highest TPC value using conventional technologies; particularly, using Soxhlet extraction when the aqueous extract was determined (4.5 g/100 g biomass). However, UAE and MAE reached up to 85% recovery values compared to Soxhlet extraction in a short time.

The phenolic profiles were qualitatively similar and the chromatographic peak corresponding to hydroxytyrosol was the major one. Its content was of 0.6 g/100 biomass in all the extracts. As an example, Figure 2 shows the phenolic profile of the extract obtained by MAE.




Figure 2. Chromatogram (280 nm) of the aqueous extract of exhausted olive pomace obtained using microwave.

3.3. Protein Solubilization

EOP was selected for further study. Figure 3 depicts the protein solubilization, taking into account the initial protein in the raw biomasses and the solid recovery after extraction. All the extraction methodologies seem to provoke the solubilization of a part of the protein content of EOP in the aqueous extracts. The highest protein solubilization was reached using MAE.




Figure 3. Protein solubilization from exhausted olive pomace after applying different extraction technologies.

4. Conclusions

The present results showed that the technologies applied to extract phenolic compounds can provoke the co-extraction of proteins, as is the case of EOP. MAE is a green method, considering that water was used as extractive agent and a shorter time was applied, which can be applied to co-extract both phenolic compounds, including hydroxytyrosol, and protein from EOP.

Author Contributions: Conceptualization, M.d.M.C.; methodology, I.G.-C. and M.d.M.C.; software, I.G.-C. and M.d.M.C.; validation, I.R.; formal analysis, I.G.-C.; investigation, E.C., I.G.-C., I.R., M.d.M.C.; data curation, M.d.M.C.; writing—original draft preparation, M.d.M.C.; writing—review and editing, E.C., I.G.-C., I.R., M.d.M.C.; visualization, M.d.M.C.; supervision, E.C., I.R., M.d.M.C.; project administration, I.R., M.d.M.C.; funding acquisition, I.R., M.d.M.C. All authors have read and agreed to the published version of the manuscript.

Funding: Financial support from Agencia Estatal de Investigación and Fondo Europeo de Desarrollo Regional (FEDER). Reference project ENE2017-85819-C2-1-R. Irene Gómez-Cruz expresses her gratitude to the Universidad de Jaén for financial support (grant R5/04/2017). Authors also thank the FEDER UJA project 1260905 funded by “Programa Operativo FEDER 2014-2020” and “Consejería de Economía y Conocimiento de la Junta de Andalucía”.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data are available on request from the corresponding author.

Acknowledgments: The technical and human support provided by CICT of the Universidad de Jaén is gratefully acknowledged.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Ciriminna, R.; Meneguzzo, F.; Delisi, R.; Pagliaro, M. Olive biophenols as new antioxidant additives in food and beverage. *ChemistrySelect* **2017**, *2*, 1360–1365, doi:10.1002/slct.201601900.
2. Contreras, M.d.M.; Romero, I.; Moya, M.; Castro, E. Olive-derived biomass as a renewable source of value-added products. *Process Biochem.* **2020**, *97*, 43–56, doi:10.1016/j.procbio.2020.06.013.
3. Contreras, M.d.M.; Gómez-Cruz, I.; Romero, I.; Castro, E. Olive pomace-derived biomasses fractionation through a two-step extraction based on the use of ultrasounds: Chemical Characteristics. *Foods* **2021**, *10*, 111, doi:10.3390/foods10010111.
4. Pavez, I.; Lozano-Sánchez, J.; Borrás-Linares, I.; Nuñez, H.; Robert, P.; Segura-Carretero, A. Obtaining an extract rich in phenolic compounds from olive pomace by pressurized liquid extraction. *Molecules* **2019**, *24*, 3108.
5. Medfai, W.; Contreras, M.d.M.; Lama-Muñoz, A.; Mhamdi, R.; Oueslati, I.; Castro, E. How cultivar and extraction conditions affect antioxidants type and extractability for olive leaves valorization. *ACS Sustain. Chem. Eng.* **2020**, *8*, 5107–5118, doi:10.1021/acssuschemeng.9b07175.

6. Taamalli, A.; Arráez-Román, D.; Barraón-Catalán, E.; Ruiz-Torres, V.; Pérez-Sánchez, A.; Herrero, M.; Ibañez, E.; Micol, V.; Zarrouk, M.; Segura-Carretero, A.; Fernández-Gutiérrez, Alberto. Use of advanced techniques for the extraction of phenolic compounds from Tunisian olive leaves: Phenolic composition and cytotoxicity against human breast cancer cells. *Food Chem. Toxicol.* **2012**, *50*, 1817–1825, doi:10.1016/j.fct.2012.02.090.
7. Contreras, M.d.M.; Lama-Muñoz, A.; Manuel Gutiérrez-Pérez, J.; Espínola, F.; Moya, M.; Castro, E. Protein extraction from agri-food residues for integration in biorefinery: Potential techniques and current status. *Bioresour. Technol.* **2019**, *280*, 459–477, doi:10.1016/j.biortech.2019.02.040.
8. Sari, Y.W.; Mulder, W.J.; Sanders, J.P.M.; Bruins, M.E. Towards plant protein refinery: Review on protein extraction using alkali and potential enzymatic assistance. *Biotechnol. J.* **2015**, *10*, 1138–1157, doi:10.1002/biot.201400569.
9. Pojić, M.; Mišan, A.; Tiwari, B. Eco-innovative technologies for extraction of proteins for human consumption from renewable protein sources of plant origin. *Trends Food Sci. Technol.* **2018**, *75*, 93–104, doi:10.1016/j.tifs.2018.03.010.
10. Gómez-Cruz, I.; Cara, C.; Romero, I.; Castro, E.; Gullón, B. Valorisation of exhausted olive pomace by an eco-friendly solvent extraction process of natural antioxidants. *Antioxidants* **2020**, *9*, 1010, doi:10.3390/antiox9101010.