

# Preliminary Characterization of a New Processive Endoglucanase from *Clostridium alkallicellulosi* DSM17461

 Techly San <sup>1</sup> and Paripok Phitsuwan <sup>1,2,\*</sup>
<sup>1</sup> Division of Biochemical Technology, School of Bioresources and Technology, King Mongkut's University of Technology Thonburi, Bangkok 10150, Thailand; techly.san@mail.kmutt.ac.th

<sup>2</sup> LigniTech-Lignin Technology Research Group, School of Bioresources and Technology, King Mongkut's University of Technology Thonburi, Bangkok 10150, Thailand

\* Correspondence: paripok.phii@kmutt.ac.th; Tel.: +662 470 7751

† Presented at the 2<sup>nd</sup> International Electronics Conference On Catalysis Sciences—A Celebration of *Catalysts* 10<sup>th</sup> Anniversary, 15-30 October 2021; Available online: <https://eccs2021.sciforum.net>

**Abstract:** The *Clostridium alkallicellulosi* DSM17461<sup>T</sup> genome contains several glucoside hydrolase encoding genes essential for cellulose degradation. Herein, the family 9 glycoside hydrolase enzyme (*CalGH9\_2089*) was cloned and expressed. The enzyme contains one GH9 catalytic module, a family 3 carbohydrate-binding module, and one Type I dockerin at its C-termini. The optimal pH and temperature for *CalGH9\_2089* to hydrolyze CMC were 55 °C and pH 6.0, with the remaining activity of more than 60% at pH 10.0. *CalGH9\_2089* produced a series of cello-oligomers (G2-G6) from CMC, suggesting that the enzyme has an endo-acting capability. When regenerated amorphous cellulose was hydrolyzed with *CalGH9\_2089*, the ratio of reducing ends in the soluble fraction to that in the insoluble pellets was 4.8, suggesting this enzyme acts processively on RAC. This work extends our knowledge of the behavior of the GH9 endoglucanase from the microorganism living in an alkaline environment.

**Keywords:** family 9 glycoside hydrolase; endoglucanase; processivity; carbohydrate-binding module; dockerin

## 1. Introduction

Cellulose is an unbranched glucose polymer that is linked by the β-1,4-glycosidic bond. Individual cellulose chains are closely associated through intra- and inter-hydrogen bonds, forming a tightly packed structure with a few amorphous regions (semi-crystalline). The crystalline structure makes cellulose insoluble, thus posing difficulty in hydrolysis [1,2]. To break down cellulose, three main enzymes, i.e., endo-1,4-β-glucanases, exo-glucanases/cellobiohydrolases, and β-glucosidases, are required to work in concert to degrade crystalline cellulose to simple sugar successfully. The endo-1,4-β-glucanases are important enzymes. They attack the β-1,4-glycosidic bonds at random sites along the cellulose chain, freeing sections of the chain from the crystal surface and creating free ends, upon which exo-glucanases can act, thereby, generating cellobioses which are a substrate for β-glucosidases to produce glucose molecules [1].

According to the carbohydrate-active enzymes (CAZy) database, glycosidases are classified on the basis of similarities in the protein sequence and three-dimensional structure. Cellulases are belonged to glycoside hydrolase families (GH) 5-10, 12, 26, 44, 45, 48, 51, 61, and 74, etc. [3]. Most endo-1,4-β-glucanases belong to GH9. GH9 endoglucanases catalyze the hydrolysis of β-1,4 linkages with the inversion of anomeric carbon configuration. Several GH9 endoglucanases are tightly associated with a carbohydrate-binding module (CBM) classified in family 3 [4].

**Citation:** Lastname, F.; Lastname, F.; Lastname, F. Title. *Chem. Proc.* **2021**, *3*, x. <https://doi.org/10.3390/xxxxx>

Published: date

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

In bacterial cellulase systems, GH9 endoglucanases are distinctive from other cellulase families. GH9 endoglucanases have shown either non-processive or processive activity. The processivity is the ability to remain to adsorb the substrate in between subsequent hydrolytic reactions, and this ability is important for the degradation of crystalline cellulose [5,6]. Therefore, the processive glucanases, behaving like exoglucanases, are thought to be a major cellulose-degrading factor besides the exoglucanase component [5,6]. A recent study showed that a single GH9 endoglucanase is essential for cellulose degradation by *Clostridium phytofermentans*, despite its production of several other cellulases [7].

Carbohydrate-binding modules (CBMs) are non-catalytic modules found in some carbohydrate-active enzymes. They play an important role in degrading insoluble substrates [8,9]. Based on the structure of the binding site and functional similarity, CBMs are grouped into three types: Type A CBMs that bind the planar surfaces of crystalline polysaccharides. Type B CBMs bind internally on the glycan chains (endo-type). Type C CBMs bind the termini of glycan (exo-type). CBMs have been classified into 81 families based on their amino acid similarities (<http://www.cazy.org>) [10], and a family 3 CBM has shown to bind cellulose effectively. There are numerous examples of fusing CBMs to catalytic modules, subsequently resulting in 2- to 10-fold increase in enzyme activities on insoluble substrates but not on soluble polysaccharides [11,12].

*Clostridium* spp. have received considerable attention as a source of cellulolytic and enzymes [13]. A number of endo-1,4- $\beta$ -glucanases from several *Clostridial* species have been studied in recent years, some of which show promising applications in biofuel and biochemical production due to their high activity and high thermostability [13]. The *Clostridium alkalicellulosi* DSM17461<sup>T</sup> genome contains several glycoside hydrolase encoding genes important for cellulose degradation. Herein, the family 9 glycoside hydrolase enzyme (*CalGH9\_2089*) was cloned and expressed. The recombinant enzyme's biochemical properties (i.e., pH and temperature optima, mode of action, and processivity) were determined.

## 2. Materials and methods

### 2.1. Cloning and protein expression

Genomic DNA of *C. alkalicellulosi* DSM17461<sup>T</sup> was purchased from Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures (Germany).

PCR amplification of the putative endoglucanase gene (*CalGH9\_2089*) was performed using the upper primer (5'-AATTAGCTAGCTCAACTTTTAACTACGGAGA-3'), and the lower primer (5'-AATTACTCGAGTTACTGGTTTACTGGGAAAA-3') using the genomic DNA as the template and the Phusion Taq High-Fidelity PCR-System (ThermoFisher Scientific, USA). The PCR amplified material was cloned in pET28a(+) (Novagen, USA) by digestion with restriction enzymes *NheI* and *XhoI*.

The purified PCR products and pET28a(+) vectors were doubly digested with *NheI* and *XhoI*. The digested PCR products and pET28a(+) vectors were ligated with T4 DNA ligase (New England Biolabs, UK). The ligated plasmids were transformed into NEB 5-alpha *E. coli* cells.

The positive clones with the target insert were transformed into *E. coli* BL21 (DE3) for protein expression following the manufacturer protocols (New England BioLabs). The recombinant *CalGH9\_2089* protein was purified from a 400 mL transformed *E. coli* BL21 (DE3) culture.

### 2.2. Enzyme Activity Assay

The enzyme assay was performed in a total reaction volume of 200  $\mu$ L. The reaction mixture consisted of 1% (w/v) substrate (i.e., CMC and RAC) and 0.25  $\mu$ M of *CalGH9\_2089* in 50 mM citrate buffer pH 6.0. The reaction mixture was incubated at 55  $^{\circ}$ C for 15 min (and 3 h for RAC). The reaction was terminated by immersion in ice water and centrifuged at 13,000 $\times$  g at 4  $^{\circ}$ C for 10 min to separate the substrate from the soluble fraction. One

hundred microliters of the soluble fraction was taken to a new 1.5-mL microcentrifuge tube and 150  $\mu$ L of dinitrosalicylic acid (DNS) [14] was added. The reaction mixture was boiled for 10 min and centrifuged at high speed. A sample was taken and measured at an absorbance of 540 nm. The amount of reducing sugar released in the mixture was estimated using a glucose standard curve. One unit of the enzyme was defined as the amount of enzyme that produces 1  $\mu$ Mol of glucose equivalent in 1 min under the assayed conditions.

### 2.3. Effect of pH and Temperature

The influence of pH on enzyme activity was determined. The activity at different pH values from pH 3.0–11.0 was measured at 55  $^{\circ}$ C for 15 min. Citrate-sodium citrate buffer for pH 3.0–7.0, Tris-HCl buffer for pH 7.0–9.0, and glycine-NaOH for pH 9.0–11.0. with a concentration of 50 mM were used as a reaction buffer. The maximum activity at the corresponding pH was taken as 100%. The pH stability assay was performed by incubating 5.5  $\mu$ L of the 11.5  $\mu$ M enzyme in 10 mM of the buffers mentioned above at 30  $^{\circ}$ C for 60 min. After that, the enzyme solution was diluted, and the residual activity was measured under optimal assay conditions.

The effect of temperature on enzyme activity was tested by incubating 0.25  $\mu$ M enzyme in a 200- $\mu$ L reaction mixture containing 1% (w/v) substrate at a temperature range of 40–70  $^{\circ}$ C, pH 6.0 (citrate buffer) for 10 min. The maximum activity at the corresponding temperature was taken as 100%. The thermal stability of the enzyme was determined by incubating 5.5  $\mu$ L of the 11.5  $\mu$ M enzyme at temperatures of 37, 55, 60, and 70  $^{\circ}$ C for 0–24 h. The enzyme solution was diluted, and the residual activity was determined under optimal assay conditions.

### 2.4. Analysis of the Hydrolysis

The reaction mixture (total volume of 200  $\mu$ L) containing 0.25  $\mu$ M enzyme and 1% (w/v) substrate (i.e., CMC) was assayed under optimal conditions with varying times from 15 min to 16 h. The reaction was stopped at specific time intervals by boiling for 5 min and centrifuging at 13,000 $\times$  g at 4  $^{\circ}$ C for 10 min to remove the undigested substrate. The soluble fraction was transferred to a new tube, and 4  $\mu$ L of the soluble fraction was mixed with 4  $\mu$ L of absolute ethanol. The mixed sample (8  $\mu$ L) was spotted on TLC plates, and the plates were immersed in a TLC chamber containing n-butanol:acetic acid:water at a ratio of 2:1:1 as the mobile phase, and heated at 90  $^{\circ}$ C in a hot air oven for visualization [15].

### 2.5. Processivity Assay

The content of reducing ends in the soluble and insoluble fractions was determined using RAC as a substrate [16]. Different concentrations of enzyme were added to a 1.5-mL microcentrifuge tube containing 1% (w/v) RAC in citrate buffer (pH 6.0) (final volume of 200  $\mu$ L). The reaction mixture was incubated at 55  $^{\circ}$ C for different incubation times. After incubation, the reaction tube was centrifuged at 13,000 $\times$  g at 4  $^{\circ}$ C for 10 min, and the supernatant was removed, of which 100  $\mu$ L was taken for measurement of reducing ends (soluble fraction) by the DNS method. The pellet was washed three times with 500  $\mu$ L citrate buffer (pH 6.0) by centrifugation at 13,000 $\times$  g at 4  $^{\circ}$ C for 5 min. After washing, the pellet was resuspended in 100  $\mu$ L citrate buffer (pH 6.0) and taken for measurement of reducing ends (insoluble fraction) by the DNS method.

## 3. Results and Discussion

### 3.1. Bioinformatic Analysis

The draft genome sequence of *C. alkalicellulosi* DSM17461 was sequenced, revealing several putative endoglucanase encoding genes essential for cellulose hydrolysis in its genome. One ORF having 2899 bp was designated as *CalGH9\_2089*. This gene was predicted to encode a putative glycoside hydrolase family 9 endoglucanase connected

with a family 3 CBM and a type I dockerin according to BlastP analysis (https://blast.ncbi.nlm.nih.gov/Blast.cgi, accessed on 6 July 2021). This gene was predicted to have 748 amino acid residues with a theoretical molecular weight of 82.86 kDa and pI of 5.1 (https://web.expasy.org/cgi-bin/protparam/protparam/, accessed on 6 July 2021) (Figure 1). The *CalGH9\_2089* was successfully expressed in *E. coli* BL21 (DE3), and it was determined for its molecular weight and purity, showing a single band with a molecular weight of ca. 83 kDa.

```

10      20      30      40      50      60
MKKIISVVIV VAVLVGLVTV TPSPAEASTF NYGEALQKSI MFYEFQMSGK LPDNLRTNWR

70      80      90      100     110     120
GDSCLDGSD VGLDLTGGWF DAGDHIKFNL PMSYTSTMLA WAVYEYKDAL ERSGQLPYIK

130     140     150     160     170     180
QQIRWATEYF IKCHPEKYVY YYQVGDGITD HRWWVPAEVI HLQSVRSHK VTLDSPGSAV

190     200     210     220     230     240
VAGTAAALAS AAIVFQSDP AYAALCLKHA KDLPDFADRT QSDAGYTAAN NYYDSWSGFW

250     260     270     280     290     300
DELSWAGVWI YMASGEKAFI DKAESYVANW NREERTNLLA YKWGHCWDDV MYGASLLLAK

310     320     330     340     350     360
ATNKSIYKEH VERHLDYWSV GYNGERITYT PKGLAHLFVW GVLRHATTTA FLASVYSDWS

370     380     390     400     410     420
ECPPAKAKTY MDTAKQVDY ALGSSGRSYV VGFGVNPQPH PHHRTAHSSW IDTMEEPSYH

430     440     450     460     470     480
RHVLYGALVG GPNQSDAYVD DIGDYITNEV ACDYNAGFVG ALAKMYDVYG GDPFIPGFNAI

490     500     510     520     530     540
EEVPPYPIYV TASLSSRTTA TEVKAFLINK SGWPARVKDT LSFKYFVDLT DFINAGHSPN

550     560     570     580     590     600
EITSSIIYSA APTAKITGPI AYDTSKNIYV FELDLKGTAI FPGSRMDHQK EVQFHIVPPN

610     620     630     640     650     660
GAPWNIPTDP SYPGTLSEDE PVPQIPVYDN GVLLFGLEPD GSTPQPTTPP TTPPTTPPP

670     680     690     700     710     720
TTTTPTPPPA IMAGDINGDG LINSTDYILL RRYILEVTPS LPTTDVSGNP YRGDLAADLN

730     740
GDGLIDSIDV ILMRRYILEI ITVFPVNO

```

Figure 1. A translated amino acid sequence of *CalGH9\_2089*.

### 3.2. Effects of pH and Temperature

The effect of pH on *CalGH9\_2089* activity was tested using CMC as a substrate. *CalGH9\_2089* exhibited an activity from pH 4.0 to 11.0, with maximal activity at pH 6.0-7.0. The enzyme showed activity of more than 50% from pH 5.0-10.0 (Figure 2A). For pH stability, the enzyme was pre-incubated in different buffers at 55 °C for 60 min and was assayed at the optimal conditions. It was found that the enzyme showed good stability from pH 5.0-10.0 with residual activity of more than 60% (Figure 2B). The pH profile suggested that *CalGH9\_2089* is active in a broad pH range and tolerates alkaline pHs.





Figure 2. Effect of pH on *CalGH9\_2089* activity towards CMC (A) and its pH stability (B).


The effect of temperature on the activity of *CalGH9\_2089* was examined with the CMC under the optimal conditions. *CalGH9\_2089* exhibited the highest activity of 1804.98  $\mu\text{mol product}/\text{min}/\mu\text{mol protein}$  on CMC. The optimal temperature of *CalGH9\_2089* toward CMC was 55-60 °C (Figure 3A). At 80 °C, the enzyme was almost completely inactivated. The thermal stability analysis of *CalGH9\_2089* was investigated at pH 6.0 with different temperatures (37, 55, 60, and 70 °C) (Figure 3B). The result showed that *CalGH9\_2089* was stable at 37 °C, retaining more than 80% activity for 24 h. At 55 °C, the residual activity remained about 60% when the enzyme was incubated for 30 min. The enzyme activity remained 40% of its maximal activity from 6-24 h-incubation. At 60 °C and 70 °C, the activity of *CalGH9\_2089* lost rapidly, retaining 20% and 10% of its maximal activity, respectively, when it was incubated for 30 min. This result likely indicates that *CalGH9\_2089* is a mesophilic enzyme and tolerates a moderate temperature around 50 °C.



**Figure 3.** (A) Effect of temperature on *CalGH9\_2089* activity towards CMC and (B) its thermal stability.

### 3.3. Mode of action

To investigate the mode of action of *CalGH9\_2089*, hydrolysis of CMC was tested at pH 6.0 and 55 °C, and the hydrolysis products at different incubation times were determined by TLC (Figure 4). It was found that *CalGH9\_2089* was able to hydrolyze CMC. At 1 min, small amounts of oligosaccharides were present around cellotriose (G3), cellotetraose (G4), compared to the standard. From 3 to 10 min, more oligosaccharides, particularly cellobiose (G2), G3, and G4, were accumulated, and they were clearly observed on the TLC plate. After 15 min to 16 h, the amounts of G2, G3, and G4 greatly increased, together with the accumulation of glucose (G1). This hydrolysis pattern likely indicates the endo-acting mode of *CalGH9\_2089*.



**Figure 4.** TLC analysis of hydrolysis product from CMC by *CalGH9\_2089*.

### 3.4. Processivity

Some GH9 endoglucanases have been shown to have processivity like exo-glucanases [17]. The processivity of endoglucanase is usually determined by the ratio of the generated soluble reducing ends to insoluble reducing ends [16]. Here, the distribution of reducing sugars generated by *CalGH9\_2089* on RAC was studied. It was found that the ratio of reducing ends in the soluble fraction to that in the insoluble fraction was increased from 1.6 to 4.8 as the incubation time was prolonged from 30 to 180 min (Figure 5). This result indicates that *CalGH9\_2089* is a processive endoglucanase, differing from the common endoglucanases, whose actions randomly cut and produce more reducing ends in the insoluble fraction than in the soluble fraction.




Figure 5. The processivity of *CalGH9\_2089* towards RAC.

## 4. Conclusion

Herein, we successfully expressed a putative GH9 endoglucanase having a modular structure containing a GH9 catalytic module, a family 3 CBM, and one Type I dockerin (*CalGH9\_2089*) from *C. alkalicellulosi* genome. The enzyme was functionally active from pH 5.0–10.0 with an optimal pH around 6.0–7.0. The optimal temperature for this enzyme is 55 °C; however, its half-life was around 1 h. The production of a series of oligosaccharides from CMC indicates an endo-acting mode of *CalGH9\_2089*. Moreover, the ratio of reducing ends in the soluble fraction to that in the insoluble fraction using RAC as a substrate was 4.8 (3 h-incubation), suggesting this *CalGH9\_2089* endoglucanase possesses processive activity.

**Author Contributions:** Conceptualization, P.P.; methodology, T.S. and P.P.; validation, T.S. and P.P.; formal analysis, P.P.; investigation, T.S.; data curation, T.S. and P.P.; writing—original draft preparation, T.S.; writing—review and editing, P.P.; supervision, P.P.; funding acquisition, P.P. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by Thailand Science Research and Innovation (TSRI), Basic Research Fund: Fiscal year 2022.

**Data Availability Statement:** All data are provided by the manuscript and the Supporting Information. Relevant data are available from the authors on request.

**Acknowledgments:** We gratefully thank Edward A. Bayer (Weizmann Institute of Science, Israel) for providing genomic information on *C. alkalicellulosi*.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Phitsuwan, P.; Sakka, K.; Ratanakhanokchai, K. Improvement of lignocellulosic biomass in planta: a review of feedstocks, biomass recalcitrance, and strategic manipulation of ideal plants designed for ethanol production and processability. *Biomass Bioenergy* **2013**, *58*, 390-405. 1-4
2. Himmel, M.E.; Ding, S.Y.; Johnson, D.K.; Adney, W.S.; Nimlos, M.R.; Brady, J.W.; Foust, T.D. Biomass recalcitrance: engineering plants and enzymes for biofuels production. *Science* **2007**, *315*, 804-807. 5-6
3. Henrissat, B. A classification of glycosyl hydrolases based on amino acid sequence similarities. *Biochem. J.* **1991**, *280*, 309-316. 7
4. Kurokawa, J.; Hemjinda, E.; Arai, T.; Kimura, T.; Sakka, K.; Ohmiya, K. *Clostridium thermocellum* cellulase CelT, a family 9 endoglucanase without an Ig-like domain or family 3c carbohydrate-binding module. *Appl. Microbiol. Biotechnol.* **2002**, *59*, 455-461. 8-10
5. Zverlov, V.V.; Schantz, N.; Schwarz, W.H. A major new component in the cellulosome of *Clostridium thermocellum* is a processive endo- $\beta$ -1, 4-glucanase producing cellotetraose. *FEMS Microbiol. Lett.* **2005**, *249*, 353-358. 11-12
6. Arai, T.; Kosugi, A.; Chan, H.; Koukiekolo, R.; Yukawa, H.; Inui, M.; Doi, R.H. Properties of cellulosomal family 9 cellulases from *Clostridium cellulovorans*. *Appl. Microbiol. Biotechnol.* **2006**, *71*, 654-660. 13-14
7. Tolonen, A.C.; Chilaka, A.C.; Church, G.M. Targeted gene inactivation in *Clostridium phytofermentans* shows that cellulose degradation requires the family 9 hydrolase Cphy3367. *Mol. Microbiol.* **2009**, *74*, 1300-1313. 15-16
8. Boraston, A.B.; Bolam, D.N.; Gilbert, H.J.; Davies, G.J. Carbohydrate-binding modules: fine-tuning polysaccharide recognition. *Biochem. J.* **2004**, *382*, 769-781. 17-18
9. Gilbert, H.J.; Knox, J.P.; Boraston, A.B. Advances in understanding the molecular basis of plant cell wall polysaccharide recognition by carbohydrate-binding modules. *Curr. Opin. Struct. Biol.* **2013**, *23*, 669-677. 19-20
10. Lombard, V.; Golaconda Ramulu, H.; Drula, E.; Coutinho, P.M.; Henrissat, B. The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res.* **2013**, *42*, D490-D495. 21-22
11. Ye, X.; Zhu, Z.; Zhang, C.; Zhang, Y.-H.P. Fusion of a family 9 cellulose-binding module improves catalytic potential of *Clostridium thermocellum* cellodextrin phosphorylase on insoluble cellulose. *Appl. Microbiol. Biotechnol.* **2011**, *92*, 551-560. 23-24
12. Thongekkaew, J.; Ikeda, H.; Masaki, K.; Iefuji, H. Fusion of cellulose binding domain from *Trichoderma reesei* CBHI to *Cryptococcus* sp. S-2 cellulase enhances its binding affinity and its cellulolytic activity to insoluble cellulosic substrates. *Enzyme Microb. Technol.* **2013**, *52*, 241-246. 25-27
13. Thomas, L.; Joseph, A.; Gottumukkala, L.D. Xylanase and cellulase systems of *Clostridium* sp.: An insight on molecular approaches for strain improvement. *Bioresour. Technol.* **2014**, *158*, 343-350. 28-29
14. Miller, G.L. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal. Chem.* **1959**, *31*, 426-428. 30
15. Phitsuwan, P.; Lee, S.; San, T.; Ratanakhanokchai, K. CalkGH9T: A glycoside hydrolase family 9 enzyme from *Clostridium alkalicellulosi*. *Catalysts* **2021**, *11*. 31-32
16. Zhang, X.Z.; Sathitsuksanoh, N.; Zhang, Y.H.P. Glycoside hydrolase family 9 processive endoglucanase from *Clostridium phytofermentans*: Heterologous expression, characterization, and synergy with family 48 cellobiohydrolase. *Bioresour. Technol.* **2010**, *101*, 5534-5538. 33-35
17. Zhang, K.D.; Li, W.; Wang, Y.F.; Zheng, Y.L.; Tan, F.C.; Ma, X.Q.; Yao, L.S.; Bayer, E.A.; Wang, L.S.; Li, F.L. Processive degradation of crystalline cellulose by a multimodular endoglucanase via a wirewalking mode. *Biomacromolecules* **2018**, *19*, 1686-1696. 36-38