



Effect of the re-growth age on the primary metabolites of *Tithonia diversifolia*, part 1: Experiment design and nitrogen metabolism.

Maidelin Paumier^a, Danis Verdecia^a, Hernán Uvidia^c, Jorge Ramírez^a, Rafael Herrera^b, Jhoel Uvidia, Edgar Chicaiza^c, Angel Santana^a.


^a Universidad de Granma, Apartado Postal 21, Bayamo Granma, Cuba (C.P. 85 100).

^b Instituto de Ciencia Animal (ICA), Apartado Postal 24 San José de las Lajas, Mayabeque; Cuba.

^c Universidad Estatal Amazónica, Puyo, Pastaza, Ecuador.

^e Universidad Politécnica de Valencia. Valencia, España.

Graphical Abstract



Abstract.

The primary metabolites, very abundant in nature, are essential for the physiological development of the plant; they are present in large quantities, they are easy to extract, and their exploitation is relatively cheap. In order to establish mathematical expressions that relate the regrowth age with the nitrogen and sugar content (glucose, fructose and sucrose) of the whole plant and its fractions. The experiment was developed, following a randomized block design, with 5 treatments (60, 90, 120, 150 and 180 d). In the first part of this communication the content of Nitrogen (N) in the integral plant, leaves and stems were evaluated. In the second part we studied sugars like Glu, Frut and Suc. Reporting that N, Glu, Frut and Suc decreased with the highest results at 60 days and quadratic and cubic equations with R² higher than 0.90 were adjusted. The established regression equations explain the close relationship between regrowth age and the contents of precursor metabolites (N, Glu, Frut and Suc), which explains the fluctuations found in sugars influenced by the

Introduction

Tithonia diversifolia is a shrubby herbaceous species with showy yellow flowers. It has great adaptability, the same can be found at sea level or at an altitude of approximately 2 400 m it can inhabit soils of high or low fertility. It has additional benefits for erosion control, it can be used as food for the animal, with high nitrogen content and a source of carbohydrates ^[1].

The role of Mathematics and its applications has been different in the spheres of human activity and in different times. It was formed historically under the considerable influence of two factors: the level of development of the mathematical apparatus and the degree of maturity of the knowledge about the object of study; as well as the possibility of describing its most important features and properties in a language of symbols and equations or, as has often been said, the possibility of building a mathematical model of the object to be studied ^[2].

Mathematical modeling is a very useful tool in different disciplines of knowledge. In Cuba, this technique has been developed in the animal branch ^[3]. However, in studies with plants, specifically in the evaluation of pastures, the works are more incipient and limited ^[1].

For all the above, the objective is to establish mathematical expressions that relate the regrowth age with the nitrogen and sugar content of the whole plant and its fractions.

Materials and Methods

The study was developed in the Teaching-Productive Department of the University of Granma, which is located in the southeast of Cuba, in the province of Granma, 17.5 km from the city of Bayamo. The experiment was carried out for two years, and two periods were considered, the rainy one (May-October) and little rainfall (November-April).

The soil present in the area was Calcium brown ^[4], with a pH of 6.2. The content of P₂O₅, K₂O and total N was 2.4; 33.42 and 3 (mg/100g of soil) respectively, with 3.6% organic matter.

Regarding the behavior of the climatic variables, during the rainy period, the precipitations were 731.4 mm; the average, minimum and maximum temperature and relative humidity registered values of 26.73; 22.31 and 33.92 °C and 80.78; 51.02 and 96.22%, respectively.

In the second period, the rainfall reached values of 270 mm; the temperature was 24.05; 18.29 and 31.58 °C and the relative humidity of 76.21; 44.16 and 97.03%, in both cases for the mean, minimum and maximum averages.

Treatment and experimental design

A randomized block design with four replications was used and the treatments were regrowth ages of 60, 90, 120, 150 and 180 days.

Procedures

The already established species had 98% population and at the beginning of each seasonal period a homogeneity cut was made at 15 cm above ground level. The samplings were carried out in 10 plants in a row, eliminating the edge effect in an area of 0.5 ha-1, according to the treatments. The sample was homogenized and weighed, later they were separated manually into leaves, petioles and stems with a diameter of less than two centimeters. Then two kilograms were taken for each of the treatments to determine the dry matter (DM). During the experimental stage neither irrigation nor fertilization was applied.

Chemical analysis

The samples were dried at room temperature in a dark and ventilated room for 12 days, then they were ground to a particle size of one millimeter and stored in amber bottles at room temperature until their analysis. The following were determined: DM and N according to ^[5], while the glucose, fructose and sucrose contents according to the Lane and Eynon titration method, which is based on the reduction of Cu +2 to Cu +1 by the reducing sugars, using methylene blue as indicator ^[5].

Statistical calculations and analysis

The relationship between age with nitrogen and sugars was established through regression analysis where the following were used: linear, quadratic, cubic, logarithmic and gompertz, where the results with the highest adjustment were used for what they were considered. the criteria of ^[2, 6, 7]. For this, the statistical program SPSS version 22 was used.

Results and Discussion (*optional*)

The contents of nitrogen (N), during the rainy period in *T. diversifolia* (Table 1) decreased with the regrowth age in (13.15; 9.99, 3.16), (0.0091, 0.0051, 0.0048), (0.0213, 0.0117, 0.0095), (0.0117, 0.0029, 0.0047 g / Kg) each one of the indicators for the integral plant (IP), leaves and stems, respectively. Quadratic equations were adjusted for all the indicators.

This species during the dry season (Table 2) maintained a similar behavior with adjustments for the quadratic equations. The highest values were obtained at 60 days for nitrogen, decrease in N (17.39; 12.47; 4.93). The set of chemical reactions that take place in an organism constitutes metabolism.

Most of the carbon, nitrogen and energy ends up in molecules common to all cells, necessary for their functioning and that of organisms. They are amino acids, nucleotides, sugars, and lipids, present in all plants and performing the same functions. They are called primary metabolites ^[8].

The adjusted models and the R² values reported in tables 1 and 2 are like those reported by ^[9, 10, 11] when evaluating the effect of age and climatic factors on the content of nitrogen in forage species in the Cauto Valley. The decrease in N with the cutting frequency may be related to a reduction in the synthesis of protein compounds, a decrease in the number of leaves, an increase in the stem fraction and an increase in the production of structural carbohydrates (cellulose and hemicellulose), although it is important to note that the values in both periods in each species exceeded 22.4 g / Kg.

Table 1. Relationship between regrowth age, nitrogen content of *Tithonia diversifolia* in the rainy season

Indicadors	A	b	SE±	C	SE±	D	SE±	R²	1-R²	MSE	SE±
Integral Plant	50,551	-0,206	0,012	0,000399	0,000052			0,99	0,01	0,12	0,347
Nitrogen leaves	42,447	-0,141	0,011	0,000235	0,000047			0,99	0,01	0,102	0,319
Stems	8,115	-0,066	0,002	0,000166	0,000008			0,99	0,01	0,003	0,054

R² todos a p<0,001

Table 2. Relationship between regrowth age, nitrogen content of *Tithonia diversifolia* in the dry season

Indicadors	A	b	SE±	C	SE±	D	SE±	R²	1-R²	MSE	SE±
Integral Plant	44,016	0,101	0,017	-0,001	0,000072			0,99	0,01	0,235	0,485
Nitrogen leaves	33,373	0,169	0,015	-0,001	0,00006			0,99	0,01	0,162	0,403
Stems	10,643	-0,068	0,003	0,000113	0,000013			0,99	0,01	0,008	0,088

R² todos a p<0,001

Other authors such as ^[12] associated this behavior with the high proportion of stems in the sample, since, usually, in the literature, N values of approximately 33.6 g / Kg are reported in the leaves, while in the stems ranges between 11.2 and 32 g / Kg. Generally, the values observed in the analyzed plants are similar to those of temperate legumes. The contents found are in the range established by^[13], for tropical legumes, which gives importance to the results of this work.

On the other hand ^[14] stated that *T. diversifolia* is a plant characterized by its excellent quality, due to its low fiber content and a good ratio of nutrients in its foliage. When conducting ^[15] a study similar to this research, they found similarities in terms of the N content of *T. diversifolia* with *G. sepium*, which emphasizes the importance of using the combination of these legume forage species in animal feed.

The quality of the trees varies in the different components of the biomass. The leaves present higher concentrations of nutrients than the branches and stems, the variation has also been related to age, with the young leaves being richer in N than the old ones ^[16]. These results coincide with those reported for this species by ^[17, 18, 19, 20].

The high variability in sugars with age is due to the different photosynthesizing capacity of the species, related to the element potassium, which is a mediator of the metabolism and transport of primary carbohydrates in plants ^[21, 22].

On the other hand, the soluble carbohydrate content is linked to the morphostructural development of plants. The concentrated reserves of these compounds, in smaller quantities, in the growth points (buds) favor the foliar concentrations of the saccharides after the regrowth emission. However, although these aspects are generally described, from the physiological point of view, the behavior of energy metabolites as a function of morphostructural variations depends on the species, nutritional status and edaphoclimatic conditions in which it is grown ^[21, 2. 3].

The sugar concentrations in *Tithonia* were similar to those obtained in the foliage of other non-legume plants such as *Morus alba*, *Trichantera gigantea*, *Cnidioscolum aconitifolium* and *Ficus carica* ^[24]. However, the amounts were lower than those reported in some legumes of traditional use as forage ^[25], in which the level of these compounds

in mature foliage's has been determined. This corroborates what has been described previously, where the intervention of different factors in the contents of said sugars is explained.

On the other hand, according to the reports of ^[26] there is a close relationship between the carbohydrate composition, age and N content, on the content of secondary metabolites.

Conclusions

The established regression equations explain the close relationship between regrowth age and the contents of precursor metabolites (N), which explains the fluctuations found influenced by the phenological state of the plants and photosynthetic activity of this.

References (mandatory)

- [1]. Ruiz TE, Torres V, Febles G, Díaz H, González J. Empleo de la modelación para estudiar el crecimiento del material vegetal 23 de *Tithonia diversifolia*. Revista Cubana de Ciencia Agrícola. 2012, 46(1): 23-29.
- [2]. Guerra C, Cabrera A, Fernández L. Criterios para la selección de modelos estadísticos en la investigación científica. Revista Cubana Ciencias Agrícolas. 2003; 37(1): 3-10.
- [3]. Torres V, Ramos N, Lizazo D, Monteagudo F, Noda A. Modelo estadístico para la medición del impacto de la innovación o transferencia tecnológica en la rama agropecuaria. Revista Cubana de Ciencia Agrícola. 2008, 42(2):133-140.
- [4]. Hernández A, Ascanio MO, Morales M, Cabrera A. Correlación de la nueva versión de clasificación genética de los suelos de cuba con las clasificaciones internacionales y nacionales: Una herramienta útil para la investigación, docencia y producción agropecuaria. EDINCA, Instituto Nacional de Ciencias Agrícolas. C. de la Habana, Cuba. 2005. 61 pp.
- [5]. AOAC. Official Methods of Analysis, Ass. Off. Agric. Chem. 2000. 17th Ed. Washington, D.C. USA. 2:777-778.
- [6]. Keviste A, Álvarez JG, Rojo A, Ruíz A D. Funciones de crecimiento de aplicación en el ámbito forestal. Ministerio de Ciencia y Tecnología. Instituto de Investigaciones y Tecnología Agraria y Alimentaria. Madrid. España. 2002. 190 pp.
- [7]. Torres V, Ortiz J. Application of modelling and simulation to the production and feeding of farm animals. Cuban J. Agric. Sci. 2005; 39:397-405.
- [8]. Ávalos A, Pérez-Urria E. Metabolitos secundarios de plantas. Reduca (Biología). Serie Fisiología Vegetal. 2009, 2: 119-145.
- [9]. Ramírez JL. Rendimiento y valor nutritivo de cinco gramíneas en el Valle del Cauto. [Tesis de Doctorado]. Universidad de Granma, Cuba. 2010. 101.
- [10]. Ramírez JL, Herrera RS, Leonard I, Cisneros M, Verdecia D, Álvarez Y, López B. Relationship between quality indicators and age indicators in *Pennisetum purpureum* vc. Cuba CT-169 in Cauto Valley, Cuba. Cuban J. Agric. Sci. 2012, 46: 315-319.

- [11]. Verdecia DM, Herrera RS, Ramírez JL, Leonard I, Álvarez Y, Bazán Y, Arceo Y, Bodas R, Andrés S, Álvarez J, Giráldez FJ, López S. Valor nutritivo de *Leucaena leucocephala* con énfasis en el contenido de metabolitos secundarios. Revista Electrónica de Veterinaria (REDVET). 2012, 13. Disponible en: <http://www.veterinaria.org/revistas/redvet/n111112/111213.pdf>. [Consultado: 5 de junio de 2017].
- [12]. McDonald P, Edwards RA, Greenhalgh JFD, Morgan CA. 1995. Animal Nutrition. Fifth Edition. Longman Scientific and Technical. 1995. 547.
- [13]. Minson PJ. Composición química y valor nutritivo de las leguminosas tropicales. En: Leguminosas Forrajeras Tropicales. FAO. Eds. Skerman, P.J., Cameron, D.E. y Riveros, F. Roma, Italia. 1991. 211-219.
- [14]. Pérez A, Montejo J, Iglesias JM, López O, Martín GJ, García DE, Milián I, Hernández A. *Tithonia diversifolia* (Hemsl.) A. Gray. Pastos y Forrajes. 2009, 32:1-15.
- [15]. García DE, Medina MG, Cova LJ, Clavero T, Torres A, Perdomo D, Santos O. Evaluación integral de recursos forrajeros para rumiantes en el estado Trujillo, Venezuela. Rev. Fac. Agron. (LUZ). 2009, 26: 555-582.
- [16]. González JC, Ayala A, Martínez E. Composición química de especies arbóreas con potencial forrajero de la región de tierra caliente, Michoacán, México. Rev. Cubana Cienc. Agríc. 2007, 41:87-93.
- [17]. Ríos CI. *Tithonia diversifolia* (Hemsl.) Gray, una planta con potencial para la producción sostenible en el trópico. En: Agroforestería para la Producción Animal en América Latina. Estudio FAO Producción y Sanidad Animal N° 143. (Eds.) Sánchez MD, Rosales M. FAO, Roma. Italia. 1999. 311-325.
- [18]. Ceballos A, Noguera RR, Bolívar DM, Posada SL. Comparación de las técnicas in situ de los sacos de nylon e in vitro (DaisyII) para estimar la cinética de degradación de alimentos para rumiantes. Livestock Research for Rural Development. 2008, 20. Disponible en: <http://www.lrrd.org/lrrd20/7/ceba20108.htm>. [Consultado 24 de mayo de 2017].
- [19]. García DE, Medina MG, Clavero T, Humbría J, Baldizán A, Domínguez C. Preferencia de árboles forrajeros por cabras en la zona baja de los andes Venezolanos. Revista Científica, FCV-LUZ. 2008, 18: 549-555.
- [20]. Verdecia DM, Ramírez JL, Bodas R, Leonard I, Giráldez FJ, Andrés S, Santana A, Herrera RS, López S. Rendimiento productivo y composición química del arbusto forrajero *Tithonia diversifolia* en una zona del valle del Cauto, Cuba. En: Pastos, paisajes culturales entre tradición y nuevos paradigmas del siglo XXI (eds. C. López, M.P. Rodríguez, A. San Miguel, F. Fernández y S. Roig), Sociedad Española para el estudio de los Pastos, Madrid. España. 2011. 613-618.
- [21]. Pineda M. 2004. Resúmenes de Fisiología vegetal. Servicios de publicaciones de la Universidad de Córdoba, Córdoba, España. 2004. 204. Disponible en: www.uco.es/publicaciones/revistas.html [Consultado 14 de octubre de 2016].
- [22]. García DE, Medina MG, Ojeda F, Humbría J, Domínguez CE, Baldizán A, Toral O. Variabilidad fitoquímica y repercusión antinutricional potencial en especies del género *Albizia*. Pastos y Forrajes. 2006, 29:231-241.

- [23]. Cao H, Hwang J, Chen X. Carbohydrate-containing natural products in medicinal chemistry. Opportunity, challenge and scope of natural products in Medicinal Chemistry. 2011, 37: 411-431.
- [24]. García C, Martínez A, Ortega JL, Castro F. Componentes químicos y su relación con las actividades biológicas de algunos extractos vegetales. Revista Química Viva. 2010, 9: 86-96.
- [25]. Benavides JE. La investigación en árboles forrajeros. En: Benavides JE. (Ed.) Árboles y Arbustos Forrajeros en América Central. Informe Téc. No. 236. CATIE. Turrialba, Costa Rica. 1994, 1: 3-21.
- [26]. Trugo LC, Farah A, Cabral L. Oligosaccharides distribution in Brazilian soya bean cultivars. Food Chem. 1988, 52: 385-387.



UEA
UNIVERSIDAD
ESTATAL AMAZÓNICA

MODECO 06

Diversidad Molecular, Producción de Energía y Ecosistemas, del Amazonas a los Océanos

invita a la a la comunidad universitaria de la UEA, instituciones de educación superior y estratégicas del Ecuador y ciudadanía en general, involucradas en áreas de:



- Conservación de Ecosistemas
- Producción de Energía
- Impacto Climático
- Desarrollo de Productos Naturales y Procesos Agroindustriales, entre otras.

Revisa las bases en el siguiente link:



<https://bit.ly/3dqvD2t>

CONVOCATORIA ABIERTA

Del 25/01/2021
hasta el 30/12/2021

#UEAesExcelencia

Presentar sus trabajos de investigación, para su difusión en el taller interuniversitario transatlántico, MODECO-06, sin costo alguno.

