

Proceeding Paper

# Effects of Chitosan Oligosaccharide Lactate on Growth and Overwintering of Evergreen Fern *Cyrtomium fortunei* var. *clivicola* †

Piotr Salachna <sup>1,\*</sup>, Anna Pietrak <sup>1</sup> and Łukasz Łopusiewicz <sup>2</sup>

<sup>1</sup> Department of Horticulture, West Pomeranian University of Technology in Szczecin, Papieża Pawła VI 3, 71-459 Szczecin, Poland; pa37778@zut.edu.pl

<sup>2</sup> Center of Bioimmobilisation and Innovative Packaging Materials, Faculty of Food Sciences and Fisheries, West Pomeranian University of Technology in Szczecin, Janickiego 35, 71-270 Szczecin, Poland; lukasz.lopusicwicz@zut.edu.pl

\* Correspondence: piotr.salachna@zut.edu.pl; Tel.: +48-914-496-359

† Presented at the 2nd International Electronic Conference on Plant Sciences—10th Anniversary of Journal Plants, 1–15 December 2021; Available online: <https://iecps2021.sciforum.net/>.

**Abstract:** Chitosan derivatives with low molecular weight can enhance plant growth and improve tolerance to various stress. Hardy ferns form a group of attractive garden perennials with unknown response to plant biostimulants. Cold winter temperature limits the growth and reduces the decorative value of evergreen fern species. This study was carried out to explore the effects of chitosan oligosaccharide lactate (COL) with Mn = 5000 on growth of evergreen fern *Cyrtomium fortunei* var. *clivicola*. COL sprayed at 50 and 100 mg/L significantly increased height of plant and fresh weight of the above-ground and the underground parts of *C. fortunei* var. *clivicola* compared to control plants. Moreover, ferns sprayed with 50 and 100 mg/L COL also had higher leaf total chlorophyll contents and value score. These results indicate that COL improved *C. fortunei* var. *clivicola* growth and overwintering and may be used for high quality hardy ferns production.

**Keywords:** hardy ferns; biostimulants; oligochitosan; low temperature stress

**Citation:** Salachna, P.; Pietrak, A.; Łopusiewicz, Ł. Effects of Chitosan Oligosaccharide Lactate on Growth and Overwintering of Evergreen Fern *Cyrtomium fortunei* var. *clivicola*. *Biol. Life Sci. Forum* **2021**, *1*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Iker Aranjuelo

Published: 2 December 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The wide demand for a plant biostimulants inspires the search for sources of innovative natural substances with high biostimulant action. Particular attention is paid to natural polysaccharides, which are eco-friendly, biocompatible, non-toxic, reactively bioactive and inexpensive. Products of polysaccharides with the low molecular weight, and therefore, with different physico-chemical properties, can affect the growth and physiological activity of plants to variable degrees [1,2]. Chitosan is an example of a widespread biostimulant obtained in the process of chitin de-N-acetylation. It has been proven that chitosan, particularly with low molecular weight, positively affects the growth and yield quality [3], regulates physiological and metabolic processes [4], and also increases plant tolerance to abiotic stress [5,6]. Chitosan oligosaccharide with a low-molecular-weight obtained by degradation of chitosan by lactate could be a promising approach in the horticultural production. This product has outstanding water-solubility and high biological activity, therefore it may be easily absorbed and utilized by plants [7]

The offer of hardy ornamental plants continues to be enriched, because the novelties always enjoy interest and higher prices are obtained. Hardy ferns are perennials still little known and very attractive [8]. *Cyrtomium fortunei*, commonly called holly fern, is an easy-to-grow evergreen fern native to Japan, Korea and China. *Cyrtomium fortunei* var. *clivicola* differs from the species by its more compact size, with lanceolate and yellow green leaves

held on nearly horizontal, arching fronds. Cold winter temperature may limit the growth and reduce the decorative value of evergreen fern species. The available literature lacks detailed data on the effects of plant-growth-regulating compounds, including biostimulants on the growth and winter hardiness of fern.

Hence, the aim of the present study is to evaluate after winter the morphological features of *C. fortunei* var. *clivicola* plants pretreated with chitosan oligosaccharide lactate (COL) with low molecular weight.

## 2. Materials and Methods

Two-years-old rhizomes of *Cyrtomium fortunei* var. *clivicola* were planted in 1.7 L pots and grown from summer 2020 to spring 2021 in a plastic tunnel of the Department of Horticulture, West Pomeranian University of Technology in Szczecin (53°25' N, 14°32' E, 25 m asl., sub-zone 7a USDA). The growing medium consisted of peat substrate (pH 6.0) added with PG Mix fertilizer at a dose of 1 g/dm<sup>3</sup>. In October 2020 the plants were sprayed twice with 50 or 100 mg/L solution of chitosan oligosaccharide lactate (C<sub>12</sub>H<sub>24</sub>N<sub>2</sub>O<sub>9</sub>)<sub>n</sub> with the number average molecular weight (Mn) = 5000. The control plants were sprayed with water. Each treatment included 20 rhizomes, planted in four replicates of 5 rhizomes. Impact of chitosan oligosaccharide lactate on plant growth was assessed after winter. In March 2021 were measured plant height from the ground level to its highest point, plant width at its widest point, leaf total chlorophyll contents in SPAD (Soil Plant Analysis Development) units with the Chlorophyll Meter SPAD-502 optical apparatus (Konica-Minolta Corporation, Osaka, Japan) determined from 5 points of leaves, and leaves fresh weight of leaves and rhizomes per plant. The visual score was rated on a scale of 1–5 by three autonomous investigators. The results were statistically analyzed using a one-way analysis of variance (ANOVA) and the means compared by Tukey's multiple comparison test.

## 3. Results and Discussion

Foliar application of COL improved the all growth attributes of plants except for leaves number (Table 1, Figure 1). The fern plants sprayed with COL at 50 and 100 mg/l showed a significant increase in plant height (by 25.4% and 28.5%, respectively) in comparison to controls. COL at 50 and 100 mg/l also improved leaf total chlorophyll contents (by 30.5% and 30.1%, respectively). COL at 100 mg/L did not change diameter of plant. Nevertheless, COL at 50 mg/L increased diameter of plant by 15.3% versus controls not treated (Table 1).

**Table 1.** Effect of foliar applications of chitosan oligosaccharide lactate on plant height, plant diameter, number of leaves per plant and leaf total chlorophyll contents (SPAD) of *Cyrtomium fortunei* var. *clivicola* plants after overwintered in a plastic tunnel.

Chitosan (mg/L)	Plant Height (cm)	Plant Diameter (cm)	Leaves (no.)	SPAD
Control (water)	13.0 b <sup>1</sup>	48.3 b	30.3 a	22.6 b
50	16.3 a	55.7 a	28.3 a	29.5 a
100	16.7 a	45.8 b	30.8 a	29.4 a

<sup>1</sup> Mean values in each column followed by a different lowercase letter are significantly different by Tukey's least significant difference test (LSD) at  $p < 0.05$ .



**Figure 1.** *Cyrtomium fortunei* var. *clivicola* at six months after treatment: (left)—untreated control; (right)—50 mg/L chitosan oligosaccharide lactate.

COL applications at 50 and 100 mg/L significantly enhanced fresh weight of leaves (by 24.8% and 27.1%, respectively) and rhizomes (by 53.4% and 14%, respectively) compared to control. The plants sprayed with COL at 50 mg/L exhibited the greatest fresh rhizomes weight (50.3 g). COL treatments also influenced visual score of plant after overwintered in a plastic tunnel. The fern plants sprayed with 50 and 100 mg/L COL had significantly greater visual score in comparison with untreated plants (Table 2).

**Table 2.** Effect of foliar applications of chitosan oligosaccharide lactate on fresh weight of leaves and rhizomes and visual score of *Cyrtomium fortunei* var. *clivicola* plants after overwintered in a plastic tunnel.

Chitosan (mg/L)	Leaves Fresh Weight (g)	Rhizomes Fresh Weight (g)	Visual Score
Control (water)	26.6 b <sup>1</sup>	32.8 c	3.7 b
50	33.2 a	50.3 a	4.5 a
100	33.8 a	37.4 b	4.9 a

<sup>1</sup> Mean values in each column followed by a different lowercase letter are significantly different by Tukey's least significant difference test (LSD) at  $p < 0.05$

The obtained results indicated the stimulating effect of COL on plant height and diameter, fresh biomass, and visual score of evergreen fern *Cyrtomium fortunei* var. *clivicola* after winter. In addition, the plants treated with COL had greater greenness index at six months after treatment. Numerous other studies have also reported that natural polysaccharides such as chitosan and their derivatives improved the growth attributes and enhanced the levels of chlorophylls in plants [2–4]. It can thus be supposed that chitosan improves the efficiency of water and individual elements' uptake, which in turn is associated with the intensification of plant growth [2,3,9].

#### 4. Conclusions

Our report is the first to show that chitosan in form of oligosaccharide lactate affects fern growth and physiological processes. The results described in the present work not only supplement the current state of knowledge on the impact of oligochitosan on plant growth, but can also contribute to the improvement and popularization of *Cyrtomium fortunei* var. *clivicola* cultivation and can be used in the production of other hardy ferns.

**Author Contributions:** Conceptualization, P.S.; methodology, P.S.; investigation, P.S., A.P. and Ł.Ł.; data curation, P.S. and A.P.; writing; P.S., A.P. and Ł.Ł.; All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Data is contained within the article.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Yang, X.-J.; Chen, Y.; Hu, Z.; Ma, S.; Zhang, J.; Shen, H. Alginate Oligosaccharides Alleviate the Damage of Rice Leaves Caused by Acid Rain and High Temperature. *Agronomy* **2021**, *11*, 500.
2. Yuan, X.; Zheng, J.; Jiao, S.; Cheng, G.; Feng, C.; Du, Y.; Liu, H. A review on the preparation of chitosan oligosaccharides and application to human health, animal husbandry and agricultural production. *Carbohydr. Polym.* **2019**, *220*, 60–70.
3. Salachna, P.; Grzeszczuk, M.; Soból, M. Effects of Chitooligosaccharide Coating Combined with Selected Ionic Polymers on the Stimulation of *Ornithogalum saundersiae* Growth. *Molecules* **2017**, *22*, 1903.
4. Salachna, P.; Pietrak, A. Evaluation of Carrageenan, Xanthan Gum and Depolymerized Chitosan Based Coatings for Pineapple Lily Plant Production. *Horticulturae* **2021**, *7*, 19.
5. Hidangmayum, A.; Dwivedi, P. Chitosan Based Nanoformulation for Sustainable Agriculture with Special Reference to Abiotic Stress: A Review. *J. Polym. Environ.* **2021**, 1–20.
6. Li, Y.; Zhang, Q.; Ou, L.; Ji, D.; Liu, T.; Lan, R.; Li, X.; Jin, L. Response to the Cold Stress Signaling of the Tea Plant (*Camellia sinensis*) Elicited by Chitosan Oligosaccharide. *Agronomy* **2020**, *10*, 915.
7. Available online: <https://agfuse.com/article/application-of-chitosan-oligosaccharide-in-agriculture> (accessed on).
8. Salachna, P.; Piechocki, R. Salinity Tolerance of Four Hardy Ferns from the Genus *Dryopteris* Adans. Grown under Different Light Conditions. *Agronomy* **2021**, *11*, 49.
9. Ha, N.M.C.; Nguyen, T.H.; Wang, S.-L.; Nguyen, A.D. Preparation of NPK nanofertilizer based on chitosan nanoparticles and its effect on biophysical characteristics and growth of coffee in green house. *Res. Chem. Intermed.* **2019**, *45*, 51–63.