

Proceeding Paper

Evaluating Seven Macadamia Seedling and Cutting Rootstocks for Their Effect on Scion Growth [†]

Pragya Dhakal Poudel ^{*}, Max Cowan, Bruce Topp and Mobashwer Alam

Queensland Alliance for Agriculture and Food Innovation, University of Queensland, Maroochy Research Facility, 47 Mayers Rd, Nambour, QLD 4560, Australia; max.cowan@uq.edu.au (M.C.); b.topp@uq.edu.au (B.T.); m.alam@uq.edu.au (M.A.)

^{*} Correspondence: p.dhakalpoudel@uq.edu.au

[†] Presented at the 2nd International Electronic Conference on Plant Sciences—10th Anniversary of Journal Plants, 1–15 December 2021; Available online: <https://iecps2021.sciforum.net>

Abstract: This study evaluated the variability of seven seedling and cutting rootstock genotypes of macadamia for their effect on scion growth. Scion height varied from 384 cm to 524 cm in cutting and 378 cm to 450 cm in seedling rootstocks. This difference was minimal at an early age (year 1–3) while being statistically significant at year 4. A highly significant positive correlation was identified between grafted and ungrafted rootstocks for canopy area ($r = 0.81$, $p < 0.001$) while scion height correlated moderately with rootstock height ($r = 0.36$, $p < 0.001$). A wider genetic range is possibly required to be studied for precise conclusions.

Keywords: Macadamia; vigour; rootstock; scion; propagation; seedling; cutting

Citation: Poudel, P.D.; Cowan, M.; Topp, B.; Alam, M. Evaluating Seven Macadamia Seedling and Cutting Rootstocks for Their Effect on Scion Growth. *Biol. Life Sci. Forum* **2021**, *1*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Feibo Wu

Published: 2 December 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Large tree size is one of the key obstacles of building an efficient orchard system in macadamia (*Macadamia integrifolia*, *M. tetraphylla*, and hybrids). Vigorous trees restrict high-density planting in commercial orchards and limits tree productivity due to intra-tree resource competition [1]. The introduction of vigour-reducing rootstocks in many horticultural crops have largely contributed to high planting density, better yield per hectare and therefore higher profitability for growers [2]. As a part of a dual plant system, rootstocks translocate water and nutrients [3] to the above ground scion and thereby regulate resource supply for growth and development of vegetative and reproductive parts. Rootstocks with strong root systems are known to provide resistance to several biotic and abiotic factors and increase adaptability to adverse environmental conditions [4]. Thus, rootstocks in tree crops like macadamia play a vital role in crop growth and vigour.

Commercial macadamias in Australia are mostly propagated through grafted seedlings while cutting (clonal) rootstocks are rarely used [5]. Seedling rootstocks are mostly preferred because of their quality root systems and ease of production in comparison to clonal rootstocks [6]. 'H2' seedlings have been the most preferred rootstock for propagation methods in Australia since the early 1990s [7] due to their uniformity and thick stem diameter which allows rapid graft formation [8]. However, seedling rootstocks are seen to influence tree nutrition which eventually leads to variation in tree size and vigour of grafted orchards [9]. This variation can be reduced by clonal rootstocks, a method thought to be more effective and cost-efficient for rapid rootstock propagation [6]. Clonal rootstock of a hybrid cultivar 'Beaumont' has been preferred in South Africa due to its high strike success and vigorous nursery growth [10]. However, one study reported that trees prop-

agated through cutting rootstocks are vulnerable to strong wind [6]. Identification of rootstocks contributing to improved productivity and resilience to biotic and climatic factors is important for industry expansion.

Study on rootstock propagation methods and its effect on scion growth in macadamia is limited. In a concurrent study by Alam et al. [11], 29 rootstock genotypes of macadamia were studied and compared for early nursery growth in both seedling and cutting rootstocks. In another study, Neal et al. [7] studied 12 macadamia cultivars and identified no significant interaction between rootstock genotype and propagation methods for tree height. Another study that evaluated the performance of seedling and cutting rootstocks found no significant variation in tree size due to rootstock effect although a significant scion effect was identified [12]. This preliminary study therefore aimed to identify the variation in growth traits and growth rates of a common scion due to the effect of seedling and cutting rootstocks of seven macadamia genotypes.

2. Materials and Methods

2.1. Rootstock Types and Plantation

Seedling and cutting rootstocks of seven macadamia genotypes ('A268', 'A4', 'BHI1', 'BHY1', 'Beaumont', '842' and 'H2') were selected from an existing macadamia rootstock trial planted in 2017 at the Maroochy Research Facility, Queensland Department of Agriculture and Fisheries Nambour QLD [13]. A common scion 'HAES 741' was grafted onto the rootstocks. Both grafted and ungrafted trees, planted side by side, were evaluated for this study.

2.2. Phenotyping

Phenotyping was performed for Scion height (ScHGT) and Canopy Area (CA). These traits were measured for 5 years following the tree establishment.

Growth rates (*GR*) for each parameter over 5 years were calculated using the following formula:

$$GR = \frac{dx}{dt} \quad (1)$$

where *GR* = Growth rate, *dx* = difference in growth over *dt* and *dt* = time between measurements (in days).

2.3. Statistical Analysis

Growth parameters were analysed using a REML (Restricted maximum likelihood) mixed model in Genstat-21 [14]. A combined analysis of seedlings and cuttings was performed for growth traits. Analysis of growth rates included combined and separate seedling and cutting analyses. Phenotypic correlations between grafted scion and ungrafted rootstock traits were estimated.

3. Results and Discussions

3.1. Effect of Rootstocks on Variability in Growth Traits

Combined analysis of both seedling and cutting rootstock effect shows no significant variation between rootstock genotypes for ScHGT and CA, while a significant variation between seedlings and cuttings for ScHGT was identified in Y4 (Table 1)—cuttings had higher average ScHGT than the seedlings of most genotypes, except for 'A4' (Figure 1). For cuttings, 'A4' (384 cm) had the lowest ScHGT, while 'H2' (524 cm) had the highest values. Similarly in seedlings, ScHGT varied from 378 cm to 450 cm with 'BHI1' and 'A4' having the highest and lowest values respectively. No significant variation between rootstock genotypes and types for CA was observed.

Table 1. F probabilities from REML mixed model analysis of growth traits (Age4) of seedlings and cutting rootstock effects.

Components	ScHGT	CA
Genotype	0.365	0.353
Type (Seedling vs Cutting)	0.025	0.196
Genotype × Type	0.15	0.342




Figure 1. Effect of rootstock propagation types and genotypes on variability in scion height. Error bars show the standard errors (SE) of each mean. Mean bars on x-axis shows average value across the genotypes.

Similarly, no significant variation for rootstock effect on ScHGT were observed in early ages (Y1–Y3) (data not shown), while the effect was significant at Y4. In a similar trial of 12 cultivars grafted as clonal and seedling rootstocks, Hardner and McConchie [12] found no significant effect of rootstocks on field growth in 2 years old trees. These findings, along with the result of the current study suggest that rootstock effect on the scion may not become apparent until later stages of maturity.

3.2. Effect of Rootstocks on Variability in Growth Rates

Initial combined analysis of seedlings and cuttings identified a significant variation between the propagation types for GR (Y0–Y4) of ScHGT and CA (Table 2). Cuttings had higher average GR for ScHGT and CA than seedlings (Figure 2). A significant Genotype X Type interaction existed for rootstock effect on GRCA. In terms of the genotypes ‘A268’, ‘BHI1’, ‘Beau’ and ‘H2’, cuttings had higher GRCA than their respective seedlings. Inversely, seedlings of ‘BHY1’, ‘A4’ and ‘842’ had higher GRCA than their cuttings (Figure 2b).

Table 2. F probabilities from REML mixed model analysis of growth rates (Y0–Y4) of seedling and cutting rootstocks effect.

Components	Seedlings Vs Cuttings		Seedlings Only		Cuttings Only	
	GRScHGT	GRCA	GRScHGT	GRCA	GRScHGT	GRCA
Genotype	0.337	0.957	0.526	0.212	0.682	0.324
Type (Seedling Vs Cuttings)	<0.001	0.036	-	-	-	-
Genotype × Type	0.214	0.015	-	-	-	-




Figure 2. Effect of rootstock propagation types and genotypes on variability in growth rates (Y0–Y4) (a) Growth Rate scion height (b) Growth Rate of canopy area. Error bars show the standard errors (SE) of each mean. Mean bars on x-axis shows average value across the genotypes.

This result is in contrast with several other findings which indicate that scions grafted on seedling rootstocks grow to be more vigorous than those on cutting rootstocks [6,12,15,16]. Conversely, a pairwise comparison of seedling and cutting rootstocks of 19 genotypes by Alam et al. [11] on early nursery growth reported similar results where relative growth rates for height and diameter of cutting rootstocks were higher than those of seedling rootstocks. The contrasting results could be due to the difference in genotypes selected for each study. Cutting propagated rootstocks in this study consisted of superior genotypes like ‘Beaumont’ and ‘A268’ with superior strike rate, growth rate and growth budding success when propagated as cuttings [12] while the seedlings are open pollinated progeny and may have shown variability due to the differences in pollen source. The significant Genotype X Type interaction showing higher GR for cuttings of ‘Beaumont’ and ‘A268’ also justifies this contrasting result.

No significant variation between rootstock types and genotypes for GRScHGT and GRCA were observed in first three consecutive years of measurement (data not shown). This study identified that growth rates measured over one year did not vary significantly between seedling and cutting rootstocks while, a significant difference was observed when a large gap of Y0–Y4 was considered. This again suggests that rootstock effect on scion growth due to propagation type may not be visible at an early age and requires a longer period of growth to be evaluated. Further monitoring over the coming years is required to reach into a more accurate conclusion.

A separate analysis of each propagation type showed no significant difference within seedling and cutting genotypes for growth rates (Table 2). This non-significant variation among the genotypes could be because the genotypes do not represent a large range of vigour. As this study did not use any dwarf genotypes, inclusion of those genotypes to increase the range of genetic material could better help to understand any variation between genotypes, if it exists.

3.3. Phenotypic Correlation between Ungrafted and Grafted Rootstock Genotypes

To investigate the transmission of trait expression from rootstock to scion, we conducted phenotypic correlations between scion growth traits with that of ungrafted rootstocks of the same genotype. A strong positive correlation ($r = 0.81$, $p < 0.001$) was identified between grafted scions and ungrafted rootstocks for CA (Table 3). Rootstock height of ungrafted genotypes were positively and moderately correlated with scion height of

the grafted ones ($r = 0.36$, $p < 0.001$). This shows that rootstock vigour is closely related with scion vigour. Thus, a scion grafted onto a vigorous rootstock is likely to grow to be vigorous and selection of rootstocks can be made accordingly. This has been widely studied and identified in many other horticultural crops [17]. However, this study consists of a single scion grafted onto all rootstock genotypes. Hence, a large combination of different rootstock and scion cultivars is recommended for further studies.

Table 3. Pearson's phenotypic correlations of growth parameters between grafted and ungrafted rootstock genotypes.

	Canopy Area-Ungrafted	Scion Height-Grafted
Canopy Area-Grafted	0.81 ***	-
Rootstock Height-Ungrafted	-	0.36 ***

*** $p < 0.001$.

4. Conclusions

This study found a significant variation in the effect of rootstock types (seedlings and cuttings) on scion vigour. Scions grown on cutting rootstocks were generally more vigorous than those on seedling rootstocks. Results here also indicate that rootstock effect on scion growth is likely to become more apparent with greater tree maturity. In addition, a strong positive relationship between grafted and ungrafted rootstock genotypes for scion height indicate that scion vigour can be predicted from the vigour of the rootstocks. Since this study was limited to few genotypes, investigating a more diverse range of germplasm and assessing growth until later stages of maturity is required for improved accuracy in future studies.

Author Contributions: Experimentation and analysis, P.D.P.; writing-original draft preparation, P.D.P.; writing- review and editing, M.C., B.T. and M.A. Supervision, B.T. and M.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research has been funded by The University of Queensland (UQ) and Hort Innovation (HI) Australia. UQ provided a higher degree research scholarship for PD. HI provided funding through MC14000, MC19000, AI18000, using the macadamia research and development levy and contributions from the Australian Government.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data availability statement: Data will be available upon request, please contact m.alam@uq.edu.au.

Acknowledgments: We acknowledge the administrative and research funding supports of UQ. We also acknowledge Hort Innovation, the grower owns, non-profit research and development corporation for Australian horticulture, for providing funding for this project.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in writing of the manuscript, or in the decision to publish the results.

References

1. McFadyen, L.M.; Robertson, D.; Sedgley, M.; Kristiansen, P.; Olesen, T. Post-pruning shoot growth increases fruit abscission and reduces stem carbohydrates and yield in macadamia. *Ann. Bot.* **2011**, *107*, 993–1001, doi:10.1093/aob/mcr026.
2. Robinson, T. Advances in Apple Culture Worldwide. *Revista Brasileira de Fruticultura* **2011**, *33*, 37–47.
3. Ferree, C. D.; Carlson, R.F. Apple Rootstocks. In *Rootstocks for Fruit Crops*, Rom, R.C., Carlson, R.F., Eds.; Wiley: Hoboken, NJ, USA, 1987; pp. 107–144.
4. Martínez-Cuenca, M.-R.; Primo-Capella, A.; Forner-Giner, M.A. Influence of Rootstock on Citrus Tree Growth: Effects on Photosynthesis and Carbohydrate Distribution, Plant Size, Yield, Fruit Quality, and Dwarfing Genotypes. *Plant Growth* **2016**, *16*, 107, doi:10.5772/64825.

5. Russell, D.; Neal, J.; Mayer, R.; Bell, D.; Topp, B. Variation of cutting rooting ability in cultivated and wild species of Macadamia. In Proceedings of the XXIX International Horticultural Congress on Horticulture: Sustaining Lives, Livelihoods and Landscapes (IHC2014): 1109, Brisbane, Australia, Aug 17, 2014.
6. Bell, H.; Bell, D. High density plantings of cuttings in macadamia. *Aust. Macadamia Soc. News Bull* **1993**, *20*, 23–24.
7. Neal, J.; Kelly, A.; Hardner, C.; McConchie, C.; Topp, B. *Preliminary Evaluation of Macadamia Rootstocks for Yield and Tree Height*; International Society for Horticultural Science (ISHS): Leuven, Belgium, 2016.
8. Stephenson, R. Macadamia nut. In *Fruits: Tropical and Subtropical*; Naya Udjog, Calcutta, India, 1990.
9. Nagao, M.A.; Hirae, H.H.; Stephenson, R. Macadamia: Cultivation and physiology. *Crit. Rev. Plant Sci.* **1992**, *10*, 441–470, doi:10.1080/07352689209382321.
10. Wiid, M.; Hobson, L. Why macadamia cultivar Beaumont produces high yields under ultra high density conditions in South Africa. In Challenges for horticulture in the tropics. In Proceedings of the 3rd Australian Society of Horticultural Science Conference, Broadbeach, Australia, 18–22 August 1996.
11. Alam, M.; Wilkie, J.; Topp, B. Early growth and graft success in macadamia seedling and cutting rootstocks. In Proceedings of the International Symposia on Tropical and Temperate Horticulture-ISTTH2016 1205, Cairns, Queensland, Australia, 30 August 2016.
12. Hardner, C.; McConchie, C. The effect of rootstock on propagation success and early field performance. In *Paper presented at: 3rd International Macadamia Symposium*; Macadamia Society: Sao Pedro, Brazil, 2006.
13. Alam, M.; Wilkie, J.; Kelly, A.; Hardner, C.; Topp, B. Genetic diversity and variability in graft success in Australian Macadamia rootstocks. In Proceedings of the International Macadamia Research Symposium, Honolulu, HI, USA, 13–14 September 2017.
14. Payne, R.; Welham, S.; Harding, S. *A Guide to REML in GenStat*; VSN International, Hemel Hempstead: Hertfordshire, UK, 2011.
15. Dogra, K.; Kour, K.; Kumar, R.; Bakshi, P.; Kumar, V. Graft-Incompatibility in Horticultural Crops. *Int. J. Curr. Microbiol. Appl. Sci.* **2018**, *7*, 1805–1820, doi:10.20546/ijcmas.2018.702.218.
16. Huett, D.O. Macadamia physiology review: A canopy light response study and literature review. *Aust. J. Agric. Res.* **2004**, *55*, 609–624, doi:10.1071/ar03180.
17. Jayswal, D.; Lal, N. Rootstock and Scion Relationship in Fruit Crops. *Agriallis*, **2020**, *2*(11), 10–16.