

Abstract

# Small Mammal Diversity and Abundance in Commercial Orchards in Relation to Habitat and Agricultural Factors <sup>†</sup>

Vitalijus Stirke, Linas Balčiauskas and Laima Balčiauskienė

Nature Research Centre, Akademijos 2, Vilnius, Lithuania; [linas.balciauskas@gamtc.lt\(L.B.\)](mailto:linas.balciauskas@gamtc.lt(L.B.)); [laima.balciauskiene@gamtc.lt\(L.B.\)](mailto:laima.balciauskiene@gamtc.lt(L.B.))

\* Correspondence: [vitalijus.stirke@gamtc.lt](mailto:vitalijus.stirke@gamtc.lt)

<sup>†</sup> Presented at the 2nd International Electronic Conference on Diversity (IECD 2022)—New Insights into the Biodiversity of Plants, Animals and Microbes, 1–15 March 2022; Available online: <https://iecd2022.sciforum.net/>.

**Abstract:** Diversity of small mammal communities (SMC) show sustainability of habitats, especially agricultural ones. Gathered over three years, data from 18 sites in Lithuania were used to analyse factors related to several dependent parameters, namely diversity (Shannon's H, dominance index and number of species trapped), the relative abundance of species in SMC and the abundances and proportions of the four most numerous species, specifically common vole, striped field mouse, yellow-necked mouse and bank vole. Using the General Linear Model, we assessed the influence of habitat type (commercial orchards, berry plantations, control meadows and control forests with at least two of these present at every investigation site), age of the orchard or plantation, intensity of agriculture, season and location (central, northern, eastern, southern and western parts of the country). To control temporal data variability, year was used as a continuous predictor. The model was valid and explained 14–31% of the listed parameters with  $p < 0.005$  or higher, with the exception of the dominance index and the proportion of the common vole. Time factor (year and season,  $p < 0.001$ ), intensity of agriculture and site location ( $p < 0.05$ ) had the highest influences on the model, while that of habitat type and its age were not significant. Univariate results suggest that old commercial orchards with low intensities of agricultural practice play a role in maintaining diversity and abundance of SMC.

Academic Editor(s): Matthieu Chauvat

Published: 15 March 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).