

# Green Biomonitoring Systems for Air Pollution <sup>†</sup>

Andreea Cozea\* and Gheorghita Tanase

National Research and Development Institute for Industrial Ecology ECOIND, 57-73 Drumul Podu Dambovitei Street, district 6, 060652, Bucharest, Romania; tanasegeoeco@gmail.com

\* Correspondence: andreea.cozea@yahoo.com

<sup>†</sup> Presented at the 1st International Electronic Conference on Processes: Processes System Innovation, 17–31 May 2022; Available online: <https://ecp2022.sciforum.net>.

**Abstract:** Human activities have led to environmental pollution also industrial growth and the gradual transition to urbanization are the main causes of the growth of different types of environmental pollutants and the continuation of studies on air quality biomonitoring in the context of “green revolution” have renew the way to use “green tools” using higher plants with specific capabilities-as biomonitors.

**Keywords:** biomonitoring; environmental pollution; enzymes; green revolution.

## 1. Introduction

Urbanization, industry and transport development causes high concentrations of air pollutants emissions, which lead to harmful effects on nature especially for all living organisms. After awareness of the danger of the ecosystems imbalance, the sought was to find ways to monitor them related with international policy efforts to limit air emissions. All that actions has been transposed into plans and agreements at international level and involved into a global movement knew as “green revolution”. Plants can also indicate pollution load in a particular area via alterations in physiological parameters such as pigment degradation, membrane damage, production of antioxidant metabolites and change in anti-oxidant enzyme activities. The aim of this first environment surveillance is to detect possible changes in the functioning and composition of biological systems, which could eventually lead to disorganization, collapse, or move them in an evolutionary unfavorable direction. Early warnings of the specialists allows policy decision makers to take action for remedy the situation, before irreversible negative effects to appear. The environment monitoring could be instrumental or biological. Biological monitoring or biomonitoring may replace or supplement instrumental monitoring and at the international level, the use of living organisms as “tools” in determining the quality of the environment in general and air contamination in particular and is in continuously increasing, [1].

As for the pollution “green” indicators, they are of two types: sensitive species, which indicate the presence of a pollutant by the appearance of lesions or malformations and accumulating species.

Most important reasons for using bioindicators wore: demonstrating how the biological effect of pollutants is integrated with the sensitivity of biological sensors and environmental (climatic) factors; through extrapolation of toxic hazard from vegetal to human, showing the toxic potential on humans; the possibility of standardization of working methods, [2].

In this paper we have conducted our own studies on air quality biomonitoring with plants in which the aim was to monitor air pollutants as dangerous factors to living organisms because of their high oxidative potential. So, the studies costing in the analysis of antioxidant compounds that are represent as structural class of chemicals, with a wide range of biological functions, with the role of inhibiting free radicals. Thus, a low level of

**Citation:** Lastname, F.; Lastname, F.; Lastname, F. Title. *Proceedings* **2022**, *69*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Lastname

Published: date

**Publisher’s Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

inhibition of free radicals indicates a reduced ability to inhibit them and increased sensitivity to leaf diseases (necrosis). Green plants can also act as indicators of air pollution by detectable metabolic product of the pollutant in their tissues or measuring indirect effects of pollutants based by the “built-in” ability to have an antioxidant response system to protect themselves from the damage caused by oxidative stress resulting from pollutant exposure. Only when the balance between the amount of antioxidant compounds and that of free radicals (caused by pollution) is altered, appear does the activation of the plant's response system, [3].

A correlation between the level of reactive oxidative system (ROS) in the cell and the activity of enzymes helps us to deduce that the antioxidant mechanism of the plant is responsible for combating oxidative stress due in this case to the stress factor: pollution. Based on that results by the experimental used in the study we developed our knowledge's in biomonitoring and we consider that is the basis for the development of new methods of use and study with application in environment assessments.

One of the aims was to change the appearance and biochemical composition of plants exposed to various pollutants compared to those not exposed, [4].

In order for them to be used in practice, species capable of providing the information necessary for the purpose of monitoring was a must in chosen plants. Sensitivity and response of plants to air pollutants is variable. The plant species which are more sensitive act as biological indicators of air. Plant material selected was: *Nicotiana tabacum*, *Petunia hybrida*, *Ricinus comunis*, *Trifolium pretense*.

In our case, the aim was to determine the antioxidant capacity of the extracts from the samples of plants exposed to pollutants compared to the control ones, in order to highlight that the degree of inhibition of free radicals decreases in the exposed plants compared to the control ones. Exposure locations used:

- 1- Site with incinerator coded Exposure site 1;
  - 2 -Site with poultry farm coded Exposure site 2;
  - 3- Site with slaughterhouse coded Exposure site 3.
- 2- Fumigation chambers.

Air pollutants generate reactive oxygen species (ROS) that leads to damage at cellular and molecular levels, [5]. Damaging effects induced by oxidative stress may get minimized by various enzymatic and non-enzymatic anti-oxidant defense mechanisms as has been suggested, [6].

The classical methods of biochemistry, the analysis of some antioxidant compounds and combined with those of biomonitoring have led to the completion of knowledge about the ways in which plants act on pollution control.

ROS OXYGEN REACTIVE SPECIES are a normal product of plant cell metabolism. A variety of environmental stresses lead to excessive ROS production, which causes oxidative damage and can eventually cause cell death. The most common reactive oxygen species are:  $O_2$ ,  $O_2$ ,  $H_2O_2$ ,  $OH$ . To avoid oxidative damage, higher plants improve their natural antioxidant defense system.

One of non-enzymatic compounds are polyphenols. Excess of phenols are considered important biomarkers for the phytotoxic effects of heavy metals and other pollutants. Increasing the production of phenolic compounds is a capability of the plant's defense mechanism. Phenolic compounds are of particular interest because of their involvement in the stress response, such as intra and / or interspecific competition or air pollution, [7]. Polyphenols are produced in the cytoplasm and form droplets in vacuoles that will later evolve into a single vacuole. If the excess polyphenols degenerate in the cytoplasm, the organs disappear and eventually the release of vacuolar contents leads to the death of mesophilic cells, [8].

Thus, in our study the amount of polyphenols in the exposed plants was analyzed compared to the Control plants.

The Antioxidant activity is determinate in our case by SOD superoxidismutase activity. The role of antioxidants is to destroy the free radicals of the cell that have a negative

effect on living organisms. A special role in neutralizing the effects of oxidative stress belongs to the enzyme superoxidismutase (SOD). Superoxidismutase (SOD) is a metal enzymes with subunit structural organization. SOD is the main regulator of oxidation processes in the cell. This enzyme catalyzes the recombination reaction of O<sub>2</sub> radicals.

Biochemical determinations of bioindicator samples can provide important information on increasing pollution.

## 2. Methods

### Determination of the total polyphenol content

The total content of polyphenolic compounds in plant extracts was determined colorimetrically using Folin-Ciocalteu reagent. Folin-Ciocalteu, also called the gallic acid equivalence method (GAE), used for colorimetric analysis of phenolic and polyphenolic antioxidants, [9].

A sample of 0.2 ml of plant extract was added to a test tube and mixed with 2 ml of Folin-Ciocalteu reagent, after 5 minutes of reaction, 1.8 ml of sodium carbonate (7.5%) was added. The absorbance was measured at 750 nm using a UV-VIS spectrophotometer.

The curve was established for analysis using gallic acid. The polyphenolic content was determined using the standard gallic acid calibration curve and expressed in mg of gallic acid equivalents.

### Determination of antioxidant capacity by determining SOD

The method described by Winterbourn et al. is based on the ability of superoxide dismutase to inhibit the reduction of tetrazolium-Nitro Blue Tetrazolium (Nitro BT) salt with superoxide radicals, [10]. An enzymatic unit is the amount of enzyme that produces 50% inhibition under standard conditions respectively speed and reaction temperature. Superoxide radicals are generated in the reaction medium by photoreduction of riboflavin. In parallel, a blank and a sample were performed, excepted for the enzymatic solution. The blank and the sample were illuminated for 5 minutes by a neon bulb, after which the extinction was read at 560 nm.

## 3. Results and Discussion

Following are presented the determination of polyphenols expressed in gallic acid and the following results were obtained (Table 1):

**Table 1.** Polyphenols expressed in gallic acid mg/g dry substance.

	<b>Nicotiana Tabacum</b>	<b>Petunia Hibrida</b>	<b>Ricinus Comunis</b>	<b>Trifolium Pratense</b>
Control	1.12	0.62	0.45	0.79
Exposure Sit 1	2.17	2.49	3.25	1.25
Exposure Sit 2	2.76	2.43	5.39	1.43
Exposure Sit 3	2.34	2.36	4.38	2.34
H <sub>2</sub> S Fumigation	1.85	2.79	1.35	2.69
NH <sub>3</sub> Fumigation	1.96	2.67	1.27	3.02

Correlating the results obtained in the previous table, we can see the high content of polyphenols in plants exposed to pollutants, especially those exposed in exposure sites and then in fumigation areas compared to the values obtained in climate control tests. This analysis gives us important information about the increase in the amount of polyphenols in plants (in excess) that increases with the degree of pollution. Table 7 shows the degree of inhibition of free radicals by the SOD method.

**Table 2.** Antioxidant activity by determining superoxidismutase. Antioxidant Activity Inhibition %.

	<b>Nicotiana Tabacum</b>	<b>Petunia Hibrida</b>	<b>Ricinus Comunis</b>	<b>Trifolium Pratense</b>
Control	84	93	75	98
Exposure Sit 1	34	27	68	35
Exposure Sit 2	57	42	67	87
Exposure Sit 3	68	54	43	96
H2S Fumigation	75	39	74	25
NH3 Fumigation	69	43	87	31

There has been observed a decrease in the ability to inhibit free radicals in plants exposed to pollutants from non-exposed plants (Control) which is explained as follows: free radicals can occur as a result of all existing forms of pollution (contaminating air, water, soil); to balance the situation, the bodies constantly produce certain substances with antioxidant effect.

In the case of pollutants in the environment, the amount of the enzyme superoxidismutase decreases and the ability of organisms, in our case of exposed plants, to defend against free radicals. In this case, there was a sensitization of plant organisms and the appearance of diseases by destroying cells.

So, according to the results obtained, the increased content of polyphenols was observed in plants exposed to pollutants, especially in those exposed in the exposure sites and then in fumigation glasshouses compared to the values obtained with the control plants in the climatic chamber.

It have been observed the decrease in the ability of plant to inhibit free radicals in plants exposed to pollutants from non-exposed plants (Control) which is explained as follows: free radicals can occur as a result of all existing forms of pollution (contaminating air, water, soil); to balance the situation, the bodies constantly produce certain substances with antioxidant effect.

#### 4. Conclusions

After those determinations we have observed that:

Polyphenols (expressed in gallic acid) are considered as important biomarkers for the phytotoxic effects of pollutants. Thus, the amount of polyphenols in the exposed plants was analyzed compared to the Control plants.

Impregnation of leaf tissues with phenols in higher quantities compared with controls are active responses. This investigation shows how the leaves adapt to increased pollution stress. This analysis gives us important information about the increase in the amount of polyphenols in plants (in excess) that increases with the degree of pollution.

The antioxidant activity (Winterbourn method) of bioindicators was analyzed, knowing the role of antioxidants in the destruction of free radicals in the cell that have a negative effect on living organisms. Superoxidismutase (SOD) is a metal enzyme with subunit structural organization. SOD is the main regulator of oxidation processes in the cell.

The pollutants in the environment, the amount of the enzyme superoxidismutase decreases and the ability of organisms, in our case of exposed plants, to defend against free radicals. In this case, there was a sensitization of plant organisms and the appearance of diseases by destroying cells.

Based on the results of this study, it is concluded that the biochemical and physiological differences can explain the effects of environmental pollution of exposed species.

Antioxidants can be used as a marker of biomonitoring pollution intensity.

**Acknowledgments:** This study was accomplishing through the “Nucleu” Research Project, developed with the Ministry of Research, Innovation and Digitalization of Romania, Agreement no. 20

N/2019, project code PN– 19 04 02 02. The authors would like to thank all those who contributed to this study.

**Conflicts of Interest:** The authors have no conflicts of interest to declare.

## References

1. Gutiérrez-Martínez, P. B.; Torres-Morán, M. I.; Romero-Puertas, M.C.; Casas-Solís, J.; Zarazúa-Villaseñor, P.; Sandoval-Pinto, E.; Ramírez-Hernández B.C. Assessment of antioxidant enzymes in leaves and roots of *Phaseolus vulgaris* plants under cadmium stress, *Biotechnia*, **2020**, Volume 22, no.2, pp. 110-118. <https://doi.org/10.18633/biotechnia.v22i2.1252>
2. Kato, M.; Shimizu, S. Chlorophyll metabolism in higher plants. VII. Chlorophyll degradation in senescing tobacco leaves: phenolic dependent peroxidative degradation. *Can. J. Bot.*, **1987**, Volume 65, pp. 729-735.
3. Liang H.; Zhao S.; Liu K.; Su Y. Roles of reactive oxygen species and antioxidant enzymes on formaldehyde removal from air by plants. *J Environ Sci Health A Tox Hazard Subst Environ Eng.*, **2019**, Volume 54, no. 3, pp. 193-201. <https://doi.org/10.1080/10934529.2018.1544477>.
4. Srivastava, H. S. Biochemical defence mechanisms of plants to increased levels of ozone and other atmospheric pollutants. *Current Science*, **1999**, Volume 76, no. 4, pp. 525–533. <http://www.jstor.org/stable/24100754>
5. Brilli, F.; Fares, S.; Ghirardo, A.; de Visser, P.; Calatayud, V.; Muñoz, A.; Annesi-Maesano, I.; Sebastiani, F.; Alivernini, A.; Variante, V.; and Menghini, F. Plants for Sustainable Improvement of Indoor Air Quality, *Trends Plant Sci.*, **2018**, Volume 23, no. 6, pp 507-512, Elsevier, <https://doi.org/10.1016/j.tplants.2018.03.004>
6. Uka U. N. ; Hogarh J.; Belford E. J. D. , Morpho-anatomical and biochemical responses of plants to air pollution, *International Journal of Modern Botany*, **2017**, Volume 7, no. 1, pp. 1-11, <https://doi.org/10.5923/j.ijmb.20160603.01>
7. Chandawat, D. K.; Verma, P. U.; Solanki, H. A.; Patel, Y. M., Role of Total Phenol in the resistance mechanism of plants against air pollution. *Biolife*, **2014**, Volume 2, no.2, pp. 586-592.
8. Krishnaveni, M. Biochemical changes in plants indicating Air pollution. *Int. J. Pharm. Pharme*, **2013**, Volume 5, pp. 585-586.
9. Blainski, A.; Lopes, G.C.; Palazzo de Mello, J.C. Application and analysis of the Folin Ciocalteu method for the determination of the total phenolic content from *Limonium brasiliense* l. *Molecules*, **2013**, Volume 18, pp.6852-6865; <https://doi.org/10.3390/molecules18066852>
10. Fan, X.; Gu, C.; Cai, J.; Bian Y.; Yang, X.; Sun, C.; Jiang X. Study on active response of superoxide dismutase and relevant binding interaction with bioaccumulated phthalates and key metabolites in *Eisenia fetida*. *Ecotoxicol. Environ. Saf.*, **2021**, Volume 223, no. 15, <https://doi.org/10.1016/j.ecoenv.2021.112559>