



Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPI.



<http://modelos.insmet.cu>




Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)

Email: aluislfh@gmail.com

ECAS 2022 - ID: sciforum-061103

June, 2022



Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPI.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Abstract

The research carries out an evaluation of the 3DVAR method with different options for the assimilation of reflectivity data, which are applied to the SisPI system with the purpose of determining which scheme presents the best results in the short-term numerical weather prediction. For this, data from 6 meteorological radars with coverage over a domain with 3km of spatial resolution are used, using the indirect method with (3DVAR-NoRain) and without (3DVAR) activate an option to also consider null-echoes of reflectivity without presence of precipitation. As a test case, the cold front that affected Cuba on December 10th, 2018 is taken.

Keywords: Data Assimilation; Numerical Weather Prediction; Mesoscale.






Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPI.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
 Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹
 (1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)




Introduction



Users:
 Forecasters,
 Transportation,
 Energy,
 Agriculture,
 Communications,
 Construction, and
 Water Resources
 Management





Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPI.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
 Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹
 (1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



SHORT-TERM NUMERICAL PREDICTION SYSTEM

http://190.115.188.235/haiti_web/sispi.php

INSMET

<http://modelos.insmet.cu>

UNITE HYDRO METEOROLOGIQUE UHM HAITI

República Dominicana

Haiti


ONAMET

<http://sispan.cathalac.int>

CATHALAC

<http://186.149.199.244/sispi.php>

Panama








Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Data Assimilation (DA) of meteorological observations



Insert the information from the meteorological observations in the Numerical Weather Forecast Models and apply consistency restrictions and dynamic balance between all the meteorological variables, to produce a field of analysis that constitutes the initialization of the model.

Radar
Armas (2015)



Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹


(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Previous studies about DA in Cuba

Study	Description
Cruz 2010	Assimilation of stations data in the MM5V3 model.
Hernández 2013, Ferrer 2013	Assimilation of radar and stations data with the ADAS module in ARPS model for high resolution domains.
Sierra et. al. 2014	Assimilation of observations from eolic gradient towers and surface stations with WRFDA module (3DVAR method)
Armas 2015	Assimilation of Radiance and GDAS prepbufr data with the WRFDA using the 3DVAR method.
Ferrer and Borrajero 2019	Assimilation of GDAS Prepbufr and Nexrad Level II radar data with the WRFDA and using the 3DVAR method. The use of CV3, CV5 and CV7 covariance matrices is compared.
Fernández 2019	Performs radar data assimilation experiments with the 3DVAR method but with covariance matrices that employ previous runs from 1 week to 45 days.






Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPI.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Study	Description
González et al. (2021)	It was evaluated for the first time in Cuba different methods such as 3DVAR, 3DENVAR and 4DENVAR, highlighting the advantages and the best results obtained with hybrid schemes, with which the advantages can be appreciated over 3DVAR method, because the contribution of the members of the ensemble provide information about hydrometeors control variables even in areas where radar data is not available.
Aguiar (2021)	The author evaluates different options implemented in the WRFDA for the assimilation of reflectivity data, highlighting the indirect method with the inclusion of hydrometeors. These last works have been limited only to the use of NOAA radar data, hence in this work it was decided to make efforts to use complete coverage throughout the maximum resolution domain of SisPI, with also data from the radar network of the INSMET.



Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹


(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Min & Kim (2016)


Recently available in
WRFDA

radar_non_precip_opt



Lee et al. (2020)

A null-echo is defined as a region with non-precipitation echoes within the radar observation range. The model removes excessive humidity and four types of hydrometeors (wet and dry snow, graupel, and rain) based on the radar reflectivity by using a three-dimensional variational (3D-Var) data assimilation technique within the WRFDA system.



Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPI.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Objectives

- The main objective of this research is to evaluate the impact of assimilating reflectivity data of non-precipitating echoes (radar_non_precip_opt parameter of WRFDA namelist) in the short-term numerical weather forecast of SisPI, which is a recently available option in the WRFDA module, and nowadays its use in the country has not been evaluated.
- On the other hand, it is also intended to carry out an experiment in which a total coverage of reflectivity information can be available in the domain of maximum spatial resolution.



Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.



Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Materials and Methods

For the accomplishment of this work the WRF model (4.1.2) and the WRFDA module (4.3) are used. The assimilation of satellite and prepbuf data from the GDAS system is carried out for all domains, and in the case of domain 3, data from the INSMET and NOAA radar network are also assimilated, guaranteeing with this a total coverage of reflectivity information on the domain of maximum spatial resolution.



A radar data conversion tool in Nexrad Level II format to FM-128 format implemented in Python by Ferrer and Borrajero (2019) is used.



Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPI.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹


(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



WRF-SisPI

Configuration of the work of Sierra et al. (2014, 2017) was taken into account

Forecast Range:
24 Hours



Materials and Methods

WRFDA Configuration (González et al. (2021) and Aguiar (2021))

DA Method: 3DVAR (3 outer loops)
var_scaling = 1.0-0.5-0.25 , len_scaling = 1.0-0.5-0.25

Reflectivity DA Method
Indirect (Wang et al., 2013)


BE Matrix Construction (GEN_BE_v3)
1 month of WRF-SisPI Data (Initialized at 00z and 12z)

BE Matrix (NMC method)
CV7 with Hydrometeors and W

Parameters	Configuration
Spatial Resolution	27, 9, 3 km
Microphysics	WSM5, WSM5, Morrison 2 moment (Morrison et. al., 2009)
Cumulos	Grell-Freitas, Grell-Freitas, No Cumulos
PBL	Mellor-Yamada-Janjic, Mellor-Yamada-Janjic, Mellor-Yamada-Janjic

OBS Types	Domain 01	Domain D02	Domain D03
Airep	8	-	-
Soundings	15	4	-
Metar	325	32	11
Ship	43	21	9
Synop	76	10	3
Buoy	123	11	-
Radiance	1909	438	153
Radar	-	-	2404

Experimental design	Description
NoDA	No data assimilation
3DVAR	GFS + GDAS + Radar
3DVAR_noRain	GFS + GDAS + Radar + Null echoes



Verification Data:

INSMET Weather Stations, GOES-16, Radar Observations, GPM-3IMERGHH product.



Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)






Materials and Methods

As a test case, the cold front that affected the western region of Cuba on December 10th, 2018 is taken, which was preceded by a prefrontal squall line that caused heavy rains on the north coast of Havana with significant accumulations in less than three hours.

Test Case: Initialization 2018-12-10 0000 UTC

GOES-16 CH-14 (IR)





Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.


Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)




Results Discussion


Cost Functions
(3DVAR RADAR DA vs 3DVAR NO-RAIN DA)



3DVAR NO-RAIN DA
Resulting analysis of
reflectivity composite (dBZ)



3DVAR NO-RAIN DA
Increments of reflectivity composite (dBZ)








Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹




(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Results Discussion



Relative Humidity (%) vertical cross-sections





Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPI.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)




Results Discussion

Reflectivity Composite (dBZ)

Observation

3DVAR RADAR DA

3DVAR NO-RAIN DA





Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)












Results Discussion

Cloud Tops Temperature
3DVAR RADAR DA

3DVAR NO-RAIN DA

GOES-16 CH-14 (IR)







Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)






Accumulated precipitation in 24 hours (from 20181210_00z to 20181211_00z)
Radar KBYX (Precipitation estimation)

WRF Experiments (mm/24 hours)

No DA simulation

3DVAR RADAR DA

3DVAR NO-RAIN DA





Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.


Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹
(¹) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)




Results Discussion

Mean Absolute Error (mm/3 hours) of precipitation forecast

Verification with all INSMET stations
(68 synoptic stations)



Verification with Casablanca Station
(WMO 78325)





Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.


Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(¹) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Conclusions

- The Null-echoes assimilation option proved to be an important option in the assimilation of radar data. It allows more realistic construction of moisture fields and its vertical structure in areas of observed reflectivity echoes.
- The use of this option contributes to improving the precipitation forecast and adequately reproduces the convective processes in the first time steps.
- The simulation with radar data assimilation but without the use of this option did not adequately represent the reflectivity echoes and the brightness temperatures of cloud tops in the first 2 hours of simulation.
- The verification with synoptic stations data showed that the use of the radar_non_precip_opt option made it possible to obtain lower Mean Absolute Error values in the forecast of accumulated precipitation in the first 3 hours of simulation.



Impact of the assimilation of non-precipitating echoes reflectivity data on the short-term numerical forecast of SisPl.

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(¹) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)




Recommendations

- Explore the impact of assimilating reflectivity data of non-precipitating echoes (radar_non_precip_opt) using hybrids DA methods (3DEnVAR and 4DEnVAR).
- Insert in the initialization the data of the network of meteorological stations of the INSMET.
- Work on implementing quality control of Cuban radars data.



Center for Atmospheric Physics Institute of Meteorology

QUESTIONS?
Thanks



**Impact of the assimilation of non-precipitating
echoes reflectivity data on the short-term
numerical forecast of SisPI.**

Adrian Luis Ferrer Hernández¹, Pedro Manuel González Jardines¹,
Maibys Sierra Lorenzo¹, Darielis C. Aguiar Figueroa¹

(1) Center for Atmospheric Physics, Institute of Meteorology, Cuba (CFA / INSMET)



Email: aluislfh@gmail.com