

Long-term (2012-2021) Variation in Carbonaceous Aerosols of PM_{2.5} at an urban site of Megacity Delhi situated over Indo-Gangetic Plain of India[†]

Sudhir Kumar Sharma^{1,2,*}, Tuhin Kumar Mandal^{1,2}, Rubiya Banoo^{1,2}, Akansha Rai^{1,2} and Martina Rani^{1,2}

¹ CSIR-National Physical Laboratory, Dr. K S Krishnan Road, New Delhi 110012, India; tuhin.npl@nic.in(TKM); rubiyabanoo31@gmail.com (RB); akansharai.may@gmail.com (AR); martina198919@gmail.com (MR)

² Academy of Scientific and Innovative Research (AcSIR), Ghaziabad 201002, India

* Correspondence: sudhircsir@gmail.com; Sudhir.npl@nic.in

[†] Presented at the 5th International Electronic Conference on Atmospheric Sciences, 16-31 July 2022; Available online: <https://ecas2022.sciforum.net>

Abstract: A long-term (January 2012 – December 2021) study on carbonaceous aerosols of fine particulates (PM_{2.5}) was done over megacity Delhi, India, to evaluate their seasonal and yearly variations. During the entire study period, the observed annual mean levels ($\mu\text{g m}^{-3}$) of PM_{2.5} and its carbonaceous components (OC, POC, SOC, EM, EC, TCM, and TC) were recorded as 126 \pm 72, 15.6 \pm 11.6, 9.3 \pm 6.3, 6.4 \pm 5.1, 8.2 \pm 5.6, 7.3 \pm 5.1, 33.2 \pm 21.9, and 23.1 \pm 16.5, respectively. On average, the CAs/TCM ratio accounts for 26% of PM_{2.5} concentrations. During the monsoon (minimum) and post-monsoon (maximum) season, significant seasonal variability of PM_{2.5} and its carbonaceous species (OC, EC, POC, SOC, and TCM) were observed. Based on the linear association (OC vs. EC) and ratios (OC/EC as well as EC/TC) of species, three significant sources of CAs [vehicular emissions (VE), fossil fuel combustion (FFC), and biomass burning (BB)] were identified.

Keywords: PM_{2.5}; aerosols; carbonaceous species; OC; IGP region

1. Introduction

Carbonaceous aerosols (CAs) are the primary contributors of ambient particulates, accounting for up to 70% of fine aerosol mass [1], and substantially influence atmospheric chemistry, climate, and human health [2]. Majorly, the emission of CAs occurs from fossil fuel combustion, biomass burning, and biogenic emissions [3]. Numerous studies have been carried out in the recent past on CAs of PM_{2.5} and their potential sources in urban [4,5], rural, remote [6] as well as high altitude atmospheres [7] of the Indian region with a yearlong dataset or less, but the limited study is available on a long-term basis. Therefore, to comprehend a more detailed knowledge of fine aerosols (PM_{2.5}) and their atmospheric processes and sources, the long-term research of CAs is crucial in the Indo-Gangetic plain (IGP) region of India. The seasonal and annual changes in PM_{2.5} and its carbonaceous components such as organic carbon (OC), primary organic carbon (POC), secondary organic carbon (SOC), elemental carbon (EC), total carbonaceous matter (TCM), and total carbon (TC) at an urban site of Delhi were reported in this paper, along with their possible sources.

2. Materials and Methods

Periodic sampling (2 samples a week) of PM_{2.5} was done from January 2012 to December 2021 at the CSIR-National Physical Laboratory (28° 38' N, 77° 10' E; 218 m above mean sea level). Delhi, the capital of India, is surrounded by different climatic zones and

Academic Editor: Anthony Lupu

Published: 31 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

is considered one of the most polluted megacities in the world [8]. The meteorology of Delhi experienced four distinct seasons, which are winter (January-February; temperature: ~3°C), summer (March-May; temperature: 47°C), monsoon (June-September; rain: 800mm), and post-monsoon (October-December). A detailed description of the sampling location is available in our earlier publication [9].

PM_{2.5} samples were collected on quartz filters (size: 47 mm) using a fine particle sampler (flow rate: 1 m³ h⁻¹) for 24 hours. A Thermo-optical carbon analyzer (Model: DRI 2001A) was used to measure OC and EC following the IMPROVE-A protocol [10]. Detailed operating procedures of the instrument are available in Sharma *et al.* [9]. The TCM concentration can be computed by the addition of OM (OM = 1.6 × OC; 1.6 factor for PM_{2.5}) and EM (EM = 1.1 × EC) of PM_{2.5} [11, 12]. The EC tracer approach [13] was used for determining the SOC (Eq. 1):

$$\text{SOC} = \text{OC} - \text{POC} ([\text{OC}/\text{EC}]_{\text{min}} \times \text{EC}) \tag{1}$$

3. Results and Discussion

Figure 1 illustrates the annual changes in PM_{2.5} and their carbonaceous components (OC, EC, and TC) during the study period (January 2012 to December 2021). In this long-term study, the annual mean (± standard deviation) concentrations (µg m⁻³) of PM_{2.5} and their carbonaceous components (OC, POC, SOC, EM, EC, TCM, and TC) were recorded as 126±72, 15.6±11.6, 9.3±6.3, 6.4±5.1, 8.2±5.6, 7.3±5.1, 33.2±21.9, and 23.1±16.5, respectively. Among the species, the contribution (w.r.t. annual mean level) to PM_{2.5} was observed highest by TCM (~26%), followed by OC (~12%) and EC (~6%). Similar observations were also stated by Jain *et al.* [5], with ~25% of TCM, ~12% of OC, and ~5.5% of EC to PM_{2.5} over Delhi.




Figure 1 Annual trend in PM_{2.5}, EC, OC, and TC levels over Delhi.

Figure 2 illustrates the seasonal variability of PM_{2.5} and its carbonaceous species (OC, EC, TC, and TCM). PM_{2.5} and their CAs (OC, EC, TCM, POC, and SOC) were recorded as minimum and maximum during the monsoon and post-monsoon season, respectively (Table 1). This might be owing to the source intensity of PM_{2.5}, prevailing meteorological circumstances, and the long-distant movement of pollutants from Punjab and Haryana to the receptor location [8].

The mass ratio of OC/EC has been used to identify the possible sources of CAs in most studies [14, 15]. The higher value of the OC/EC ratio that ranges from 4 to 12 indicates the dominance of biomass burning (BB) [15]. On the other hand, when the OC/EC ratio ranged between 1.4 to 4, it indicates the occurrence of VE together with BB [16]. During the study period, the observed average ratio of OC/EC of PM_{2.5} were 2.3±0.7, 2.1±0.6, 2.2±0.8, and 2.5±0.8 during winter, summer, monsoon, and post-monsoon seasons, respectively (Table 1). From the observed OC/EC mass ratio, BB and FFC are the major sources of CAs over study site in Delhi. Jain et al. [5] have also reported that BB, FFC, and VE are the main sources of PM_{2.5} in megacity Delhi. Sharma et al. [17] investigated the stable carbon and nitrogen isotope of aerosols at a Delhi metropolitan location and revealed that the VE, BB, and FFC are the most likely main sources of aerosols in the city. These sources are also influenced by regional and long-distant transport of pollutants in Delhi.

Table 1. Seasonal mean of PM_{2.5} and their carbonaceous species (µg m⁻³) in Delhi.

Species	Seasons			
	Winter	Summer	Monsoon	Post-Monsoon
PM _{2.5}	157±64	94±35	66±28	189±92
OC	20.6±10.9	9.4±4.3	6.7±2.8	25.8±14.2
EC	9.3±5.2	4.9±2.7	2.8±1.5	10.8±6.5
TC	29.7±15.5	14.4±6.5	9.5±4.3	36.5±19.4
POC	11.8±5.9	6.2±3.1	3.4±1.7	13.2±7.1
SOC	8.3±6.6	3.5±1.8	3.0±1.6	12.3±8.6
OC/EC	2.3±0.7	2.1±0.6	2.2±0.8	2.5±0.8




Figure 2 Seasonal mean of (error bar: ± SD) OC, EC, TC, and TCM of PM_{2.5} in Delhi.

4. Conclusions

In the present study of 10-year long-term dataset, the annual and seasonal mean concentrations of PM_{2.5} and all carbonaceous species vary significantly in megacity Delhi. In total, about 26% of CAs accounts to PM_{2.5} mass concentration. A substantial variability in seasonal concentrations of PM_{2.5} and its associated CAs were observed as follows: post-

monsoon > winter > summer > monsoon seasons. A linear relationship was observed between OC and EC which demonstrates that pollutants generated from similar combustion sources in Delhi, India. In addition, BB, VE and FFC are the prominent sources of CAs of PM_{2.5} suggested by the OC/EC as well as EC/TC ratios. Thus, the observed results states the significant effect of CAs from different sources.

Author Contributions: Conceptualization and the first draft of the manuscript is prepared by SKS. PM_{2.5} samples collection and chemical analysis were performed by SKS, RB, AR, MR and TKM. All the authors read and reviewed the manuscript before communication

Funding: The authors also acknowledge the CSIR-NPL, New Delhi for financial support for this study.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The datasets are available with corresponding author and will be provided on reasonable request.

Acknowledgments: The authors are thankfully acknowledging to the Director, CSIR-NPL, New Delhi and Head, ES&BMD, CSIR-NPL, New Delhi for their support of this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lelieveld, J.; Evans, J.S.; Fnais, M.; Giannadaki, D.; Pozzer, A. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* **2015**, *525*, 367–371.
2. Dockery, D.W.; Stone, P.H. Cardiovascular risks from fine particulate air pollution. *N Engl J Med* **2007**, *356*, 511–513.
3. Venkataraman, C.; Habib, G.; Eiguren-Fernandez, A.; Miguel, A. H.; Friedlander, S. K. Residential biofuels in south Asia: carbonaceous aerosol emissions and climate impacts. *Science* **2005**, *307*, 1454–1456.
4. Sharma, S.K.; Mandal, T.K.; Jain, S.; Saraswati.; Sharma, A.; Saxena, M. Source apportionment of PM_{2.5} in Delhi, India using PMF model. *Bull Environ Contam Toxicol* **2016**, *97*, 286–293.
5. Jain, S.; Sharma, S. K.; Vijayan, N.; Mandal, T.K. Seasonal characteristics of aerosols (PM_{2.5} and PM₁₀) and their source apportionment using PMF: A four year study over Delhi, India. *Environ Poll* **2020**, *262*, 114337.
6. Begam, G. R.; Vachaspati, C.V.; Ahammed, Y.N.; Kumar, K.R.; Reddy, R.R.; Sharma, S.K.; Saxena, M.; Mandal, T.K. Seasonal characteristics of water soluble inorganic ions and carbonaceous aerosols in total suspended particulate matter at a rural semi-arid site, Kadapa (India). *Environ Sci Poll Res* **2017**, *24*(2), 1719–1734.
7. Sharma, S.K.; Mukherjee, S.; Choudhary, N.; Rai, A.; Ghosh, A.; Chatterjee, A.; Vijayan, N.; Mandal, T.K. Seasonal variation and sources of carbonaceous species and elements of PM_{2.5} and PM₁₀ over the eastern Himalaya. *Environ Sci Poll Res* **2021a**, *28*, 51642–51656.
8. Sharma, S.K.; Banoo, R.; Mandal, T.K. Seasonal characteristics and sources of carbonaceous components and elements of PM₁₀ (2010–2019) in Delhi, India. *J Atmos Chem* **2021b** *78*, 251–270.
9. Sharma, S.K.; Mandal, T.K.; Saxena, M.; Rashmi.; Rohtash.; Sharma, A.; Gautam, R. Variation of OC, EC, WSIC and trace metals of PM₁₀ in Delhi. *J Atmos Sol Terr Phy* **2014**, *113*, 10–22.
10. Chow, J. C.; Watson, J.G.; Chen, L. W. A.; Arnott, W. P.; Moosmuller, H. Equivalence of elemental carbon by thermal/optical reflectance and transmittance with different temperature protocols. *Environ Sci Technol* **2004**, *38*, 4414–4422.
11. Amato, F.; Pandolfi, M.; Escrig, A.; Querol, X.; Alastuery, A.; Pey, J.; Perez, N.; Hopke, P.K. Quantification of road dust resuspension in urban environment by multilinear engine: a comparison with PMF2. *Atmos Environ* **2009**, *43*, 2770–2780.

12. Malm, W.C.; Schichtel, B.A.; Pitchford, M.L.; Ashbaugh, L.L.; Eldred, R.A. Spatial and monthly trends in speciated fine particle concentration in the United States. *J Geophys Res:Atmos* **2004**, *16*,109(D3).
13. Castro, L.M.; Pio, C.A.; Harrison, R.M.; Smith, D.J.T. Carbonaceous aerosol in urban and rural European atmospheres: estimation of secondary organic carbon concentrations. *Atmos Environ* **1999**, *33*, 2771–2781.
14. Ram, K.; Sarin, M.M.; Tripathi, S.N. One-year record of carbonaceous aerosols from an urban location (Kanpur) in the Indo-Gangetic Plain: characterization, sources and temporal variability. *J Geophys Res* **2010**, doi:10.1029/2010JD014188
15. Szidat, S.; Jenk, T.M.; Synal, H.A.; Kalberer, M.; Wacker, L.; Hajdas, Kasper-Gieblt. A.; Baltensperger, U. Contribution of fossil fuel, biomass burning and biogenic emissions to carbonaceous aerosols in Zurich as traced by ¹⁴C. *J Geophys Res* **2006**, *111*:D07206.
16. Salameh, D.; Detournay, A.; Pey, J.; Pérez, N.; Liguori, F.; Saraga, D.; Bove, M.C.; Brotto, P.; Cassola, F.; Massabò, D.; Latella, A. PM_{2.5} chemical composition in five European Mediterranean cities: a 1-year study. *Atmos Res* **2015**, *155*, 102-117.
17. Sharma, S.K.; Karapurkar, S.G.; Shenoy, D.M.; Mandal, T.K. Stable carbon and nitrogen isotopic characteristics of PM_{2.5} and PM₁₀ in Delhi, India. *J Atmos Chem* **2022**, doi.org/10.1007/s10874-022-09429-0.