

Proceeding Paper

# Chemical and Nutritional Characterization of Various by-Products of the Industry *Olea europea* L. Source of Healthy Ingredients <sup>†</sup>

F. Chamorro <sup>1</sup>, Lucia Cassani <sup>1,2</sup>, Pauline Donn <sup>1</sup>, Sepidar Seyyedi Mansour <sup>1</sup>, M. Fraga-Corral <sup>1,2</sup>, Jianbo Xiao <sup>1</sup>, Jesus Simal-Gandara <sup>1</sup>, M.A. Prieto <sup>1,2,\*</sup> and Paz Otero <sup>1,2,\*</sup>

<sup>1</sup> Nutrition and Bromatology Group, Department of Analytical and Food Chemistry, Faculty of Food Science and Technology, University of Vigo, Ourense Campus, E32004 Ourense, Spain

<sup>2</sup> Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolonia, 5300-253 Bragança, Portugal

\* Correspondence: author: mprieto@uvigo.es (M.A.P.); paz.otero@uvigo.es (P.O.)

† Presented at the 3rd International Electronic Conference on Foods: Food, Microbiome, and Health—A Celebration of the 10th Anniversary of Foods' Impact on Our Wellbeing; Available online: <https://foods2022.sciforum.net>.

**Abstract:** Currently, the food industry is facing a wide demand for natural foods that provide benefits for the consumer's health. In this sense, a trend in food technology is to obtain bioactive compounds from different food by-products with antioxidant, anti-inflammatory, antiviral properties, among others, that allow their recovery through the circular economy model. In this study, the chemical and nutritional composition was evaluated in terms of phenolic compounds (PC), fatty acid profile (FA) and mineral composition, in several by-products of *Olea europaea* L.: pomace, olive water, olive leaves, fiber crude, and an extract obtained by extraction with supercritical fluid (SCFE) at 20 and 24 MPa of the olive fiber. PC quantification was performed by liquid chromatography-mass spectrometry (LC-MS/MS), the fatty acid profile was obtained by gas chromatography coupled to a flame ionization detector (GC-FID) and the mineral content was determined by inductively coupled plasma optical emission spectrometry (ICP-OES). In all the byproducts studied, dihydroxybenzoic acid, hydroxytyrosol and oleacein were found as main PCs. Olive fiber presented the highest content of hydroxytyrosol (171.2 mg/kg) and oleacein (150 mg/kg). Regarding the content of FA, a high concentration of monounsaturated fatty acids (MUFA) between 71–73% was reported in all the by-products of the olive, highlighting oleic acid as the main FA. The fraction of polyunsaturated fatty acids (PUFA) corresponded to 10–12%, with  $\alpha$ -linolenic acid being the main compound. With respect to the mineral profile, some differences were observed, the pomace and the fiber presented relevant values of calcium (4–6.5 g/kg), potassium (0.5 g/kg) and magnesium (0.5 g/kg). while the oily extract did not report significant levels of minerals. Taking into account the previous results, the by-products resulting from the olive industry can be considered as a valuable source of bioactive molecules, including PC, AG and minerals, which provide important beneficial effects for health, mainly for the prevention of diseases related to oxidative stress, and that could be used as novel natural ingredients for incorporation into functional foods or nutraceuticals, however, both in vitro and in vivo studies are needed to confirm the potential in humans and demonstrate the safety of these ingredients.

**Keywords:** *Olea europaea* L; by-products; bioactive compounds profile; LC-MS; GC-FID; ICP-OES

**Citation:** Chamorro, F.; Cassani, L.; Donn, P.; Mansour, S.S.; Fraga-Corral, M.; Xiao, J.; Simal-Gandara, J.; Prieto, M.A.; Otero, P. Chemical and Nutritional Characterization of Various by-Products of the Industry *Olea europea* L. Source of Healthy Ingredients. *Biol. Life Sci. Forum* **2022**, *2*, x. <https://doi.org/10.3390/xxxxx> Published: 1 October 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Acknowledgments:** The research leading to these results was supported by MICINN supporting the Ramón y Cajal grant for M.A. Prieto (RYC-2017-22891) and Jianbo Xiao (RYC-2020-030365-I), by Xunta de Galicia for supporting the program EXCELENCIA-ED431F 2020/12, the post-doctoral grant of M. Fraga-Corral (ED481B-2019/096), and L. Cassani (ED481B-2021/152). The authors thank

the program BENEFICIOS DO CONSUMO DAS ESPECIES TINTORERA-(CO-0019-2021) that supports the work of F. Chamorro. The research leading to these results was supported by the European Union through the “NextGenerationEU. Authors are grateful to Ibero-American Program on Science and Technology (CYTED—AQUA-CIBUS, P317RT0003), to the Bio Based Industries Joint Undertaking (JU) under grant agreement No 888003 UP4HEALTH Project (H2020-BBI-JTI-2019) that supports the work of P. Otero. The JU receives support from the European Union’s Horizon 2020 research and innovation program and the Bio Based Industries Consortium. The project SYSTEMIC Knowledge hub on Nutrition and Food Security, has received funding from national research funding parties in Belgium (FWO), France (INRA), Germany (BLE), Italy (MIPAAF), Latvia (IZM), Norway (RCN), Portugal (FCT), and Spain (AEI) in a joint action of JPI HDHL, JPI-OCEANS and FACCE-JPI launched in 2019 under the ERA-NET ERA-HDHL (n° 696295).