

Effect of lemon balm and spearmint extracts on the survival of *S. aureus* in goat's raw milk cheese [†]

Beatriz Nunes Silva ^{1,2,3,4}, Sara Coelho-Fernandes ^{1,2}, José António Teixeira ^{3,4}, Vasco Cadavez ^{1,2} and Ursula Gonzales-Barron ^{1,2*}

¹ Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal; vcadavez@ipb.pt (V.C.); sara.coelho@ipb.pt (S.C.-F.)

² Laboratório para a Sustentabilidade e Tecnologia em Regiões de Montanha, Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

³ CEB – Centre of Biological Engineering, University of Minho, Campus Gualtar, 4710-057 Braga, Portugal; beatriznunesilva@ceb.uminho.pt (B.N.S.); jateixeira@deb.uminho.pt (J.A.T.)

⁴ LBBELS – Associate Laboratory, Braga 4710-057 Guimarães, Portugal

* Correspondence: ubarron@ipb.pt; Tel.: +35-12-7330-3325

[†] Presented at the 3rd International Electronic Conference on Foods: Food, Microbiome, and Health, 01-15 October 2022; Available online: <https://sciforum.net/event/Foods2022>

Abstract: Previous investigation from our research group has revealed that lemon balm and spearmint hydroethanolic extracts present high inhibitory capacity in-vitro against *S. aureus*. Raw milk cheeses have shown moderate prevalence of *S. aureus*, thus imposing a safety issue for consumers. In this sense, our work aimed to evaluate the antibacterial effect of lemon balm and spearmint extracts against *S. aureus* in goat's raw milk cheeses during maturation and characterise the survival kinetic parameters of this pathogen by extended Bigelow models.

Keywords: Log-decay model with tail; Bigelow model; D-value; Z_{pH} ; dairy; predictive microbiology

1. Introduction

A meta-analysis study by Gonzales-Barron et al. [1] revealed *S. aureus* as the pathogen of highest incidence in goat raw milk (35.2%; 95% CI: 23.2–49.3%), and the overall occurrence of *S. aureus* in goat milk cheeses to be noticeably high (16.0%; CI: 7.92–29.8%). Moreover, cheeses made of raw milk, regardless of origin, presented an even higher prevalence of *S. aureus* (38.7%; 95% CI: 9.28–79.6%). These concerning prevalence values underscore the importance of improving cheese manufacture to control *S. aureus* development. For this, biopreservative agents such as plant extracts can be used [2].

Previous investigation from our research group has demonstrated the antimicrobial capacity of several plant extracts against *S. aureus in vitro* by determination of their minimum inhibitory concentration (MIC) [3]. Lemon balm extract resulting from hydroethanolic (70% v/v) solid-liquid extraction presented a MIC of 2.5 mg/mL against *S. aureus*, whereas the equivalent extract obtained from spearmint showed a MIC of 1.25 mg/mL against this pathogen [3]. Such outcomes suggest the potential of lemon balm and spearmint extracts to be incorporated in foods as preservatives against microbial spoilage.

In this sense, the objective of this work was to assess the antimicrobial effect against *S. aureus* of lemon balm and spearmint extracts in goat's raw milk cheeses, when directly incorporated in curd, and to characterise the survival kinetic parameters of this pathogen by means of an extended Bigelow model. With this approach, values of decimal reduction time (*D*) can be described as a function of pH and incorporation of plant extract, and the inactivation parameters of *S. aureus* may help on the optimisation of the manufacturing process to ensure microbial safety of cheeses.

Citation: Silva, B.N.;

Coelho-Fernandes, S.; José António

Teixeira; Cadavez, V.;

Gonzales-Barron, U. Effect of lemon

balm and spearmint extracts on the

survival of *S. aureus* in goat's raw

milk cheese. 2022, volume number, x.

<https://doi.org/10.3390/xxxxx>

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

2. Materials and Methods

2.1. Plant Material and Extraction Procedure

Dried lemon balm and spearmint aerial parts were provided by Pragmático Aroma Lda. (“Mais Ervas”, Trás-os-Montes, Portugal) and mechanically ground. Extracts were obtained using ethanol 70% (*v/v*) as solvent in a shaking water bath (150 rpm) at 60 °C for 90 minutes. The sample/solvent ratio was 1:20. The mixtures were filtrated (7–10 µm), and the ethanolic fraction was evaporated. The remaining aqueous fraction was frozen and lyophilised.

2.2. Inoculation of *S. aureus* in milk and cheese production

Staphylococcus aureus ATCC 6538 kept on a fresh slant was cultivated twice at 37 °C, 200 rpm, for 16h, first on tryptic soy broth (TSB) and then on tryptic soy broth with pH adjusted to 6.34, to mimic goat’s milk pH. On the day of inoculation, the second subculture was centrifuged at 10640g at 4 °C for 10 minutes, for removing debris and residual culture media. After centrifugation, the supernatant was discarded, and pellets were washed with sterile 0.9% physiological solution. Centrifugation and washing procedures were repeated twice and cells were re-suspended in sterile 0.9% physiological solution to reach ~7 log CFU/mL.

Lab-scale cheeses were prepared by adding the rennet (0.75 mL/L milk) and inoculum (5 mL/L milk) to milk at ~34 °C. Through this procedure, each cheese reached a *S. aureus* target concentration of 4 to 5 log CFU/g, depending on the milk initial contamination. After 30 minutes at 34 °C, curdled milk was cut and drained, and 1% (*w/w*) of lyophilised spearmint or lemon balm extract was added to the curd and mixed, while an inoculated control without extract was kept. The curd was then placed in 50 mL tubes and centrifuged at 6000 rpm at 20 °C for 3.5 minutes. The supernatant (whey) was removed, and cheeses of ~5 g were cut from the compacted curd and placed in a 15% brine solution for 10 minutes for salting. Finally, the weight in g of each cheese was annotated and cheeses were kept in a climate-controlled chamber (10 °C, 98% RH) for fermentation and maturation to take place for 15 days.

2.3. Microbiological and Physicochemical Analyses

Analyses were conducted on days 0 (day of cheese production), 2, 4, 7, 10, 13 and 15. For the microbiological determinations, for every test unit, appropriate serial dilutions were prepared by homogenising the cheese in 50 mL of buffered peptone water for 60 seconds. To determine the concentration of *S. aureus*, 0.1 mL aliquot was plated on Baird-Parker agar, supplemented with Egg Yolk Tellurite, following ISO norm [4]. Typical colonies were counted after 48 h following incubation at 37 °C.

Physicochemical analyses comprised the measurement of pH, which was carried out using a pH meter (Hanna Instruments, model HI5522, USA) equipped with a HII1131 glass penetration probe.

2.4. Modelling

For every treatment, a log-decay function with tail in differential form as primary model, with varying *D*-value, coupled to a secondary model Bigelow equation of *D*-value as a function of pH (with parameters log *D*_{ref} at pH 7.0 and *z*_{pH}) was adjusted, as follows:

$$\frac{dN}{dt} = -kN \left(\frac{1}{1 + C_c} \right) \left(1 - \frac{N_{res}}{N} \right) \quad (1)$$

$$\frac{dC_c}{dt} = -kC_c \tag{2}$$

$$D = \frac{\ln(10)}{k} \tag{3}$$

$$\log D = \log D_{ref} - \left(\frac{pH - pH_{ref}}{z_{pH}} \right)^2 \tag{4}$$

Herein, N is the population density (CFU/g), k is the inactivation rate (1/day), C_c is the physiological state of the cells, N_{res} is the residual population density (CFU/g), D (days) is the decimal reduction time at the constant temperature T (10 °C) and at the pH of the cheese, pH_{ref} is the reference pH (set to 7), z_{pH} is the distance of pH from pH_{ref} which leads to a ten-fold change in decimal reduction time, and D_{ref} is the decimal reduction time at pH_{ref} (days).

3. Results and Discussions

Bigelow-type secondary models were used to describe the inactivation of *S. aureus* in goat’s raw milk cheeses during maturation as affected by spearmint and lemon balm extracts. The survival curves of *S. aureus* in cheese without and with plant extracts, as depicted by dynamic models, are presented in Figure 1 (spearmint) and Figure 2 (lemon balm). The results of the Bigelow parameters for each treatment are shown in Table 1.




Figure 1. Survival of *S. aureus* in goat’s raw milk cheese without (left) and with spearmint extract (right), as depicted by dynamic models with 95% CI.




Figure 2. Survival of *S. aureus* in goat’s raw milk cheese without (left) and with lemon balm extract (right), as depicted by dynamic models with 95% CI.

Table 1. Effect of adding 1% spearmint extract and 1% lemon balm extract in curd on the Bigelow’s inactivation parameters of *Staphylococcus aureus* in goat’s raw milk cheese during maturation.

Treatment	Bigelow parameters	Mean (SE)	Pr (> t)	Goodness-of-fit measures
Spearmint 0% ($C_c(0) = 1.5$)	$\log D_{ref}$	0.932 (0.166)	<.0001	$S^2 = 0.0017$
	z_{pH}	1.727 (0.392)	0.001	RMSE = 0.0403 MAE = 0.0357
Spearmint 1% ($C_c(0) = 0.1$)	$\log D_{ref}$	0.621 (0.061)	<.0001	$S^2 = 0.0147$
	z_{pH}	3.172 (0.660)	<.0001	RMSE = 0.1172 MAE = 0.0978
Lemon balm 0% ($C_c(0) = 1.5$)	$\log D_{ref}$	0.996 (0.056)	<.0001	$S^2 = 0.0015$
	z_{pH}	1.851 (0.007)	<.0001	RMSE = 0.0374 MAE = 0.0330
Lemon balm 1% ($C_c(0) = 0.1$)	$\log D_{ref}$	1.189 (0.200)	<.0001	$S^2 = 0.0042$
	z_{pH}	2.339 (0.835)	0.019	RMSE = 0.0633 MAE = 0.0556

S^2 : variance; RMSE: root mean square error; MAE: mean absolute error.

The dynamic models adequately fitted the survival curves with root mean square errors (RMSE) of 0.1172 and 0.0633 for spearmint and lemon balm, respectively (Table 1), producing significant parameter estimates ($p < 0.05$).

From Table 1, the parameter $\log D_{ref}$ was affected by the addition of extracts (0.621 [SE = 0.061] for spearmint; 1.189 [SE = 0.200] for lemon balm) in comparison to the controls (0.932 [SE = 0.166] and 0.996 [SE = 0.056]).

In the case of cheeses with spearmint extract, $\log D_{ref}$ was lower than that of the control, implying a greater inactivation rate of the pathogen. The survival curves presented in Figure 1 show that the incorporation of this extract reduced the initial lag phase (the “shoulder”) and promoted *S. aureus* inactivation earlier in maturation.

Oppositely, when adding lemon balm extract to the cheese, the estimated $\log D_{ref}$ was higher than that of the control, thus suggesting a slower inactivation rate. However, *S. aureus* inactivation was steadier and more prolonged throughout maturation, compared to control cheeses, in which *S. aureus* inactivation phase was rather short and stationary phase (“tail”) was reached sooner (Figure 2).

In both cases, the addition of plant extracts significantly decreased the time to achieve one log reduction, which in practical terms corresponded to up to 1.36 log CFU/g reduction by the end of maturation. Such outcomes support the usefulness of incorporating spearmint and lemon balm extracts to reduce *S. aureus* burden in this dairy product.

The fermentation process was affected by the presence of extracts in the sense that the pH drop in the beginning of maturation (until day 4) was slower, and the pH value reached after 14 days was slightly higher, compared to the corresponding control group (data not shown). This is reflected by the higher z_{pH} values of cheeses with spearmint and lemon balm extracts in Table 1 (3.172 [SE = 0.660] and 2.339 [SE = 0.835, respectively) and means that a greater difference between pH and pH_{ref} is necessary to lead to a tenfold change in D when incorporating plant extracts in cheese, than the one needed for the same variation in D in the controls. This outcome is likely a result of plant extracts affecting, to some extent, the production of organic acids by bacteriocinogenic lactic acid bacteria that drop the pH during fermentation. Nonetheless, after day 4, the pH trend for both treatments and controls were similar.

The results of this work indicate that the main effect of adding 1% lemon balm extract in curd is on *S. aureus* lag phase and the z_{pH} ; while 1% spearmint extract affects lag phase, z_{pH} and $\log D_{ref}$. Therefore, spearmint extract is more efficient in controlling *S. aureus* in goat's raw milk cheese.

4. Conclusions

Using Bigelow-type secondary models, this work characterised *S. aureus* survival parameters in goat's raw milk cheese. The results indicate that both parameters, $\log D_{ref}$ and z_{pH} , were affected by the addition of extracts. z_{pH} values are increased by the addition of extracts due to their interaction with the ongoing fermentation.

The dynamic models also demonstrated that the addition of lemon balm and spearmint extracts reduced the time needed to achieve one log reduction of *S. aureus*, thus showing their ability to act as biopreservatives against this pathogen during cheese maturation.

Author Contributions: Conceptualization, V.C., and U.G.-B.; Data curation, B.N.S. and U.G.-B.; Formal analysis, B.N.S., V.C. and U.G.-B.; Funding acquisition, V.C., J.A.T. and U.G.-B.; Investigation, B.N.S. and S.C.-F.; Methodology, B.N.S., S.C.-F., V.C. and U.G.-B.; Project administration, V.C. and U.G.-B.; Resources, V.C. and U.G.-B.; Software, B.N.S., and U.G.-B. Supervision, V.C., J.A.T. and U.G.-B.; Validation, U.G.-B.; Visualization, B.N.S., V.C. and U.G.-B.; Writing—original draft, B.N.S.; Writing—review & editing, B.N.S. and U.G.-B. All authors have read and agreed to the published version of the manuscript.

Funding: The authors are grateful to the EU PRIMA program and the Portuguese Foundation for Science and Technology (FCT) for funding the ArtiSaneFood project (PRIMA/0001/2018). The authors are grateful to the Foundation for Science and Technology (FCT, Portugal) for financial support through national funds FCT/MCTES (PIDDAC) to CIMO (UIDB/00690/2020 and UIDP/00690/2020) and SusTEC (LA/P/0007/2021). This study was supported by FCT under the scope of the strategic funding of UIDB/04469/2020 unit and Bio-TecNorte operation (NORTE-01-0145-FEDER-000004) funded by the European Regional Development Fund under the scope of Norte2020—Programa Operacional Regional do Norte.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Summary data available upon request.

Acknowledgments: BNS acknowledges the financial support provided by the Foundation for Science and Technology (FCT, Portugal) through the PhD grant SFRH/BD/137801/2018. The authors are grateful to EU PRIMA programme and FCT for funding the ArtiSaneFood project (PRIMA/0001/2018), and to FCT for financial support through national funds FCT/MCTES (PIDDAC) to CIMO (UIDB/00690/2020 and UIDP/00690/2020) and SusTEC (LA/P/0007/2021). Dr. Gonzales-Barron acknowledges the national funding by FCT, P.I., through the Institutional Scientific Employment Programme contract.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Gonzales-Barron, U.; Gonçalves-Tenório, A.; Rodrigues, V.; Cadavez, V. Foodborne pathogens in raw milk and cheese of sheep and goat origin: a meta-analysis approach. *Curr. Opin. Food Sci.* **2017**, *18*, 7-13. 1
2. Bondi, M.; Lauková, A.; de Niederhausern, S.; Messi, P.; Papadopoulou, C. Natural Preservatives to Improve Food Quality and Safety. *J. Food Qual.* **2017**, 2017, 1–3. 2
3. Silva, B.N.; Cadavez, V.; Ferreira-Santos, P.; Alves, M.J.; Ferreira, I.C.F.R.; Barros, L.; Teixeira, J.A.; Gonzales-Barron, U. Chemical Profile and Bioactivities of Extracts from Edible Plants Readily Available in Portugal. *Foods* **2021**, *10*, 673. 3
4. ISO 6888-1:2021. *Microbiology of the Food Chain - Horizontal Method for the Enumeration of Coagulase-Positive Staphylococci (Staphylococcus aureus and Other Species) - Part 1: Method Using Baird-Parker Agar Medium*. International Organization for Standardization: Geneva, Switzerland, 2021. 4