

Proceeding Paper

Indoor Air Quality (PM_{2.5} and PM₁₀) and Toxicity Potential at a Commercial Environment in Akure, Nigeria[†]

Francis Olawale Abulude^{1,*}, Kikelomo Mabinuola Arifalo², Ademola Adamu², Amoke Monisola Kenni³, Akinyinka Akinnusotu⁴, Samuel Dare Oluwagbayide⁵, Sunday Acha⁶

Citation: Abulude, F.O., Arifalo, K.M., Adamu, A., Kenni, A.M., Akinnusotu, A., Oluwagbayide, S.D., Acha, S. Indoor Air Quality (PM_{2.5} and PM₁₀) and Toxicity Potential at a Commercial Environment in Akure, Nigeria. *Environ. Sci. Proc.* **2022**, *4*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Jon Ø. Odland

Published: 26 October 2023

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

¹ Science and Education Development Institute, Akure, Ondo State, Nigeria

² Department of Chemistry, Bamidele Olumilua University of Education, Science and Technology, Ikere - Ekiti, Ekiti State, Nigeria

³ Department of Science Education, Bamidele Olumilua University of Education, Science and Technology, Ikere - Ekiti, Ekiti State, Nigeria

⁴ Department Science Laboratory Technology, Rufus Giwa Polytechnic, P.M.B. 1019, Owo, Ondo State, Nigeria

⁵ Department of Agricultural and Bio-environmental Engineering, Federal Polytechnic, P.M.B. 50, Ilaro, Ogun State, Nigeria

⁶ Department of Geography, Federal University of Technology, Minna, Niger State, Nigeria

* Correspondence: walefut@gmail.com; Tel.: (+2348034458674)

† Presented at the 4th International Electronic Conference on Environmental Research and Public Health — Climate Change and Health in a Broad Perspective 15–30 Oct 2022

Abstract: According to studies, indoor air quality is a major concern because of the health risks it poses. In Nigeria, little is done to improve indoor air quality and the toxicity potentials (TP) of PM_{2.5} and PM₁₀. We assessed the levels of PM_{2.5}, PM₁₀, the PM_{2.5}/PM₁₀ ratio, and the toxicity potential of a commercial area in Akure, Ondo State, Nigeria, in this study. For the three-month assessment of the study area (March to May 2022), a low-cost sensor (Canãree A1) was used. The results depict the following: 73.23±53.94 µg/m³ (PM_{2.5}), 68.58±50.64 µg/m³ (PM₁₀), 0.93±0.02 (PM_{2.5}/PM₁₀ ratio), and toxicity potentials (PM_{2.5} - 2.74±0.04 and PM₁₀ - 1.47±0.02). Both PM values exceed the WHO standard limits. The PM values differ significantly. The average ratio value indicates that anthropogenic activities in the area contribute significantly to the high PM_{2.5} levels. It should be noted that TP greater than 1 indicates a potential health risk. The TP values obtained in this study are greater than one, indicating that the environment may be harmful to the vulnerable. Based on these findings, efforts should be directed toward continuous monitoring of this study area and Akure as a whole.

Keywords: Indoor; health risks; low-cost sensor, WHO standard, Nigeria

1. Introduction

In general, neither developing nor developed countries take the effects of air pollution for granted. Attempts are being made to reduce or entirely eliminate it. Pollutant gases (O₃, NO₂, SO₂, and CO) and particulate matter (PM - PM₁₀ and PM_{2.5}) are the main culprits of air pollution. The size of particles in PM is directly related to their likelihood of triggering health problems [1] Small particles less than 10 micrometers in diameter cause the most issues because they can penetrate deep into lungs and, in some cases, enter into bloodstream. Exposure to such particles can harm the lungs as well as the heart. Various studies have shown particle pollution exposure to a variety of problems, including: premature death in people with heart or lung disease, nonfatal heart attacks, irregular heartbeat, aggravated asthma, decreased lung function, and increased respiratory symptoms such as airway irritation, coughing, or difficulty breathing. Particle pollution exposure is most likely to affect vulnerable people (the sick, children, and the elderly).

Researchers have discovered that human activity is a major source of heavy metals and fine particulate matter contaminants in the air [2-4]. Industrial activities in both developed and developing nations are documented to contribute significantly to human induced air pollution, with particulate matter and gaseous pollutants being produced at mostly undesirable levels. Heavy metal air pollution is a worldwide issue because most metals are inextinguishable and can endanger human health, plants, animals, ecosystems, or other media [4]. The studies on pollutant toxicity potential have been motivated by growing human health concerns about PM inhalation [3]. Particulate matter toxicities are known to pose serious health risks, and a calculated toxicity potential value that exceeds unity when using the threshold limits is concerning. The objective of this study was to assess the levels of PM_{2.5}, PM₁₀, the PM_{2.5}/PM₁₀ ratio, and the toxicity potential of a commercial area in Akure, Ondo State, Nigeria.

2. Materials and Methods

Akure is the capital city of Nigeria's Ondo state. Every year, the city experiences dry (November to March) and wet (April to October) seasons. The dry season (harmattan) is typically cold (9-16 °C), with dusty northeasterly trade winds from the Sahara desert transporting large amounts of dust for several days. The season is typically distinguished by high solar radiation and clear skies, moderate air temperatures, and no precipitation. Between April and mid-October, southwesterly winds from the Atlantic Ocean predominate. The research was carried out at the Federal College of Agriculture, Akure, REC campus commercial area (5 14 23.94 E 7 5 49.34 N). There are motorcycle mechanic, shoe making workshops, as well as a commercial shop (typing, book binding, etc.). A lot of generator usage occurs in these places. A low-cost Canāree A1 sensor an Intelligent Particle Sensor was used for the three-month monitoring (March to May 2022) of PM₁₀ and PM_{2.5} in the study for 6 h each day. The manufacturer's standard protocols were strictly followed. The generated data was statistically manipulated using Minitab and Excel 2013 software, producing basic summary reports and a bar chart, respectively.

2.1. Toxicity Potential

Toxicity potential (TP), is the ratio of evaluated ambient PM mass to the standard limit of ambient concentration [5]. It is useful in determining the harmful effects of pollutants on human health. It was calculated using Eqn (1), taking into account the World Health Organization 2021 Guidelines for PM₁₀ and PM_{2.5} ambient air quality standards: PM₁₀ (24h - 45 µg/m³ and Annual - 15 µg/m³) PM_{2.5} (24h - 5 µg/m³ and Annual - 5 µg/m³) [6].

$$\text{Toxicity Potential} = \frac{MPM}{SPM} \quad (1)$$

where: MPM is the measured particulate matter and SPM is the guideline limit set for PM₁₀ and PM_{2.5}

3. Results and Discussion




Figure 1: The Summary Report of PM₁₀, PM_{2.5}, and Their Ratios.




Figure 2: The PM₁₀ and PM_{2.5} Toxicity Potentials.

Figure 1 depicts the summary reports for PM₁₀, PM_{2.5}, and their respective ratios. The recorded PM_{2.5} and PM₁₀ levels for the study periods ranged from 1.26 to 469.80 µg/m³, PM_{2.5} - 1.26 to 419.13 µg/m³, and their ratios were observed as 0.49 and 1.00. It should be noted that the WHO 2021 guidelines assert that annual average PM_{2.5} and PM₁₀ concentrations must not exceed 5 and 15 µg/m³, in both, while 24-h average risks must not surpass 15 and 45 µg/m³, respectively. The majority of the 24-h PM₁₀ and PM_{2.5} values in the commercial area were discovered to be greater than the WHO recommendations in this study. Figure 1 depicts the summary reports for PM₁₀, PM_{2.5}, and their respective ratios. The high Particulate matter levels are expected due to the many anthropogenic activities within the study area, including the use of perfumes by people (customers and shop owners) within the surroundings, generator fume, and the center's proximity to a high traffic location, vehicular movements on the airport road. During this time, there were also many land preparations for the start of the new planting season. There was a lot of bush burning and soil dust movement. Long-distance transport from nearby

Table 1: Comparison of the results of our study with previous ones in Nigeria and Other countries.

S/N	Location	PM ₁₀ (µg/m ³)	PM _{2.5} (µg/m ³)	PM _{2.5} /PM ₁₀	Toxicity Potential	References
1.	FECA, Akure, Nigeria	73.23±53.94	68.58±50.64	0.93±0.02	PM ₁₀ -1.47 PM _{2.5} -2.74	This Study
2.	Kano, Nigeria	22.70-57.00	11.71-25.30	-	PM ₁₀ : 0.45-1.14 PM _{2.5} : 0.45-1.08	Ayua et al. [7]
3.	Chelyabinsk, Russia	6 and 64	5 to 56	0.85	-	Krupnova et al. [8]
4.	Helsinki and Stockholm	-	-	0.30-0.80	-	Adães and Pires [9]
5.	Tianjin, China	40.09 to 746	13.03 to 309	Average (0.41 ± 0.22 (Spring) – 0.43 ± 0.19 (winter),	-	Zhang et al. [10]
6.	China mainland	74.37-85.13	38.08-48.63	0.54-0.57	-	Fan et al., [11]
7.	Omu-Aran, Nigeria	59.32 – 473.52	11.42 – 32.40	-	PM ₁₀ : 0.13 – 0.36,	Fakinle et al. [12]

					PM _{2.5} : 0.33 – 2.64
8.	Idiroko Road, Nigeria	-	43.0±1.0 to 91.0±5.0 (day) 42.0±2.0 to 53.0±3.0 (evening)	-	PM _{2.5} : 1.36-3.64 Oghenovo et al. [13]

forest fires could have also contributed to mass concentrations during this time period. Table 1 compares our findings to previous research in Nigeria and other countries. Figure 1 depicts the summary reports for PM₁₀, PM_{2.5}, and their respective ratios. The average PM_{2.5}/PM₁₀ ratios in other countries were consistent to our studies, which range from 0.3 to 0.85 in the winter, spring, and summer seasons. The consistency could be explained by the fact that traffic-related sources are dominated by particle emissions and the constituents of PM_{2.5} and PM₁₀.

Figure 2 depicts the mean values of PM₁₀ and PM_{2.5} Toxicity Potentials (1.47 and (1.47 and 2.74). Our results were similar to those of Fakinle et al. [12] - PM_{2.5}: 0.13 - 0.36, and PM₁₀: 0.33 - 2.64 - but Oghenovo et al. [13] obtained a higher maximum level (3.64). The similarities in TP results revealed that the sources of particulate matter may not differ between locations. A TP value greater than one indicates that PM₁₀ and PM_{2.5} with a location pose a risk to people in the area. Several epidemiologic studies and toxicity research studies have found that a variety of chemical components and sources can have a negative impact on people's health. It is crucial to mention that the area surrounding this study is healthy, but good tracking of the environment in terms of PM should be done on a regular basis.

4. Conclusion

The PM_{2.5}, PM₁₀, PM_{2.5}/PM₁₀ ratio, and TP of a commercial area in Akure, Ondo State, Nigeria were measured for three months using a Canāree A1 low-cost sensor. The findings revealed that the WHO 2021 guidelines were exceeded. The PM_{2.5}/PM₁₀ ratios revealed that the presence of PM could be attributed to fumes from the generator and vehicles in the study area. A TP greater than one is a health concern, especially for the vulnerable (the sick, children, and the elderly). Constant monitoring is advised.

Author Contributions: Conceptualization, F.O.A.; methodology, F.O.A, K.M.A and A.A.; investigation and formal analysis, AaA, S.D.O, and S.A.; data curation, A.A. and S.D.O; writing—original draft preparation, F.O.A.; writing—review and editing, S.A and K.M.A.; visualization, F.O.A and A.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

Acknowledgments: The authors are grateful to Piera Systems, Canada for the provision of the low-cost sensor used for the monitoring.

References

1. Ndamitso, M.M., Iyaka, Y.A., Abdulkadir, A., and Abulude, F.O. Evaluation of Atmospheric PM₁₀ in the Southwest Region of Nigeria. *Current Applied Science and Technology*, 2021, Vol. 21 No. 2, 215-223
2. Zhang, M.-K., Liu, Z.-Y., and Wang, H. Use of Single Extraction Methods to Predict Bioavailability of Heavy Metals in Polluted Soils to Rice. *Communications in Soil Science and Plant Analysis*, 2010. 41, 820-831. <https://doi.org/10.1080/00103621003592341>

3. Fakinle, B.S., Sonibare, J.A., Akeredolu, F.A., Okedere, O.B., and Jimoda, L.A. Toxicity Potential of Particulates in the Airshed of Haulage Vehicle Park. *Global NEST Journal*. 2013. Vol 15, No 4, pp 466-473.
4. Abah, J., Mashebe, P., and Onjefu, S. A. Heavy Metals Pollution Status of the Katima Mulilo Urban Open Land Wastewater Disposal Centre and the Immediate Vicinity. 2020. *Cogent Environmental Science*, 6, Article ID: 1726093. <https://doi.org/10.1080/23311843.2020.1726093>
5. Sonibare J.A., Akeredolu F.A., Osibanjo O. and Latinwo I. ED-XRF Analysis of Total Suspended Particulates from Enamelware Manufacturing Industry, *American Journal of Applied Science*. 2005. 2(2), 573-578.
6. WHO. WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. World Health Organization. 2021. <https://apps.who.int/iris/handle/10665/345329>.
7. Ayua, T., Tyovenda, A. , Sombo, T. , Tikyaa, E. and Igbawua, T. Fine Particulate Matter and Heavy Metals Pollution Status in Ambient Air of Some Selected Industrial Sites in Northern Nigeria. *Journal of Geoscience and Environment Protection*. 2020. 8, 1-13. doi: [10.4236/gep.2020.88001](https://doi.org/10.4236/gep.2020.88001).
8. Krupnova, T.G.; Rakova, O.V.; Bondarenko, K.A.; Saifullin, A.F.; Popova, D.A.; Potgieter-Vermaak, S.; Godoi, R.H.M. Elemental Composition of PM_{2.5} and PM₁₀ and Health Risks Assessment in the Industrial Districts of Chelyabinsk, South Ural Region, Russia. *Int. J. Environ. Res. Public Health*. 2021, 18, 12354. <https://doi.org/10.3390/ijerph182312354>
9. Adães J, Pires JCM. Analysis and Modelling of PM_{2.5} Temporal and Spatial Behaviors in European Cities. *Sustainability*. 2019; 11(21):6019. <https://doi.org/10.3390/su11216019>
10. Zhang, J., Wu, L., Fang, X., Li, F., Yang, Z., Wang, T., Mao, H., and Wei, E.. Elemental Composition and Health Risk Assessment of PM₁₀ and PM_{2.5} in the Roadside Microenvironment in Tianjin, China. *Aerosol and Air Quality Research*, 2018, 18: 1817–1827. doi: 10.4209/aaqr.2017.10.0383
11. Fan, H., Zhao, C., Yang, Y., and Yang, X. Spatio-Temporal Variations of the PM_{2.5}/PM₁₀ Ratios and Its Application to Air Pollution Type Classification in China. *Front. Environ. Sci.*, 14 June 2021, Sec. *Atmosphere and Climate*. 2021, <https://doi.org/10.3389/fenvs.2021.692440>
12. Fakinle, B.S., Adebayo, B.M., Aremu, C.O., and Sonibare, J.A. Toxicity potential of particulate in the airshed of a university farm. IOP Conf. Series: Earth and Environmental Science, 2020, 445. 012036. doi:10.1088/1755-1315/445/1/012036
13. Oghenovo, I., Benson, N., Adedapo, A., and Ahmadu, O.F. Assessment of Particulate Matter (PM_{2.5}) in Residential Staff Quarters of Covenant University, Nigeria. OP Conf. Series: Journal of Physics: Conf. Series, 2019, 1299 012101. doi:10.1088/1742-6596/1299/1/012101