

Eucalyptus Genotypes Water Use and Site Interactions on Water Resource Sustainability and Productivity[†]

Rafael Rubilar ^{1,2,*}, Daniel Bozo ^{1,2}, Juan Valverde ¹, Matias Pincheira ³, Veronica Emhart ³, Alex Medina ³ and Hector Valenzuela ⁴

¹ Cooperativa de Productividad Forestal, Departamento de Silvicultura, Facultad de Ciencias Forestales, Universidad de Concepción, Chile; rafaelrubilar@udec.cl

² Centro Nacional de Excelencia para la Industria de la Madera (CENAMAD), Pontificia Universidad Católica de Chile, Vicuña Mackenna 4860, Santiago, Chile

³ Forestal Minico SpA, CMPC Forestal, Los Angeles, Chile

⁴ Forestal Arauco Co., Constitución, Chile

* Correspondence: rafaelrubilar@udec.cl; Tel.: +56 412204980

[†] Presented at the 3rd International Electronic Conference on Forests — Exploring New Discoveries and New Directions in Forests, 15 to 31 October 2022. Available online: <https://iecf2022.sciforum.net>

Abstract: Water use, water use efficiency and water use sustainability are challenging issues for expansion of Eucalyptus plantations around the world nowadays. Fast-growing species effect on water resources are of concern to society and forest industry wonders about effects on productivity and water sustainability under climate change. We evaluated growth, water use and water use efficiency response at canopy closure (3 to 4 years of age) of eight selected *E.globulus*, *E.nitens* and *E.nitens* × *globulus* (*E.gloni*) genotypes at four contrasting environments. Across sites & genotypes growth rate and water use ranged 35-70- m³/ha/yr and 350 to 900 mm ha⁻¹. Despite specific genotype cases, our results showed similar water use vs productivity relationships for a single taxon (90 m³ of water per m³ of wood) but suggested large differences among taxas across sites. Our results suggest that hybrids may provide large productivity benefits but may need a more precise understanding on their physiological mechanisms. Interesting opportunities exist for matching genotypes to sites in order to reduce environmental concerns on water resources use for Eucalyptus plantations.

Keywords: Water use; water sustainability; clonal; growth efficiency; eucalyptus

Citation: Rubilar, R.; Bozo, D.; Valverde, J.; Pincheira, M.; Emhart, V.; Medina, A.; Valenzuela, H. Eucalyptus genotypes water use and site interactions on water resource sustainability and productivity. *Environ. Sci. Proc.*, **2022**, *4*, x. <https://doi.org/10.3390/xxxxx>

Received: date

Accepted: date

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Eucalyptus is one of the most extensively planted genera worldwide and several concerns have been raised about the effects on water resources, water sustainability and survival and productivity of intensively managed plantations under climate change ([1, 2]). Despite several detailed ecophysiological assesments have been made over the last decade on the most extensively planted species[3-6], scarce information exist about the opportunity on selecting specific genotypes that may reducing the potential effects of plantations on water resources and their risk for sustained production of forest resources [7, 8]. The objective of our study was to understand the differential growth response, water use and water use efficiency of mediterranean Eucalyptus in order to evaluate opportunities on specific taxa and genotypes selection for increasing site water sustainability of intensivley managed plantations.

2. Materials and Methods

2.1. Site Characteristics

Four sites, from the regional validation genetic trials of CMPC and ARAUCO forest companies in Chile representing a gradient of soils of contrasting soil water holding capacity and atmospheric water demand, this last characteristic represented by the regional gradient in summer atmospheric vapor pressure deficit (Figure 1) were selected for evaluation of *Eucalyptus globulus*, *Eucalyptus nitens* and *Eucalyptus nitens* × *globulus* hybrids top genotypes. North sites considered evaluation of five *E. globulus* and three *E. nitens* × *globulus* hybrids clones at “Los Patos” (LP) trial farm located on granitic derived clay soils of the Coastal Range and “Vegas de Saldia” (VS) trial farm located on recent volcanic ash loamy soils of the Piedmont of the Andes. South sites considered evaluation of two *E. globulus* and four *E. nitens* × *globulus* clones, and two *E. nitens* family genotypes at “El Derrumbe” (ED) trial farm located on metamorphic derived clay soils of the Coastal Range and “San Andrés” (SA) trial farm located on red clay old volcanic ash of the Central Valley. All stands were established with the best early silvicultural treatments to secure full survival and early growth by considering subsoiling, fertilization and weed control up to crown closure (preplanting and postplanting). North stands had 4 years old and were planted at 1666 trees per hectare; South sites had 5 years old and were planted at 1430 trees per hectare.




Figure 1. Mean daily vapour pressure deficit during February summer month map representing the regional gradient in atmospheric water demand for all sites selected and genotypes within each site.

2.2. Annual growth & sapflow assessments

For each selected genotype and three replicates at each site, 36 trees (measurement plots of 6 × 6 trees) were measured for diameter at breast height (1.3 m height) and total height in march 2017, march 2018 and march 2019. Genotype annual volume growth ($\text{m}^3 \text{ wood ha}^{-1} \text{ yr}^{-1}$) was estimated from annual plot volume estimates scaled to an hectare level. From the diameter distribution of each genotype at each site, measured in 2017, six trees were selected for sapflow assessments. Granier sensors, located at DBH height on each selected tree, were monitored continuously and sap flux density records were obtained every 15 minutes. Annual sapwood area wood cores were obtained at DBH height from three buffer trees of each plot in order to develop a DBH vs sapwood area relationship. Estimates of water use at individual tree level were obtained from sapwood area and sap flux density estimates and scaling estimates to a plot and an hectare level to obtain water use estimates ($\text{m}^3 \text{ water ha}^{-1} \text{ yr}^{-1}$).

3. Results and Discussion

Across sites & genotypes growth rate ranged 35-70- $\text{m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$. *Eucalyptus nitens* showed the largest growth of all genotypes at southern sites ranging from 50 to 70 $\text{m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$, and *E. globulus* genotypes showed the highest growth at northern sites ranging from 30 to 65 $\text{m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$. The lowest average site productivity was observed at Los Patos

North clayey soils across sites and the highest average site productivity was observed at San Andrés sites mainly sustained by *Eucalyptus nitens* growth (Figure 2).




Figure 2. Average volume growth ($m^3\ ha^{-1}\ yr^{-1}$) of each genotype by taxa at a) Los Patos and b) Vegas de Saldia North sites, and c) El Derrumbe and d) San Andrés sites from 2017 to 2018.

Across sites & genotypes annual water used ranged 300 to 950 mm of water transpired annually. *Eucalyptus nitens* x *globulus* hybrid clones showed, in average, the largest water use ($675\ mm\ year^{-1}$), and except for a single genotype *E. globulus* showed the lowest average water use ($535\ mm\ year^{-1}$).




Figure 3. Average annual water use ($mm\ ha^{-1}$) for each selected genotype evaluated across all evaluated sites from 2017 to 2018.

Despite specific genotype cases our results showed similar water use vs productivity relationships for a single taxon. In fact, average annual water use reached in average 90

m³ per m³ of annual wood growth. However, analysis of specific genotypes within each taxa and among taxa suggest large differences across sites.

4. Conclusions

Our results suggest that hybrids may provide large productivity benefits but may need a more precise understanding on their physiological mechanisms. Interesting opportunities exist for matching genotypes to sites in order to reduce environmental concerns on water resources use for Eucalyptus plantations.

Author Contributions: Contributions for this manuscript consider Conceptualization, RR, VE; methodology, RR, AM, MP, HV; formal analysis, RR, DB, JV; investigation, RR, JV; writing—original draft preparation, RR; writing—review and editing, RR; supervision, RR, AM, HV; project administration, RR, AM, HV. All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded by the government of Chile via CONICYT Fondecyt Regular Project 1190835, and CONICYT FONDEF Project IT16I10087 and ANID BASAL FB210015, also funding for maintenance of this trials was provided by CMPC Forestal Mininco S.A., ARAUCO Co., the Forest Productivity Cooperative at Universidad de Concepción Chile and support of National Agency for Research and Development (ANID), Scholarship Program, DOCTORADO BECAS CHILE/2020-21202023.

Data Availability Statement: “Not applicable”.

Acknowledgments: We acknowledge the support of several colleagues of Forestal Mininco and Arauco forest companies that supported field work activities and maintaining field installations..

Conflicts of Interest: “The authors declare no conflict of interest.” “The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results”.

References

1. Ferraz, S.F.D., et al., *Effects of Eucalyptus plantations on streamflow in Brazil: Moving beyond the water use debate*. Forest Ecology and Management, 2019. **453**.
2. Tupinamba-Simoes, F., et al., *Assessment of drought effects on survival and growth dynamics in eucalypt commercial forestry using remote sensing photogrammetry. A showcase in Mato Grosso, Brazil*. Forest Ecology and Management, 2022. **505**.
3. Battaglia, M. and P. Sands, *Application of sensitivity analysis to a model of Eucalyptus globulus plantation productivity*. Ecological Modelling, 1998. **111**(2-3): p. 237-259.
4. Hingston, F.J., J.H. Galbraith, and G.M. Dimmock, *Application of the process-based model BIOMASS to Eucalyptus globulus subsp. globulus plantations on ex-farmland in south western Australia - I. Water use by trees and assessing risk of losses due to drought*. Forest Ecology and Management, 1998. **106**(2-3): p. 141-156.
5. Dye, P., *Water use efficiency in South African Eucalyptus plantations: A review*. Southern African Forestry Journal, 2000: p. 17-26.
6. Whitehead, D. and C.L. Beadle, *Physiological regulation of productivity and water use in Eucalyptus: a review*. Forest Ecology and Management, 2004. **193**(1-2): p. 113-140.
7. Rubilar, R., et al., *Climate and water availability impacts on early growth and growth efficiency of Eucalyptus genotypes: The importance of GxE interactions*. Forest Ecology and Management, 2020. **458**.
8. Butler, J.B., et al., *Climate Adaptation, Drought Susceptibility, and Genomic-Informed Predictions of Future Climate Refugia for the Australian Forest Tree Eucalyptus globulus*. Forests, 2022. **13**(4).