



Type of the Paper (Proceedings)

Potential geography and conservation of *Ipomoea beninensis*, an Endangered plant species for Benin (West Africa) [†]

Rodrigue Idohou ^{1,2,*}, Hospice Dassou ³, Gafarou Agounde², Guillaume Hounsou-Dindin² and Aristide Adomou ³

¹ Ecole de Gestion et de Production Végétale et Semencière, Université Nationale d'Agriculture, Bénin

² Laboratoire de Biomathématiques et d'Estimations Forestières, Faculté des Sciences Agronomiques, Université d'Abomey-Calavi, Bénin

³ Laboratoire de Botanique et Ecologie Végétale, Faculté des Sciences et Techniques, Université d'Abomey-Calavi, Bénin

* Correspondence: rodrigodohou@gmail.com ; Tel.: +229 97938303

† Presented at The 3rd International Electronic Conference on Forests — Exploring New Discoveries and New Directions in Forests

Abstract: The only endemic plant of Benin, *Ipomoea beninensis* Akoègn. Lisowski & Sinsin, is threatened in its natural habitats. This study assessed the suitability of the current and future habitat for its conservation countrywide. Maxent models were run using records added to environmental variables under present and two climates. The results showed that the most suitable areas for *I. beninensis* will be mainly in the phytodistrict of southern and northern Borgou. The species could lose 9% and 13.6% of its suitable habitats under RCP4.5 and RCP8.5, respectively. Urgent and timely strategies are needed to save the remaining population of the species.

Keywords: conservation; climate changes; threatened taxa; West Africa.

Citation: Idohou, R. ; Dassou, H. ; Agounde, G. ; Hounsou-Dindin, G. ; Adomou, A. Potential geography and conservation of *Ipomoea beninensis*, an Endangered plant species for Benin (West Africa). *Environ. Sci. Proc.* **2022**, *4*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor:

Published:

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Worldwide, the biological diversity is evolving in an environment that is changing at an unstoppable and disturbing pace [1]. The world is undergoing significant lifestyle change, closely linked to both natural phenomena and anthropogenic factors [2]. In the natural ecosystems, the function and structure have been affected by human activities and climate changes, the impacts of which will get worse in the near future [3]. As a result, the world is expected to experience an unexpected vulnerability of all features of biodiversity, climate systems, and the people who depend on them for their daily needs.

Nonetheless, there is increasing evidence that variations in climate variables such as precipitation and temperature will affect the biological biodiversity and the geographic distribution of suitable habitats for species [4]. *Ipomoea beninensis* Akoègninou, Lisowski, and Sinsin is an endemic plant (Convolvulaceae), first described in Benin in 1999 [5]. It grows in forests and wooded savannas in the Guineo-Sudanese and Sudanese zones in Benin [5]. It plays an important role in meeting the needs of local populations, especially in rural areas, whether environmental, social, or cultural, with its rhizome being used to treat chickenpox and malaria and as fodder [6]. For some years, its populations have also been exposed to bushfires that affect the availability of its above-ground parts, justifying its poor management and leading to its rapid decline in

natural habitats [6]. To better understand the very poorly known distribution of the species and improve its monitoring in biodiversity monitoring campaigns, this study assessed the current distribution of *Ipomoea beninensis* and projected its future distribution under climate scenarios.

2. Materials and Methods

2.1 Study area

The study was conducted in the two ecological zones of the Guinea-Sudanese transition area (7°30'N to 9°45'N) and the Sudanese area (9°45'N to 12°25'N) in Benin.

2.2 Occurrence Data

In whole 189 occurrence records were gathered from the field, and supplemented by those (69) downloaded from online resources (GBIF: <http://www.gbif.org>). Data were quality checked [7]. In addition, we discarded duplicate records and those with obvious errors.

2.3 Environmental variables

In this study, 21 bioclimatic variables were downloaded (≈ 1 km) from the Africlim website (www.africlim.org) for both baseline and future climates. For future climate projections, the Ensemble model [8] was used and the projections were made for 2055 under RCP 4.5 and 8.5 scenario. In addition, soil layers were obtained at the same resolution from (<https://databasin.org>) and from Africa Soil Profiles Database (<https://www.isric.org>), respectively. We discarded highly correlated variables (≥ 0.8) by using Pearson's rank correlation coefficients and converted to Ascii format.

2.4 Species distribution models' development

The Maximum Entropy modeling approach, MaxEnt version 3.4.4k [8] was used to map the present-day and future distribution of *I. beninensis*. Linear, quadratic, product, hinge, and threshold functions of predictor variables were checked and variable importance was assessed using a jackknife analysis [9]. We restricted the selection of background points using the regularization of 10,000 background points. To easy the interpretation of the findings, we specified the Maxent' output format to a logistic form. All models were run ten (10) times. We computed the performance of the models using the area under the receiver operating characteristics curve (AUC) and the partial receiver operator characteristics (pROC) [10]. The dynamic of these patterns of suitability classes were quantified.

3. Results

3.1 Variables' contribution and models performance

The 10-time replications highlighted five (05) variables as most contributing to the distribution of *I. beninensis* : mean diurnal range in temperature, Temperature seasonality, mean temperature of warmest quarter), and rainfall of the wettest month), and elevation. Overall, variables related to temperature were more represented in the selection. The performance of the models assessed through AUC and partial ROC revealed a value of AUC of 0.84 and a value of AUC ratio was 1.6; thus, indicating that our models have very good predictive power.

3.2 Current and future distribution of *I. beninensis* across Benin

The models showed under the present condition (Fig. 4a) that the highly suitable habitat for *I. beninensis* were mainly located in the phytodistricts of Borgou (south and north). However, the phytodistrict of Bassila as well as the Atacora chain phytodistrict were found to be favorable for the conservation of the species. The moderately suitable habitats were located in the same phytodistricts

abovementioned. Analyses based on projecting the final models into the future (by 2055) scenarios revealed that (Fig. 1) the highly suitable habitats for the species will be more concentrated in the phytodistricts of Borgou (south and north) for both scenarios. In addition, part of the highly suitable area will remain alongside the East sides of the study area. Assessment of the dynamics of the suitability under future scenarios revealed that more suitable areas will be lost under RCP8.5 compared to RCP4.5.




Figure 1. Distribution maps for of *Ipomoea beninensis*; left: present-day distribution; middle: future distribution under RCP 4.5; right: future distribution under RCP 8.5.

4. Discussion

This study assessed the potential geography and conservation areas of *Ipomoea beninensis* in Benin. Globally, models yielded very good performance (AUC value >0.80 and AUC ratio was 1.6). Our findings joined the previous reported by [1], [7] and [11] on tropical species.

Especially, five variables were identified as the most contributing to the distribution of *I. beninensis*. The temperature (85.5%) through its mean diurnal range, seasonality, and its mean warmest quarter, as well as the rainfall wettest month (1.1%), highly contributed to the models. These were associated with the elevation, which showed a high contribution (13.4%). This indicates that the species can withstand the annual temperature between 26 and 28°C, annual rainfall between 1150 and 1300 mm, and the elevation range between 150 and 200 m. Overall, these variables largely represent climatic conditions of *I. beninensis* trees distribution, providing insights into underlying climatic factors which delimit its suitable habitats. These results were in line with the findings of [1], [12] and [13], who state that abiotic conditions represent one of the most important factors that

determine the area in which a species is found. Moreover, high elevation locations have tended to be less impacted by habitat degradation due to direct human impact, nevertheless, climate change threatens to increase these risks as upland areas become more suitable for agriculture [14]. The species thus has a restricted ecological amplitude for environmental conditions and consequently a very weak ecological plasticity highlighting its vulnerability to climate threats challenge.

The present results aligned with previous studies that reported the populations of *I. beninensis* as declining, due to the conversion of lands where the species is found into farmlands and the over-exploitation of its organs for the treatment of human pathologies and as fodder for animals [6] and will serve for its management practices (in-situ and ex-situ). In addition, under the future climate conditions, only one-third of the protected areas (N'dali and Ouémé Supérieur protected areas) could be suitable for its distribution. In fact, when habitats of an endemic species are lost by mismanagement and various other human activities combined with the climate change threats, the distribution ranges, population sizes, and genetic variability of the species will be reduced and will become vulnerable to extinction at a faster rate than other species [15]. Consequently, this species must be carefully monitored and managed to maintain biodiversity.

Results of this study did not address the relationship between biotic factors and their complex interactions with the abiotic factors, which influence the distribution range of species with much of the conditions that make up niche [16]. Thus, the limiting factors of species distribution are far more than environmental factors, highlighting the impossible to measure the true niche of a species completely. Therefore, modelling approaches should always be made with caution, and choosing modelling variables must be prioritized [16]. Even though these variables may not address those relations, the colonization and extinction dynamics assessed in this study will be relevant to *I. beninensis* management policy, in particular, to identify new areas potentially suitable for its conservation.

5. Conclusion

The study suggests that despite the projected spatiotemporal dynamics, linked to climate change, the environmental conditions in Benin will remain favorable to the cultivation and conservation of *I. beninensis* by year-2055. This argues that conservation efforts should be focused on species and habitats threatened by human activities, through the conservation of suitable habitats. We encourage the creation of a reserve or the reintroduction of *I. beninensis* in areas with high environmental suitability within the area of occurrence including the phytodistrict of Borgou (southern and northern), Bassila, and Atacora chain.

Author Contributions: Conceptualization, R.I. and D.H.; methodology, R.I.; software, A.G.; validation, R.I. and D.H.; formal analysis, R.I.; investigation, D.H.; resources, R.I. and D.H.; data curation, R.I.; writing—original draft preparation, A.G.; writing—review and editing, R.I.; visualization, H.D.G.; supervision, A.A.; funding acquisition, R.I. and D.H. All authors have read and agreed to the published version of the manuscript.”

Funding: This work was supported by the Rufford Small Grant Foundation through the grant N° 20742-1. RI acknowledges support from the Rufford Foundation (Grant 31042-D).

Institutional Review Board Statement: Not applicable

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable

Acknowledgments: In this section, you can acknowledge any support given which is not covered by the author contribution or funding sections. This may include administrative and technical support, or donations in kind (e.g., materials used for experiments).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Assogba, D.; Idohou R.; Chirwa P. Assogbadjo A. On opportunities and challenges to conserve the African baobab under present and future climates in Benin (West Africa). *J Arid Environ* **2022**, *198*, 104692. 171-178
2. Pereira, H.M.; Navarro, L.M.; Martins, I.S. Global biodiversity change: the bad, the good, and the unknown. *Annu Rev Environ Res* **2012**, *37*; 25-50. 179-180
3. Ogle, J.; Delporte, D.; Sanger, H. Quantifying the sustainability of urban growth and form through time: An algorithmic analysis of a city's development. *Appl Geogr* **2017** *88*; 1-14. 181-182
4. IPCC. Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte; V.; P. Zhai; A. Pirani; S.L. Connors; C. Péan; S. Berger; N. Caud; Y. Chen; L. Goldfarb; M.I. Gomis; M. Huang; K. Leitzell; E. Lonnoy; J.B.R. Matthews; T.K. Maycock; T. Waterfield; O. Yelekçi; R. Yu; and B. Zhou (eds.)]. Cambridge University Press. **2021** 183-186
5. Akoègninou, A.; Van Der Burg, J.; Van Der Maesen, Le. Flore analytique du Bénin, Cotonou & Wageningen: Backhuys Publishers **2006**. 1063 p 187-188
6. Dassou, H.G.; Idohou, R.; Adomou, A.C.; Ouachinou, J.M.-AS; Yédomonhan, H. Indigenous knowledge and local practices concerning the endemic plant *Ipomoea beninensis* Akoègn.; Lisowski & Sinsin (Convolvulaceae): an initial assessment for its conservation in Benin. *Flora et Vegetatio Sudano-Sambesica* **2020**, *23*, 3-13. 189-191
7. Idohou, R.; Assogbadjo, A.E.; Glèlè Kakai, R.; Peterson, A.T. Spatio-temporal dynamic of suitable areas for species conservation in West Africa: eight economically important wild palms under present and future climates. *Agroforest Syst* **2017**, *91*(3), 527-540. 192-194
8. Platts, P.J.; Omeny, P.A.; Marchant, R. AFRICLIM: high-resolution climate projections for ecological applications in Africa. *Afr J Ecol* **2015**, *53*, 103-108. 195-196
9. Phillips, S.J.; Anderson, R.P.; Schapire, R.E. Maximum entropy modelling of species geographic distributions. *Ecol Model* **2006**, *190*, 231-259. 197-198
10. Peterson, A.T.; Papeş, M.; Soberón, J. Rethinking receiver operating characteristic analysis applications in ecological niche modeling. *Ecol Model* **2008**, *213*(1): 63-72. 199-200
11. Wouyou, H.G.; Lokonon, B.E.; Idohou, R.; Zossou-Akete, A.G.; Assogbadjo, A.E.; Glèlè Kakai, R. Predicting the potential impacts of climate change on the endangered *Caesalpinia bonduc* (L.) Roxb in Benin (West Africa). *Heliyon* **2022**; *8*(3); e09022. 201-203
12. Krigas, N.; Tsiafouli, M.A.; Katsoulis, G.; Votsi, N-E.; van Kleunen, M. Investigating the invasion pattern of the alien plant *Solanum elaeagnifolium* Cav. (Silverleaf Nightshade): environmental and human-induced drivers. *Plants* **2021**, *10*(4), 805. 204-206
13. Moshobane, M.C.; Mudereri, B.T.; Mukundamago, M.; Chitata, T. Predicting future distribution patterns of *Jatropha gossypifolia* L. in South Africa in response to climate change. *S Afr J Bot* **2022**, *146*, 417-425. 207-208
14. Pradhan A; Chan C; Roul PK; Halbrendt J; Sipes B (2018) Potential of conservation agriculture (CA) for climate change adaptation and food security under rainfed uplands of India: A transdisciplinary approach. *Agric Syst* **2018**; *163*; 27-35. 209-211
15. İşik, K. Rare and endemic species: why are they prone to extinction? *Turk J Bot* **2011**; *35*(4); 411-417. 212
16. Zangiabadi, S.; Zaremaivan, H.; Brotons, L.; Mostafavi, H.; Ranjbar, H. Using climatic variables alone overestimate climate change impacts on predicting distribution of an endemic species. *PLoS ONE* **2021**; *16*(9); e0256918. 213-215