

Proceeding Paper

Synthesis and Chemosensory Studies of a Heterocyclic Thiosemicarbazone as a New Tributyltin Optical Chemosensor [†]

Rui P. C. L. Sousa, Susana P. G. Costa, Rita B. Figueira and M. Manuela M. Raposo *

Centre of Chemistry, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal

* Correspondence: mfox@quimica.uminho.pt

[†] Presented at the 26th International Electronic Conference on Synthetic Organic Chemistry; Available online: <https://ecsoc-26.sciforum.net>.

Abstract: Thiosemicarbazones are a versatile type of organic compounds which are known by their coordination ability with different types of analytes, due to sulfur and nitrogen heteroatoms presence. Therefore, the functionalization of thiosemicarbazones with heterocyclic moieties can be a promising route to develop new optical chemosensors. Tributyltin (TBT) is an antifouling component of paints which is acutely toxic to aquatic environments, being quickly absorbed by microorganisms and causing problems such as *imposex*. Herein, the synthesis of a novel heterocyclic thiosemicarbazone, functionalized with a quinoline moiety, is reported to assess the potential of this recognition moiety for TBT optical chemosensing. A preliminary chemosensory study in acetonitrile solution was performed showing that 50 equivalents of TBT were needed to induce a change of color from colorless to yellow. Spectrophotometric titration was performed to assess the concentration of TBT necessary for a maximum optical signal, revealing that 100 equivalents of TBT were necessary to reach maximum absorbance, although it was able to respond with a detectable color change to a TBT concentration as low as 10 μ M.

Keywords: Tributyltin; biocide; colorimetric chemosensor; thiosemicarbazone; quinoline; synthesis

Citation: Sousa, R.P.C.L.; Costa, S.P.G.; Figueira, R.B.; Raposo, M.M. Synthesis and Chemosensory Studies of a Heterocyclic Thiosemicarbazone as a New Tributyltin Optical Chemosensor. *2022*, *4*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s):

Published: 15 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Optical chemosensors are pursued worldwide due to advantages over other types of sensors such as low-cost and selectivity [1]. Within this field, colorimetric chemosensors show the possibility of “naked eye” detection [1,2], which is extremely relevant and advantageous to reach a preliminary qualitative detection of a desired analyte. Besides, this type of sensor also may allow a fast and simple quantitative detection [2]. Colorimetric optical chemosensors are often based on organic chromophores, that can be modified towards the detection of a particular analyte.

Thiourea derivatives, particularly thiosemicarbazones, are molecules of particular interest in the sensing field due to the conjugation of heteroatoms with electronic properties that can be tuned by the presence of electron donor/withdrawing groups [3–5]. The combination of this core with π -conjugated bridges can yield selective and sensitive optical chemosensors for different ions. Particularly, the functionalization of a thiosemicarbazone with a heterocyclic ring such as quinoline, can be interesting for the development of colorimetric chemosensors, due to the combination of the coordination ability of sulfur and nitrogen heteroatoms from the thiosemicarbazone with a π -conjugated heterocyclic group.

Tributyltin (TBT) is an antifouling component of paints used on ships, vessels, and submersed structures to avoid biofouling [6]. This compound is a biocide that prevents accumulation of microorganisms on the abovementioned structures. However, it was

found that TBT is extremely toxic to aquatic organisms as bacteria, fishes or algae [7–12]. TBT is quickly absorbed by microorganisms and induces long-term problems in aquatic living beings such as *imposex* i.e., superimposition of male sexual characteristics on female marine gastropods [13,14]. The bioaccumulation of TBT in gastropods results in higher testosterone levels causing an endocrine disrupting effect [15].

Nowadays, TBT monitoring is only detected and quantified by sampling and analysis in laboratory by chromatographic techniques, such as Liquid Chromatography-Mass Spectrometry (LC-MS) and Gas Chromatography-Mass Spectrometry (GC-MS)- These techniques require expensive equipment, expert operators, and long procedures with several steps, such as extraction, preconcentration and derivatization [16]. Other attempts to monitor TBT include analysis by graphite furnace atomic absorption spectrometry with Zeeman correction (ZGFAAS), but this method share the same disadvantages [17]. Therefore, TBT optical chemosensors would be a huge and interesting advantage to build devices capable of in-situ TBT detection.

This work reports the synthesis of a novel heterocyclic thiosemicarbazone, functionalized with a quinoline moiety, to assess the potential of this recognition moiety for TBT optical chemosensing. The new compound was obtained by a condensation reaction, between quinoline-2-carboxaldehyde and *N*-phenylhydrazinecarbothioamide, and its chemosensory ability was studied in the presence of TBT in acetonitrile solution. A spectrophotometric titration was also performed to assess the concentration of TBT necessary for a maximum optical signal.

2. Experimental Section

2.1. Methods and Materials

Melting points were measured on a Stuart SMP3 melting point apparatus. TLC analysis was carried out on 0.20 mm thick precoated silica plates (Macherey-Nagel), and spots were observed under UV light on a CN-15 camera (Vilber Lourmat). UV-Vis absorption spectra (200–700 nm) were obtained using Shimadzu UV/3101PC spectrophotometer. NMR spectra were obtained on a Bruker Avance III 400 at an operating frequency of 400 MHz for ^1H using the solvent peak as internal reference at 25 °C. All chemical shifts are given in ppm using $\delta_{\text{H Me}_4\text{Si}} = 0$ ppm as reference and *J* values are given in Hz. Assignments were supported by spin decoupling-double resonance and bidimensional heteronuclear correlation techniques. All commercial reagents and solvents were used as received.

2.2. Synthesis of *N*-phenyl-2-(quinolin-2-ylmethylene)hydrazine-1-carbothioamide 1

Quinoline-2-carboxaldehyde **2** and *N*-phenylhydrazinecarbothioamide **3** in equal amounts (0.318 mmol) were dissolved in 10 mL of MeOH at room temperature. The reaction mixture was stirred for 8 h, and then for more 6 h at 60 °C. After cooling, the precipitated compound was filtered, dried and obtained as a brown solid in 17% yield (0.017 g).

Mp: 160 °C. ^1H NMR (DMSO-*d*₆, 400MHz): δ = 7.24 (t, *J* = 7.6 Hz, 1H, H4), 7.40 (dt, *J* = 7.6 and 1.6 Hz, 2H, H3 + H5), 7.55 (dd, *J* = 8.0 and 0.4 Hz, 2H, H2 + H6), 7.62 (dt, *J* = 7.2 and 1.2 Hz, 1H, H6'), 7.78 (dt, *J* = 7.2 and 1.2 Hz, 1H, H7'), 7.99 (dd, *J* = 8.0 and 0.4 Hz, 1H, H5'), 8.02 (d, *J* = 8.4 Hz, 1H, H8'), 8.33 (s, 1H, N=CH), 8.38 (d, *J* = 8.8 Hz, 1H, H4'), 8.59 (d, *J* = 8.4 Hz, 1H, H3'), 10.38 (s, 1H, NH-C=S), 12.17 (s, 1H, NH-N) ppm.


2.3. Preliminary Chemosensory Studies

For the preliminary test, 50 equivalents of TBT (50 μL , 1×10^{-1} M) were added to an acetonitrile (ACN) solution of the new thiosemicarbazone **1** (1 mL, 1×10^{-4} M). The color/fluorescence changes were assessed by visual inspection and in a UV-vis chamber under ultraviolet light at 312 nm. Spectrophotometric titration was performed with a sequential addition of TBT (10^{-2} M) to an ACN solution of compound **1** (3 mL, 1×10^{-5} M). Absorbance spectra were collected, until a plateau was reached.

3. Results and Discussion

3.1. Synthesis of *N*-phenyl-2-(quinolin-2-ylmethylene)hydrazine-1-carbothioamide **1**

The synthesis of the new thiosemicarbazone **1** was performed with a mixture of quinoline-2-carboxaldehyde **2** and *N*-phenylhydrazinecarbothioamide **3** in equal amounts (0.318 mmol). The two precursors were dissolved in MeOH (10 mL) and stirred at room temperature for 8 h. However, TLC analysis showed that the reaction did not progress, so it was stirred for more 6 h at 60 °C. After this time, the pure product **1**, was obtained as a brown solid, which was filtered, from the cold reaction mixture, in 17% yield (Scheme 1). The novel thiosemicarbazone **1** was characterized by ¹H NMR.



Scheme 1. Synthesis of thiosemicarbazone **1**.

¹H NMR spectrum shows the characteristic signals for NH protons at 12.17 and 10.38 ppm. Imine N=CH proton appears as a singlet at 8.33 ppm. Phenyl group protons appear at a smaller chemical shift at 7.24 (H₄), 7.40 (H₃ and H₅), and 7.55 (H₂ and H₆) ppm, while quinoline protons show higher chemical shifts with H_{6'} at 7.62 ppm, H_{7'} at 7.78 ppm, H_{5'} at 7.99 ppm, H_{8'} at 8.02 ppm, H_{4'} at 8.38 ppm, and H_{3'} at 8.59 ppm.

3.2. Studies of the Chemosensory Ability of Thiosemicarbazone **1** for TBT

The chemosensory ability of the new thiosemicarbazone **1** was studied in the presence of TBT. To an ACN solution of the compound (1 mL, 10⁻⁴ M), 50 equivalents of TBT (50 μL, 10⁻¹ M) were added. The optical response was analyzed by visual inspection and in a UV-vis chamber under ultraviolet light at 312 nm. A color change from colorless to yellow was observed (vide Figure 1). No changes in fluorescence were found.




Figure 1. Preliminary TBT chemosensory test for the new thiosemicarbazone **1**: left—compound **1**; right—compound **1** + 50 equivalents of TBT.

A spectrophotometric titration was performed to assess the number of equivalents necessary to reach the maximum optical signal (vide Figure 2). A sequential addition of a TBT solution (10⁻² M) to an ACN solution of the compound (10⁻⁵ M) was conducted and

the absorbance spectra were collected. It was observed that 100 equivalents were necessary to reach the absorbance plateau. However, the addition of 1 equivalent (10 μM) was enough to detect a change in the optical signal, showing this compound has a large range of concentration in which TBT can be detected.




Figure 2. Spectrophotometric titration of thiosemicarbazone **1** with TBT.

4. Conclusions

In this work, a new thiosemicarbazone was successfully synthesized by a condensation reaction between quinoline-2-carboxyaldehyde and N-phenylhydrazinecarbothioamide. The compound was tested in the presence of the biocide TBT and the preliminary test revealed a color change from colorless to yellow. A spectrophotometric titration was performed to assess the number of equivalents necessary to reach the plateau of the absorbance spectra. It was found that 100 equivalents of TBT were necessary to reach the maximum optical signal. However, this colorimetric probe was able to detect a change in the optical signal with a concentration as low as 10 μM . The wide range of concentration that can result in optical changes, allow to conclude that compound **1** shows potential to be used as a TBT chemosensor.

Author Contributions: Conceptualization, S.P.G.C., M.M.M.R.; methodology, R.P.C.L.S., S.P.G.C., M.M.M.R.; formal analysis, R.P.C.L.S., S.P.G.C., M.M.M.R.; investigation, R.P.C.L.S.; resources, S.P.G.C., R.B.F., M.M.M.R.; writing—original draft preparation, R.P.C.L.S.; writing—review and editing, S.P.G.C., R.B.F., M.M.M.R.; supervision, S.P.G.C., R.B.F., M.M.M.R.; project administration, S.P.G.C., R.B.F., M.M.M.R.; funding acquisition, S.P.G.C., R.B.F., M.M.M.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Fundação para a Ciência e Tecnologia (FCT) and FEDER (European Fund for Regional Development)-COMPETE-QRENEU for financial support through the Chemistry Research Centre of the University of Minho (Ref. CQ/UM UID/QUI/00686/2020), and a PhD grant to R. P. C. L. Sousa (SFRH/BD/145639/2019).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement:

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kaur, B.; Kaur, N.; Kumar, S. Colorimetric metal ion sensors—A comprehensive review of the years 2011–2016. *Coord. Chem. Rev.* **2018**, *358*, 13–69, <https://doi.org/10.1016/j.ccr.2017.12.002>.
2. Wang, B.; Anslyn, E. Colorimetric sensor design. In *Chemosensors: Principles, Strategies, and Applications*; 2011; pp. 275–295.
3. Raposo, M.M.M.; García-Acosta, B.; Ábalos, T.; Calero, P.; Martínez-Máñez, R.; Ros-Lis, J.V.; Soto, J. Synthesis and Study of the Use of Heterocyclic Thiosemicarbazones As Signaling Scaffolding for the Recognition of Anions. *J. Org. Chem.* **2010**, *75*, 2922–2933, <https://doi.org/10.1021/jo100082k>.
4. Santos-Figueroa, L.E.; Moragues, M.E.; Raposo, M.M.M.; Batista, R.M.F.; Costa, S.P.G.; Ferreira, R.C.M.; Sancenón, F.; Martínez-Máñez, R.; Ros-Lis, J.V.; Soto, J. Synthesis and evaluation of thiosemicarbazones functionalized with furyl moieties as new chemosensors for anion recognition. *Org. Biomol. Chem.* **2012**, *10*, 7418–7428, <https://doi.org/10.1039/c2ob26200b>.
5. Santos-Figueroa, L.E.; Moragues, M.E.; Raposo, M.M.M.; Batista, R.M.; Ferreira, R.C.M.; Costa, S.P.; Sancenón, F.; Martínez-Máñez, R.; Soto, J.; Ros-Lis, J.V. Synthesis and evaluation of fluorimetric and colorimetric chemosensors for anions based on (oligo)thienyl-thiosemicarbazones. *Tetrahedron* **2012**, *68*, 7179–7186, <https://doi.org/10.1016/j.tet.2012.06.021>.
6. Sousa, R.P.C.L.; Figueira, R.B.; Costa, S.P.G.; Raposo, M.M.M. Optical Fiber Sensors for Biocide Monitoring: Examples, Transduction Materials, and Prospects. *ACS Sens.* **2020**, *5*, 3678–3709, <https://doi.org/10.1021/acssensors.0c01615>.
7. Antizar-Ladislao, B. Environmental levels, toxicity and human exposure to tributyltin (TBT)-contaminated marine environment. A review. *Environ. Int.* **2008**, *34*, 292–308, <https://doi.org/10.1016/j.envint.2007.09.005>.
8. Fent, K. Ecotoxicology of Organotin Compounds. *Crit. Rev. Toxicol.* **1996**, *26*, 3–117, <https://doi.org/10.3109/10408449609089891>.
9. Cooke, G.M. Toxicology of Tributyltin in Mammalian Animal Models. *Immunol. Endocr. Metab. Agents Med. Chem.* **2006**, *6*, 63–71, <https://doi.org/10.2174/187152206775528815>.
10. Boyer, I.J. Toxicity of dibutyltin, tributyltin and other organotin compounds to humans and to experimental animals. *Toxicology* **1989**, *55*, 253–298, [https://doi.org/10.1016/0300-483x\(89\)90018-8](https://doi.org/10.1016/0300-483x(89)90018-8).
11. Tornero, V.; Hanke, G. Chemical contaminants entering the marine environment from sea-based sources: A review with a focus on European seas. *Mar. Pollut. Bull.* **2016**, *112*, 17–38, <https://doi.org/10.1016/j.marpolbul.2016.06.091>.
12. Cima, F.; Bragadin, M.; Ballarin, L. Toxic effects of new antifouling compounds on tunicate haemocytes: I. Sea-Nine 211™ and chlorothalonil. *Aquat. Toxicol.* **2008**, *86*, 299–312, <https://doi.org/10.1016/j.aquatox.2007.11.010>.
13. Artifon, V.; Castro, B.; Fillmann, G. Spatiotemporal appraisal of TBT contamination and imposex along a tropical bay (Todos Santos Bay, Brazil). *Environ. Sci. Pollut. Res.* **2016**, *23*, 16047–16055, <https://doi.org/10.1007/s11356-016-6745-7>.
14. Laranjeiro, F.; Sánchez-Marín, P.; Oliveira, I.B.; Galante-Oliveira, S.; Barroso, C. Fifteen years of imposex and tributyltin pollution monitoring along the Portuguese coast. *Environ. Pollut.* **2018**, *232*, 411–421, <https://doi.org/10.1016/j.envpol.2017.09.056>.
15. Matthiessen, P.; Gibbs, P.E. Critical appraisal of the evidence for tributyltin-mediated endocrine disruption in mollusks. *Environ. Toxicol. Chem.* **1998**, *17*, 37–43. [https://doi.org/10.1897/1551-5028\(1998\)017<0037:CAOTEF>2.3.CO;2](https://doi.org/10.1897/1551-5028(1998)017<0037:CAOTEF>2.3.CO;2).
16. Cole, R.F.; Mills, G.A.; Parker, R.; Bolam, T.; Birchenough, A.; Kröger, S.; Fones, G.R. Trends in the analysis and monitoring of organotins in the aquatic environment. *Trends Environ. Anal. Chem.* **2015**, *8*, 1–11, <https://doi.org/10.1016/j.teac.2015.05.001>.
17. Oliver, A.L.-S.; Sanz-Landaluze, J.; Muñoz-Olivas, R.; Guinea, J.; Cámara, C. Zebrafish larvae as a model for the evaluation of inorganic arsenic and tributyltin bioconcentration. *Water Res.* **2011**, *45*, 6515–6524, <https://doi.org/10.1016/j.watres.2011.09.052>.