

Relationship between precipitation just above the lava dome and displacement of the dome using X-band MP radar at Unzen Fugendake

Rikuto Daikai^{1,*}, Gomez Christopher^{1,2}, Balazs Bradak¹, Norifumi Hotta³ and Yoshinori Shinohara⁴

¹ Graduate School of Oceanology, Sediment Hazards and Disaster Risk Laboratory, Kobe University, Kobe, Japan; christophergomez@bear.kobe-u.ac.jp (C.G.); bradak.b@port.kobe-u.ac.jp (B.B.)

² Faculty of Geography, Universitas Gadjah mada, Yogyakarta, Indonesia

³ Faculty of Agriculture, The University of Tokyo, Tokyo, Japan, hotta.norihumi@fr.a.u-tokyo.ac.jp

⁴ Faculty of Agriculture, University of Miyazaki, Miyazaki, Japan, shinohara@cc.miyazaki-u.ac.jp

* Corresponding: rikuto.d221@gmail.com

Abstract

Since its last eruption from 1990-1995, Unzen Volcano (Shimabara Peninsula, Japan) has been quiescent. At its summit, a complex Dacitic dome spreads towards the East, in the direction of the Mizunashigawa-valley. In precarious equilibrium, sliding over previously erupted material, the dome has been generating rockfalls, and as the surrounding gullies have been eroding headwards, the stability of the dome is further reduced. Even if the volcano is in a dormant stage, its monitoring is therefore essential for disaster risk management. Therefore, the present contribution aims to (1) quantify the minute dome movement as a whole and (2) divided by lobe, in order to understand deformation; and (3) calculate what is the link between rainfalls and the dome movement. The method relies on the Unzen GBSAR system (Ground Based radar interferometry system) and on hourly rainfalls from raingages stations at Unzen Volcano. As a result, the authors have identified that (1) there is a time-delay between rainfall events and dome movements, and that peak rainfall alone is not sufficient to trigger dome movement; (2) the lower part of the dome rises and falls more rapidly than the upper part of the dome when rainfall is less than 100 mm/48 hours, and (2) the upper and lower parts of the dome move up and down at the same level when rainfall exceeds 100 mm/48 hours. In turn, when rainfall exceeds 250 mm/48 hours, then the upper part of the dome also displays further downward movement, so that the entire dome might be moving down like an accordion.

Keywords: lave dome; lave dome collapse; Continuous monitoring; Unzen volcano

Citation: Rikuto, D.; Gomez, C.; Balazs, B.; Norifumi, H.; Yoshinori, S. Relationship between precipitation just above the lava dome and displacement of the dome using X-band MP radar at Unzen Fugendake. *Proceedings* **2023**, *69*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Firstname Last-name

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Lava-dome collapse is a serious threat in volcanic regions [1]. The collapse of the dome can be devastating to settlements and infrastructure at the foot of the mountain. There are four possible causes of lava dome collapse: (1) excessive steepening due to gravity, (2) dome failure due to volcanic gas pressure, (3) rainfall, and (4) the dome and its underlying structure [2]. Among these, volcanic domes are considered susceptible to rainfall, and the dome collapse of March 20th, 2000, at Soufriere Volcano, Montserrat, (Lesser Antilles, UK) was considered to have been caused by rainfall before and during the collapse [3]. The mechanism of dome collapse due to rainfall is thought to be a combination of destabilization by erosion of the dome front, steam and water action on the potential failure surface inside the dome, rapid cooling of the lava, and small phreatomagmatic explosions [3], but the details in individual lava domes are unknown. At Unzen Volcano

(Kyushu-Island, Japan), the lava dome that developed at the summit crater of Mount Unzen during its last eruption between 1991 and 1995 has remained stable since the end of the eruption without major collapse. However, it continues to slide downwards. In quiescent periods, gravity waxed by water from rainfalls or other sources can be hypothesized to be the main origin of dome movement. Especially because even during the eruption, rainfall is thought to have helped triggering some of the dome collapse and pyroclastic flow [4].

Scientists are therefore agreeing that rainfall plays a role on volcanic dome collapses, but rainfall data measurement frequency, location of raingages and other limitations inherent to volcanoes make these correlations still difficult to quantify. For this purpose, accurate precipitation measurements are needed to predict dome fluctuations, and even with such dataset, several limitations persist, such as the difficulty to account for local convective precipitation systems near the top of volcanoes [5]. In fact, precipitation measurements at Soufriere volcano [3] and Unzen [4] were made at distances of about 5.7 km from the lava dome and 5 km from the crater, respectively, and precipitation falling directly on the dome was not accurately measured. Therefore, there is a research gap in the quantification of relations between dome movement and rainfalls for scientific and technical reasons.

Therefore, in the present study, high-resolution dome movement was recorded using ground-based radar, and the precipitation directly over the lava dome of Fugendake were collected using the Japanese rainfall radar data (XRAIN) operated by the Ministry of Land, Infrastructure, Transport and Tourism, with the aim to quantify the relationship between rainfalls and dome movement.

2. Research location and Method

To reach this aim, we worked from the lava dome of Mt. Unzen-Fugendake, located on the Shimabara Peninsula in the Kyushu region, southwest of the Japanese archipelago (Fig. 1).




Figure 1. Regional Map of the research area (a) map of Japan with the survey region squared in red, (b)Unzen-Fugendake in Shimabara Peninsula located west of Kumamoto City.

Unzen Fugendake is mainly composed of andesite and dacites (SiO₂ 58-68%) [6]. It has grown in its own graben separated by two parallel faults roughly oriented East West in Shimabara Peninsula (Fig. 1). The volcano has erupted frequently during the historical period, the oldest recorded eruption being in 1663, when basaltic andesite lava containing olivine flowed down a slope about 1 km from the crater. In 1792, dacite-type lava erupted and flowed 2 km to the north. A few months after this eruption, Mayu-yama collapsed, causing a landslide that devastated Shimabara City and further entered the sea, hitting the opposite shore of Shimabara Bay and generating a massive tsunami, taking the lives of 15,000 people [6][7]. The 1991-1995 eruption formed a lava dome on the east side of Fugendake. Pyroclastic-flows and lahars frequently swept this area, and the Mizunashi River (Yamamizu Valley and Jigoku Valley) to the south of the lava dome (Fig. 2).

The displacement of the lava dome is minute and to account for these small variations, we have used a ground-based synthetic aperture radar (GBSAR). It has the further advantage of being able to measure areas of relatively large displacement and not requiring the installation of equipment in inaccessible locations. In the present study, we used the GBSAR at the Akamatsu Valley Observatory operated by Pasco Corporation (Fig. 2). As the dome of Mt. Unzen is recognized to be generated from several lobes, the observation system was calibrated to observe them (as attested by the radar swath range – Fig. 2(a)). Therefore, we have analyzed the data separately for each of the five domes, using dataset collected with a return period of 48 hours. For the present research, we were provided with the data from June 2, 2018 to December 31, 2020. The reference for displacement here is the position of the dome 48 hours ago, respectively. In this paper, this position is defined as 0 mm, and the displacement is defined as the distance moved during the 48 hours.




Figure 2. (a) Location of GBSAR, the area between the orange lines is the observation range, and the lava dome is in the red frame. (b) Block division of the lava dome data used in this study.

At Unzen Volcano, as it is the case of all active volcanoes, instrumentation is always a trade-off between ideal position and safety and practical aspects, often leading to data gaps. This issue is particularly acute with rainfalls as complex topography lead to numerous local effects.

In this study, we used an X-band MP radar-based rainfall observation system called XRAIN (eXtended RAdar Information Network) operated by the Ministry of Land, Infrastructure, Transport and Tourism. This system combines the high-resolution features of X-band radar with the high real-time performance of MP-band radar, enabling observations at higher frequencies (5 times more frequently than conventional radar at 1-minute intervals) and higher resolution (16 times more frequently than conventional radar at 250-meter intervals) than conventional C-band radar. This rainfall observation system was applied to an area of approximately 1,000 m square directly above the lava dome (red frame in Fig. 3), enabling measurement of localized rainfall at the summit of the mountain.

The XRAIN precipitation data used in this study is hourly precipitation data. In addition, each cycle contains precipitation data for a total of 16 areas: 4 areas from east to west and 4 areas from north to south. The data for these 16 areas are then averaged to calculate the hourly precipitation for the area circled by the red box in Fig. 3. In addition, since the lava dome displacement data is for two days, the hourly average precipitation data for the two days (48 hours) is added to calculate the precipitation for the two days.




Figure 3. (a) map of Shimabara peninsula with the Unzen-Fugendake region squared in red, (b) Map of lava dome with XRAIN observation area circled in red

3. Results

3.1. Overall displacement

The 48 hours displacement for each dome ranges from ~8 millimeters to ~51 millimeters for the period June 2, 2018 to December 31st, 2020, with domes 1 and 2 moving twice to 6 times faster than the domes 3 to 5. This movement is the result of a total rainfall > 9.4 thousands millimeters. These values can be divided between downward movement and upward movement. Domes 1 to 3 are moving ‘upward’ (positive values) while dome 4 and 5 are moving ‘downward’ (negative values in Table 1). During this period, the total displacements of Dome 1 to 3 are 49.581 mm, 51.226 mm, and 16.410 mm, respectively, indicating that the dome descended the slope, whereas Dome 4 and 5 are -8.454 mm and -24.236 mm, respectively, indicating an upward slope movement (Table 1). This equates

to an average velocity of 0.0529 mm/day to -0.009 mm/day for the two extreme values provided by dome 1 and dome 4.

Table 1. Total displacement for Dome 1 to Dome 5 for the period June 2, 2018 to December 31, 2020.

	Dome 1	Dome 2	Dome 3	Dome 4	Dome 5
Total displacement (mm)	49.581	51.226	16.410	-8.454	-24.236
Total precipitation (mm)	9428.709				

3.2. Temporal Distribution of displacement and apparent relations with rainy events

During the study-period, four distinct sets of events with rainfalls exceeding 150 mm/48 hours were recorded with three of these events having recurring peaks within a couple of weeks periods (Fig. 4). Event 1 (Ev1) occurs in June 2018 with at least two peaks of rainfalls exceeding 150 mm (arbitrary thresholds used for the present study, further work on its significance is necessary). This event has created for dome 3, 4 and 5 visible increase in the GBSAR data with sets of pronounced peaks of dome movement. The lower domes have shown the highest change in amplitude, and the lower the dome the longer the sequence of change is lasting. At dome number 5, the change lasted for almost 6 months. Event number 2 occurs in July-August 2019 and similarly it presents visible excitation in the signal for domes 3 to 5 with the highest amplitude and duration for the lower dome. For Event three, change in all the domes from 1 to 5 occurs almost at the same time as the rainfall, and although there are no clear observations of these changes at dome 1 and dome 2, domes 3 to 5 display important movement until the end of 2020. Another >150 mm rainfall event was also detected at the end of March 2020, but it is isolated with less “background rainfall” compared to the other events 1 to 3 and it is difficult to know whether it had an impact on the dome as it does not clearly show in the signal. This result shows that it is the combination of high-rainfall volume concentrated over a few days combined to “background” rainy events that are best linked to displacement of the dome.




Figure 4. Dome displacement and rainfall events plotted over time during the study period. The y-axis of Dome1-5 is the displacement (mm) and the y-axis of precipitation is the 48-hour precipitation (mm).

3.3. Upward and Downward displacement

Combining the displacement and the rainfall data for every 48 hours periods confirm the data provided temporally (Fig. 5). Combined with the correlation coefficients in Table 2, both upward and downward displacement are not correlated with the amount of rainfall at the time of displacement, but the displacement tends to be smaller with more rainfall and larger with less rainfall. This indicates that precipitation for a short period of time

(48 hours) alone is unlikely to trigger the displacement of the dome. This result further emphasizes the importance of background rainfall, which was shown earlier in 3.2.

Focusing only on dome rise, the displacement of domes 1 and 2, which are located at the top of the slope, is within approximately 2mm, while the displacement of domes 3, 4, and 5, which are located at the bottom of the slope, is within approximately 4mm, indicating that domes 3, 4, and 5, which are located at the bottom of the slope, tend to rise in a short period of time (Figure 5(a),(b)). In contrast, all the domes have a similar scattering pattern in terms of descent, indicating that there is no significant difference in descent in 48 hours regardless of the slope location.

Table 2. Correlation coefficient between rainfall and displacement (upward/downward).

For downwards displacement					
	Dome 1	Dome 2	Dome 3	Dome 4	Dome 5
correlation coefficient	0.047	0.036	0.056	-0.049	-0.005
For upwards displacement					
	Dome 1	Dome 2	Dome 3	Dome 4	Dome 5
correlation coefficient	0.124	0.099	0.116	0.054	0.077




Fig. 5. Scatterplot of Figure 4 further subdivided by dome location: (a)Dome 1,2(upward) , (b)Dome 3,4,5 (upward), (c)Dome 1,2 (downward), (d)Dome 3,4,5 (downward).

4. Discussion

The comparison of rainfall events and dome displacement have shown that the dome is moving at a different rhythm in its upper area, compared to the lower area, with vertical changes in the down part that is almost twice the change in the upper part. This change is however not immediate after heavy rainfall events and they can also last for months after a rainfall event where peaks exceed 150 mm/48 hours. We have also seen that the rainfall peaks that are associated with background rainfall are more effective in triggering dome movement.

The present movement is thus showing a time-delay, which is then not a similar process to the one of dome collapse [4], as it was pointed out that the relation between “hot” dome collapse generating pyroclastic density currents and rainfall was almost immediate. Carn et al. (2004) showed that rainfall and collapse responded quickly to the collapse of the lava dome at Soufriere volcano on July 29, 2001, but this result is different from the Unzen displacement data.

As explained in Kelfoun et al. (2021) four causes of lava dome collapse and thus movement can be inferred: (1) excessive steepening of the lava dome due to gravity, (2) dome failure due to volcanic gas pressure, (3) precipitation, (4) the structure of the dome and its lower layer, but the slow response of precipitation and displacement in the Unzen lava dome suggests that the displacement of the Unzen lava dome may have been caused by a combination of (1), (3), and (4), especially because the dome position on the slope seems to define the amount of movement recorded.

This result is therefore essential for its use in hazard and disaster risk research because it shows that volcanic domes during quiescent time are showing a different reaction time to rainfall events and that in critical case it would be important to extend temporally the alert time to account for the slow response of the dome.

5. Conclusion

In conclusion, the dome of Unzen-Fugendake is reacting to a complex of high-intensity and background rainfall, and so over several month periods, suggesting that the vadose zone and the groundwater movement may play an important controlling role. Furthermore, this time-lag suggests that hazards and disaster risk alert need to be adapted to account for those delayed changes. The next step in this research will now be to model the time-lag of the dome to rainfall events.

Acknowledgments: In this study, XRAIN precipitation data were downloaded from DIAS. In addition, GBSAR data was provided by the Kyushu Regional Development Bureau, Ministry of Land, Infrastructure, Transport and Tourism, Unzen Sediment Control Management Center and Pasco Corporation.

References

1. Herd, R. A.; Edmonds, M.; Bass, V. A. Catastrophic Lava Dome Failure at Soufrière Hills Volcano, Montserrat, 12–13 July 2003. *Journal of Volcanology and Geothermal Research* **2005**, *148* (3), 234–252. <https://doi.org/10.1016/j.jvolgeores.2005.05.003>.
2. Kelfoun, K.; Santoso, A. B.; Latchimy, T.; Bontemps, M.; Nurdien, I.; Beauducel, F.; Fahmi, A.; Putra, R.; Dahamna, N.; Laurin, A.; Rizal, M. H.; Sukmana, J. T.; Gueugneau, V. Growth and Collapse of the 2018–2019 Lava Dome of Merapi Volcano. *Bull Volcanol* **2021**, *83* (2), 8. <https://doi.org/10.1007/s00445-020-01428-x>.
3. Carn, S. A.; Watts, R. B.; Thompson, G.; Norton, G. E. Anatomy of a Lava Dome Collapse: The 20 March 2000 Event at Soufrière Hills Volcano, Montserrat. *Journal of Volcanology and Geothermal Research* **2004**, *131* (3), 241–264. [https://doi.org/10.1016/S0377-0273\(03\)00364-0](https://doi.org/10.1016/S0377-0273(03)00364-0).
4. Yamasato, H.; Kitagawa, S.; Komiya, M. Effect of Rainfall on Dacitic Lava Dome Collapse at Unzen Volcano, Japan. *Pap. Met. Geophys.* **1998**, *48* (3), 73–78. <https://doi.org/10.2467/mripapers.48.73>.
5. Matthews, A. J.; Barclay, J.; Carn, S.; Thompson, G.; Alexander, J.; Herd, R.; Williams, C. Rainfall-Induced Volcanic Activity on Montserrat. *Geophysical Research Letters* **2002**, *29* (13), 22-1-22–24. <https://doi.org/10.1029/2002GL014863>.
6. Otani Yasuo, Tsuchide Masakazu, Shibata Atsushi, Kato Shigeru, Iwabuchi Yoshio. (2004) The List of Volcanoes and Their Activities Records in the Adjacent Seas of Japan 3rd Edition. Report of Hydrographic and Oceanographic Reserchies No.40 March, 2004.
7. Hideo Hoshizumi, Kozo Uto, Kazunori Watanabe. Geology and eruptive history of Unzen volcano, Shimabara Peninsula, Kyushu, SW Japan. 1999. *Journal of Volcanology and Geothermal Research* Volume 89, Issues 1–4, April 1999, pp.81-94.